



Extremal RN/CFT in both hands revisited

En-Jui Kuo, Yi Yang*

Department of Electrophysics, National Chiao-Tung University, Hsinchu, Taiwan, ROC



ARTICLE INFO

Article history:

Received 11 December 2015
 Received in revised form 16 January 2016
 Accepted 9 February 2016
 Available online 10 February 2016
 Editor: M. Cvetič

ABSTRACT

We study RN/CFT correspondence for four dimensional extremal Reissner–Nordstrom black hole. We uplift the 4d RN black hole to a 5d rotating black hole and make a geometric regularization of the 5d space–time. Both hands central charges are obtained correctly at the same time by Brown–Henneaux technique.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

To fully understand Bekenstein–Hawking entropy of a black hole in a microscopic point of view is still a challenge. An important progress has been made in [1] by using the Brown–Henneaux technique [2], i.e. the Kerr/CFT correspondence. The central charge of the dual CFT reproduces the exact Bekenstein–Hawking entropy of the 4-dimensional extremal Kerr black hole by Cardy's formula. This method has been extensively studied for Kerr black hole and other more general rotating black holes [3–16]. However, in the original Kerr/CFT method, the Virasoro algebra was realized only from an enhancement of the rotational $U(1)$ isometry, which corresponds to left hand central charge, not from the $SL(2, R)$. Later, it was found that the right hand central charge of rotating black holes can be obtained by delicately choosing appropriate boundary conditions, [17–28].

Nevertheless, the method in Kerr/CFT correspondence cannot be directly applied to non-rotating charge black holes, such as Reissner–Nordstrom (RN) black hole. One found that the central charge for 4d RN black hole vanishes by directly using the Brown–Henneaux technique. One possible way to solve the problem is to uplift the space–time to higher dimensions. The left hand central charge $c_L = 6Q^2$ of 4d extremal RN black hole was obtained by uplifting it to 5 dimensions [4,9,24,29–31]. The other possible way is to consider a 2-dimensional effective theory of 4d extremal RN black hole via a dimensional reduction following the idea in [19], by which the right hand central charge $c_R = 6Q^2$ of 4d extremal RN was obtained in [31,32]. Although both hands central charges can be obtained, but one has to use difference method. It

is an interesting question whether we are able to calculate both hands central charges for extremal RN black hole at the same time by choosing appropriate boundary conditions as in the case of Kerr/CFT. The answer is yes, but with a geometric regularization.

In this paper, we calculate both hands central charges for 4d extremal RN black hole by using the Brown–Henneaux technique. We first uplift the 4d RN black hole to a 5d black hole, following the idea in [27], we then deform the 5d black hole by a geometric regularization. Both hands central charges $c_R = c_L = 6Q^2$ are obtained in this way at the same time.

The paper is organized as follows. In section 2, we briefly review the near horizon geometry of 4d extremal RN black hole. We then uplift 4d RN black hole to 5d and calculate both hands asymptotic Killing vectors in section 3. In section 4, we deform the 5d black hole by the geometric regularization to obtain both hands central charges. Our conclusion and discussion are included in section 5.

2. Near horizon geometry of 4d extremal RN black hole

In four dimensions, the Einstein–Maxwell theory

$$S_4 = \frac{1}{16\pi G_4} \int d^4x \sqrt{-g_4} (R_4 - F^2), \quad (1)$$

admits a unique spherical electro-vacuum solution, the RN black hole. The metric of 4d non-extremal RN black hole is

$$ds^2 = -f(r) dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega_2^2, \quad (2)$$

where $d\Omega_2^2 = d\theta^2 + \sin^2\theta d\phi^2$ and

$$f(r) = 1 - \frac{2m}{r} + \frac{q^2}{r^2}, \quad A = -\frac{q}{r} dt. \quad (3)$$

* Corresponding author.

E-mail addresses: eric1028.ep01@nctu.edu.tw (E.-J. Kuo), yyang@mail.nctu.edu.tw (Y. Yang).

Two horizons are obtained by solving $f(r) = 0$,

$$r_{\pm} = m \pm \sqrt{m^2 - q^2}. \quad (4)$$

The Hawking temperature and black hole entropy can be calculated as follows,

$$T_H = \frac{\kappa}{2\pi} = \frac{r_+ - r_-}{4\pi r_+^2}, \quad S_{BH} = \frac{A}{4G_4} = \frac{\pi r_+^2}{G_4}. \quad (5)$$

In the extremal limit of RN black hole, i.e. $r_+ = r_- = m = q$, we have

$$f(r) = \left(1 - \frac{q}{r}\right)^2, \quad (6)$$

and

$$T_H = 0, \quad S_{BH} = \frac{\pi q^2}{G_4} = \pi Q^2, \quad (7)$$

where we defined $Q \equiv q/\sqrt{G_4}$.

To consider the near horizon limit, $r \rightarrow r_+$, we make the following coordinates transformation,

$$\rho = \frac{r - r_+}{\epsilon q}, \quad \tau = \frac{t}{q}, \quad (8)$$

and take the limit $\epsilon \rightarrow 0$. Finally, the metric of 4d near horizon extremal RN black hole is obtained as

$$ds^2 = q^2 \left(-\rho^2 d\tau^2 + \frac{d\rho^2}{\rho^2} + d\Omega_2^2 \right), \quad (9)$$

$$A = q\rho d\tau. \quad (10)$$

3. Central charges by uplifting to five dimensions

Direct calculation following the method of Kerr/CFT in [1] shows that the central charges of the 4d extremal RN black hole always vanish. This is because the $U(1)$ symmetry of rotation in Kerr black hole does not exist in the RN black hole, in which the $U(1)$ symmetry behaves as gauge symmetry. To get non-zero central charge, a possible way is to uplift the 4d RN black hole into 5-dimensional space-time with the extra dimension compactified on a circle as in [4,31]. The metric of the uplifted 5-dimensional black hole can be expressed as,

$$ds^2 = q^2 \left(-\rho^2 d\tau^2 + \frac{d\rho^2}{\rho^2} + d\Omega_2^2 \right) + (dy + A)^2 \\ = \frac{q^2 d\rho^2}{\rho^2} + q^2 d\Omega_2^2 + dy^2 + 2q\rho d\tau dy, \quad (11)$$

$$A_{(2)} = \sqrt{3}q\rho d\tau \wedge dy, \quad (12)$$

where $A_{(2)}$ is a gauge 2-form and the new coordinate y is compactified on a circle with a proper period [30] $y = y + 2\pi q$. The metric (11) with the 2-form (12) is a solution of 5d Einstein-Maxwell theory,

$$S_5 = \frac{1}{16\pi G_5} \int d^4x \sqrt{-g_5} \left(R_5 - \frac{1}{12} F_{(3)}^2 \right), \quad (13)$$

where $G_5 = 2\pi q G_4$ and $F_{(3)} = dA_{(2)}$. We should notice that the 2-form field $A_{(2)}$ has no contribution to the central charge based on the argument in [6,29].

To get the expected leading behavior of asymptotic Killing vector (AKV) with both hands, we carefully choose the following boundary condition,

$$\begin{pmatrix} h_{\tau\tau} = \mathcal{O}(1) & h_{\tau\rho} = \mathcal{O}(\rho^{-2}) & h_{\tau\theta} = \mathcal{O}(\rho^{-1}) & h_{\tau\phi} = \mathcal{O}(\rho^{-1}) & h_{\tau y} = \mathcal{O}(1) \\ & h_{\rho\rho} = \mathcal{O}(\rho^{-3}) & h_{\rho\theta} = \mathcal{O}(\rho^{-2}) & h_{\rho\phi} = \mathcal{O}(\rho^{-2}) & h_{\rho y} = \mathcal{O}(\rho^{-1}) \\ & & h_{\theta\theta} = \mathcal{O}(\rho^{-1}) & h_{\theta\phi} = \mathcal{O}(\rho^{-1}) & h_{\theta y} = \mathcal{O}(\rho^{-1}) \\ & & & h_{\phi\phi} = \mathcal{O}(\rho^{-1}) & h_{\phi y} = \mathcal{O}(\rho^{-1}) \\ & & & & h_{yy} = \mathcal{O}(1) \end{pmatrix}. \quad (14)$$

The AKV is obtained by solving the asymptotic Killing equation $L_{\zeta}(g_{\mu\nu} + h_{\mu\nu}) \sim h_{\mu\nu}$ with the corresponding boundary condition. We note that a weaker boundary condition will cause the AKV being ill-defined and further restriction will rule out the interesting excitations. The AKV associated to the above boundary condition (14) is

$$\zeta = \epsilon(\tau) \partial_{\tau} + [-\rho \partial_{\tau} \epsilon(\tau) - \rho \partial_y \eta(y)] \partial_{\rho} \\ + [\eta(y) - q \partial_{\tau}^2 \epsilon(\tau) / \rho] \partial_y, \quad (15)$$

where $\epsilon(\tau)$ and $\eta(y)$ are arbitrary functions of τ and y , respectively. The left hand and right hand AKVs read

$$\zeta^L = -\rho \partial_y \eta(y) \partial_{\rho} + \eta(y) \partial_y, \quad (16a)$$

$$\zeta^R = \epsilon(\tau) \partial_{\tau} - \rho \partial_{\tau} \epsilon(\tau) \partial_{\rho} - \frac{q}{\rho} \partial_{\tau}^2 \epsilon(\tau) \partial_y. \quad (16b)$$

Since $y = y + 2\pi q$ is periodic, we can express the function $\eta(y)$ as its Fourier bases $e^{-iny/q}$. Similarly, if we assign a period for τ with $\tau = \tau + 2\pi\beta$ with β an arbitrary real number, the function $\epsilon(\tau)$ can be expressed by $e^{-in\tau/\beta}$. With appropriate normalizations, we can write

$$\eta_n(y) = -q e^{-in\frac{y}{q}}, \quad \epsilon_n(\tau) = -\beta e^{-in\frac{\tau}{\beta}}. \quad (17)$$

The corresponding AKVs

$$\zeta_n^L = -in\rho e^{-in\frac{y}{q}} \partial_{\rho} - q e^{-in\frac{y}{q}} \partial_y, \quad (18a)$$

$$\zeta_n^R = -\beta e^{-in\frac{\tau}{\beta}} \partial_{\tau} - in\rho e^{-in\frac{\tau}{\beta}} \partial_{\rho} - \frac{n^2 q}{\rho\beta} e^{-in\frac{\tau}{\beta}} \partial_y, \quad (18b)$$

compose two copies of Virasoro algebra without central extension as expected,

$$i \left[\zeta_m^L, \zeta_n^L \right] = (m - n) \zeta_{m+n}^L, \quad (19a)$$

$$i \left[\zeta_m^R, \zeta_n^R \right] = (m - n) \zeta_{m+n}^R. \quad (19b)$$

To obtain the central charges, we define the asymptotic charge Q_{ζ} associated to the AKV ζ as,

$$Q_{\zeta} = \frac{1}{8\pi G_5} \int_{\partial\Sigma} k_{\zeta} [h, g], \quad (20)$$

where the 2-form k_{ζ} is defined for a perturbation $h_{\mu\nu}$ around the background metric $g_{\mu\nu}$ [3],

$$k_{\zeta} [h, g] = \frac{1}{2} \left[\zeta_{\nu} \nabla_{\mu} h - \zeta_{\nu} \nabla_{\sigma} h_{\mu}^{\sigma} + \zeta_{\sigma} \nabla_{\nu} h_{\mu}^{\sigma} \right. \\ \left. + \frac{1}{2} h \nabla_{\nu} \zeta_{\mu} - h_{\nu}^{\sigma} \nabla_{\sigma} \zeta_{\mu} \right. \\ \left. + \frac{1}{2} h_{\nu\sigma} (\nabla_{\mu} \zeta^{\sigma} + \nabla^{\sigma} \zeta_{\mu}) \right] * (dx^{\mu} \wedge dx^{\nu}), \quad (21)$$

where $*$ denotes the Hodge dual, and Σ is the 4-dimensional equal-time hypersurface at $\tau = \text{const}$.

The Dirac bracket of the asymptotic charge is given by

$$\{Q_{\zeta_m}, Q_{\zeta_n}\}_{DB} = Q_{[\zeta_m, \zeta_n]} + \frac{1}{8\pi G_5} \int_{\partial\Sigma} k_{\zeta_m} [L_{\zeta_n} \bar{g}, \bar{g}]. \quad (22)$$

We define the generators L_n of Virasoro algebra as

$$L_n \equiv Q_{\zeta_n} + \alpha \delta_n,$$

with α an arbitrary c -number. With the usual rule of Dirac brackets $\{Q_{\zeta_m}, Q_{\zeta_n}\}_{DB} \rightarrow -i[Q_{\zeta_m}, Q_{\zeta_n}]$, the commutation relations of the generators L_n can be calculated as

$$\begin{aligned} [L_m, L_n] &= i \{Q_{\zeta_m}, Q_{\zeta_n}\}_{DB} \\ &= (m-n)L_{m+n} + \frac{i}{8\pi G_5} \int_{\partial\Sigma} k_{\zeta_m} [\mathcal{L}_{\zeta_n} \bar{g}, \bar{g}], \end{aligned} \quad (23)$$

where \mathcal{L} is Lie derivative. Plugging AKVs (18) into the 2-form (21), we obtain

$$\begin{aligned} \frac{i}{8\pi G_5} \int_{\partial\Sigma} k_{\zeta_m}^L [\mathcal{L}_{\zeta_n} \bar{g}, \bar{g}] &= \frac{q^2}{2G_4} (m^3 + m) \delta_{m+n} \\ &= \frac{Q^2}{2} (m^3 + m) \delta_{m+n}, \end{aligned} \quad (24a)$$

$$\frac{i}{8\pi G_5} \int_{\partial\Sigma} k_{\zeta_m}^R [\mathcal{L}_{\zeta_n} \bar{g}, \bar{g}] = 0. \quad (24b)$$

Comparing to the standard commutation relations of Virasoro algebra,

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{c}{12} (m^3 - m) \delta_{m+n}, \quad (25)$$

the central charges are read as,¹

$$c_L = 6Q^2, \quad c_R = 0. \quad (26)$$

We see that the above calculation only gives us the correct left hand central charge with the right hand central charge vanishing. However, because there is no gravitational anomaly, we expect that $c_L = c_R$. In next section, we will make a geometric regularization of the 5d black hole and obtain the correct both hands central charges together.

4. Geometric regularization

At the boundary $\rho = \text{const} \rightarrow \infty$, the 5d metric (11) reduces to

$$ds^2 = \frac{q^2 d\rho^2}{\rho^2} + q^2 d\Omega_2^2 + dy^2 + 2q\rho d\tau dy \rightarrow 2q\rho d\tau dy. \quad (27)$$

In the above calculation, we took the “equal-time hypersurface” Σ with $\tau = \text{const}$ when we defined the asymptotic charge Q_ζ in Eq. (20). However, it is easy to see that $ds^2 \rightarrow 0$ for $\tau = \text{const}$ from Eq. (27), so that Σ is not a space-like surface but a light-like one on the boundary. Therefore our naive choice of the “equal-time hypersurface” Σ causes problem when we consider the asymptotic behavior near the boundary. To define a space-like surface at the boundary, we follow [27] to regularize the 5d geometry by a coordinates transformation,

$$\tau' = \tau - ay, \quad y' = y, \quad (28)$$

where a is the regularization parameter. We should remark that the resulted central charges should be independent of the regularization parameter a .

With the coordinates transformation (28), the 5d black hole metric (11) becomes

$$ds^2 = \frac{q^2 d\rho^2}{\rho^2} + q^2 d\Omega_2^2 + (1 + 2aq\rho) dy'^2 + 2q\rho d\tau' dy', \quad (29)$$

which, at the boundary $\rho = \text{const} \rightarrow \infty$, reduces to

$$ds^2 \rightarrow 2aq\rho dy'^2 + 2q\rho d\tau' dy'. \quad (30)$$

Now the shifted “equal-time hypersurface” Σ' with $\tau' = \text{const}$ becomes a space-like surface $ds^2 \rightarrow 2aq\rho dy'^2$.

We also note that, to respect the periodicity $y = y + 2\pi q$, we should fix the period of τ by $\beta = aq$ in Eq. (17) and the AKVs can be rewritten by the new coordinates (τ', y') as

$$\xi_n^L = aqe^{-in\frac{y'}{q}} \partial_{\tau'} - in\rho e^{-in\frac{y'}{q}} \partial_\rho - qe^{-in\frac{y'}{q}} \partial_{y'}, \quad (31a)$$

$$\begin{aligned} \xi_n^R &= -\left(aq + \frac{n^2}{\rho}\right) e^{-in\left(\frac{\tau'}{a} + y'\right)/q} \partial_{\tau'} - in\rho e^{-in\left(\frac{\tau'}{a} + y'\right)/q} \partial_\rho \\ &\quad - \frac{n^2}{a\rho} e^{-in\left(\frac{\tau'}{a} + y'\right)/q} \partial_{y'}, \end{aligned} \quad (31b)$$

which also satisfy Virasoro algebra without central charge as in Eq. (19). Similar calculations can be carried on with the results,

$$\begin{aligned} \frac{i}{8\pi G_5} \int_{\partial\Sigma} k_{\zeta_m}^L [L_{\zeta_n} \bar{g}, \bar{g}] &= \frac{q^2}{2G_4} (m^3 + m) \delta_{m+n} \\ &= \frac{Q^2}{2} (m^3 + m) \delta_{m+n}, \end{aligned} \quad (32a)$$

$$\frac{i}{8\pi G_5} \int_{\partial\Sigma} k_{\zeta_m}^R [L_{\zeta_n} \bar{g}, \bar{g}] = \frac{q^2}{2G_4} m^3 \delta_{m+n} = \frac{Q^2}{2} m^3 \delta_{m+n}. \quad (32b)$$

Comparing to the standard commutation relations of Virasoro algebra (25), the above results lead to the correct both hands central charges of 4d extremal RN black hole,

$$c_L = c_R = 6Q^2. \quad (33)$$

We see that the central charges are indeed independent of the regularization parameter a as we promised.

Together with the CFT temperature of both hands [31,32],

$$T_L = \frac{1}{2\pi}, \quad T_R = 0, \quad (34)$$

we obtain the microscopic entropy by Cardy's formula from the central charges (33),

$$S = \frac{\pi^2}{3} (c_L T_L + c_R T_R) = \pi Q^2, \quad (35)$$

which correctly reproduces the Bekenstein–Hawking entropy of the extremal RN black hole (7).

5. Conclusion

In this paper, we studied near horizon geometry of 4d extremal RN black hole. We uplifted the 4d extremal RN black hole to a 5d black hole, and deformed the 5d black hole by a geometric regularization. Both hands central charges of 4d extremal RN black hole $c_L = c_R = 6Q^2$ were correctly obtained by using the Brown–Henneaux technique. The crucial step to get the result is the geometric regularization by which the shifted “equal-time hypersurface” becomes space-like. The resulted central charges are independent of the regularization as we expected.

¹ The term linear in m plays no essential role here, since it can be shifted arbitrarily by the c -number α in the definition of the generators L_n .

Acknowledgements

We thank Chiang-Mei Chen, Jia-Rui Sun and Xiaoning Wu for useful discussions. This work is supported by the National Science Council (NSC 101-2112-M-009-005) and National Center for Theoretical Sciences, Taiwan.

References

- [1] Monica Guica, Thomas Hartman, Wei Song, Andrew Strominger, The Kerr/CFT correspondence, arXiv:0809.4266, 2008.
- [2] J. David Brown, Marc Henneaux, Central charges in the canonical realization of asymptotic symmetries: an example from three dimensional gravity, *Commun. Math. Phys.* 104 (2) (1986) 207–226.
- [3] H. Lü, Jianwei Mei, C.N. Pope, Kerr–AdS/CFT correspondence in diverse dimensions, *J. High Energy Phys.* 2009 (04) (2009) 054.
- [4] Thomas Hartman, Keiju Murata, Tatsuma Nishioka, Andrew Strominger, CFT duals for extreme black holes, *J. High Energy Phys.* 2009 (04) (2009) 019.
- [5] David D.K. Chow, Mirjam Cvetič, H. Lü, C.N. Pope, Extremal black hole/CFT correspondence in (gauged) supergravities, *Phys. Rev. D* 79 (8) (2009) 084018.
- [6] Geoffrey Compère, Keiju Murata, Tatsuma Nishioka, Central charges in extreme black hole/CFT correspondence, *J. High Energy Phys.* 2009 (05) (2009) 077.
- [7] Wen-Yu Wen, Holographic descriptions of (near-) extremal black holes in five dimensional minimal supergravity, arXiv:0903.4030, 2009.
- [8] Steven Carlip, Extremal and nonextremal Kerr/CFT correspondences, *J. High Energy Phys.* 2011 (4) (2011) 1–17.
- [9] Bin Chen, Jia-ju Zhang, Novel CFT duals for extreme black holes, *Nucl. Phys. B* 856 (2) (2012) 449–474.
- [10] Jianwei Mei, On the general Kerr/CFT correspondence in arbitrary dimensions, *J. High Energy Phys.* 2012 (4) (2012) 1–18.
- [11] Bin Chen, Jia-ju Zhang, RN/CFT correspondence from thermodynamics, *J. High Energy Phys.* 2013 (1) (2013) 1–30.
- [12] Cheng-Yong Zhang, Yu Tian, Xiao-Ning Wu, Generalized Kerr/CFT correspondence with electromagnetic field, *Class. Quantum Gravity* 31 (8) (2014) 085009.
- [13] Cheng-Yong Zhang, Yu Tian, Xiao-Ning Wu, Shao-Jun Zhang, Entropy of the isolated horizon from the surface term of gravitational action, *Class. Quantum Gravity* 31 (19) (2014) 195002.
- [14] Avirup Ghosh, Note on Kerr/CFT correspondence in a first order formalism, *Phys. Rev. D* 89 (12) (2014) 124035.
- [15] G. Compère, K. Hajian, A. Seraj, M.M. Sheikh-Jabbari, Extremal rotating black holes in the near-horizon limit: phase space and symmetry algebra, arXiv:1503.07861, 2015.
- [16] Marco Astorino, Magnetised Kerr/CFT correspondence, *Phys. Lett. B* 751 (2015) 96–106.
- [17] Yoshinori Matsuo, Takuya Tsukioka, Chul-Moon Yoo, Another realization of Kerr/CFT correspondence, *Nucl. Phys. B* 825 (1) (2010) 231–241.
- [18] Yoshinori Matsuo, Takuya Tsukioka, Chul-Moon Yoo, Yet another realization of Kerr/CFT correspondence, *Europhys. Lett.* 89 (6) (2010) 60001.
- [19] Alejandra Castro, Finn Larsen, Near extremal Kerr entropy from AdS2 quantum gravity, *J. High Energy Phys.* 2009 (12) (2009) 037.
- [20] Jørgen Rasmussen, Isometry-preserving boundary conditions in the Kerr/CFT correspondence, *Int. J. Mod. Phys. A* 25 (08) (2010) 1597–1613.
- [21] Jørgen Rasmussen, A near-NHEK/CFT correspondence, *Int. J. Mod. Phys. A* 25 (30) (2010) 5517–5527.
- [22] Jørgen Rasmussen, On the CFT duals for near-extremal black holes, *Mod. Phys. Lett. A* 26 (22) (2011) 1601–1611.
- [23] Monica Guica, Andrew Strominger, Microscopic realization of the Kerr/CFT correspondence, *J. High Energy Phys.* 2011 (2) (2011) 1–20.
- [24] Bin Chen, Chiang-Mei Chen, Bo Ning, Holographic Q-picture of Kerr–Newman–AdS–dS black hole, *Nucl. Phys. B* 853 (1) (2011) 196–209.
- [25] Tatsuo Azeyanagi, Noriaki Ogawa, Seiji Terashima, Emergent AdS3 in the zero entropy extremal black holes, *J. High Energy Phys.* 2011 (3) (2011) 1–27.
- [26] Yoshinori Matsuo, Tatsuma Nishioka, New near horizon limit in Kerr/CFT, *J. High Energy Phys.* 2010 (12) (2010) 1–24.
- [27] Tatsuo Azeyanagi, Noriaki Ogawa, Seiji Terashima, On non-chiral extension of Kerr/CFT, *J. High Energy Phys.* 2011 (6) (2011) 1–25.
- [28] Chang-Young Ee, Myungseok Eune, Nonextremal Kerr/CFT on a stretched horizon, arXiv:1212.3031, 2012.
- [29] Ahmad Ghodsi, Mohammad R. Garousi, The RN/CFT correspondence, *Phys. Lett. B* 687 (1) (2010) 79–83.
- [30] Chiang-Mei Chen, Ying-Ming Huang, Shou-Jyun Zou, Holographic duals of near-extremal Reissner–Nordström black holes, *J. High Energy Phys.* 2010 (3) (2010) 1–14.
- [31] Chiang-Mei Chen, Jia-Rui Sun, Holographic dual of the Reissner–Nordström black hole, *J. Phys. Conf. Ser.* 330 (2011) 012009, IOP Publishing.
- [32] Chiang-Mei Chen, Jia-Rui Sun, Shou-Jyun Zou, The RN/CFT correspondence revisited, *J. High Energy Phys.* 2010 (1) (2010) 1–13.