

# The Mutually Independent Bipanconnected Property for Hypercube

Yuan-Kang Shih

Department of Computer Science,  
National Chiao Tung University,  
Hsinchu, Taiwan 30010, R.O.C.  
Email: ykshih@cs.nctu.edu.tw

Jimmy J. M. Tan

Department of Computer Science,  
National Chiao Tung University,  
Hsinchu, Taiwan 30010, R.O.C.  
Email: jmtan@cs.nctu.edu.tw

Lih-Hsing Hsu

Department of Computer Science  
and Information Engineering  
Providence University,  
Taichung, Taiwan 43301, R.O.C.  
Email: lhhsu@cs.pu.edu.tw

**Abstract**—A graph is denoted by  $G$  with the vertex set  $V(G)$  and the edge set  $E(G)$ . A path  $P = \langle v_0, v_1, \dots, v_m \rangle$  is a sequence of adjacent vertices. Two paths with equal length  $P_1 = \langle u_1, u_2, \dots, u_m \rangle$  and  $P_2 = \langle v_1, v_2, \dots, v_m \rangle$  from  $a$  to  $b$  are independent if  $u_1 = v_1 = a$ ,  $u_m = v_m = b$ , and  $u_i \neq v_i$  for  $2 \leq i \leq m - 1$ . Paths with equal length  $\{P_i\}_{i=1}^n$  from  $a$  to  $b$  are mutually independent if they are pairwise independent. Let  $u$  and  $v$  be two distinct vertices of a bipartite graph  $G$ , and let  $l$  be a positive integer length,  $d_G(u, v) \leq l \leq |V(G) - 1|$  with  $(l - d_G(u, v))$  being even. We say that the pair of vertices  $u, v$  is  $(m, l)$ -mutually independent bipanconnected if there exist  $m$  mutually independent paths  $\{P_i\}_{i=1}^m$  with length  $l$  from  $u$  to  $v$ . In this paper, we explore yet another strong property of the hypercubes. We prove that every pair of vertices  $u$  and  $v$  in the  $n$ -dimensional hypercube, with  $d_{Q_n}(u, v) \geq n - 1$ , is  $(n - 1, l)$ -mutually independent bipanconnected for every  $l$ ,  $d_{Q_n}(u, v) \leq l \leq |V(Q_n) - 1|$  with  $(l - d_{Q_n}(u, v))$  being even. As for  $d_{Q_n}(u, v) \leq n - 2$ , it is also  $(n - 1, l)$ -mutually independent bipanconnected if  $l \geq d_{Q_n}(u, v) + 2$ , and is only  $(l, l)$ -mutually independent bipanconnected if  $l = d_{Q_n}(u, v)$ .

## I. INTRODUCTION

For the graph definitions and notations we refer the reader to [6]. A graph is denoted by  $G$  with the vertex set  $V(G)$  and the edge set  $E(G)$ . The simulation of one architecture by another is an important issue in interconnection networks. The problem of simulating one network by another is also called embedding problem. One particular problem of path embedding deals with finding all the possible length of paths in an interconnection network.

A path  $P = \langle v_0, v_1, \dots, v_m \rangle$  is a sequence of adjacent vertices. We also write  $P = \langle v_0, \dots, v_i, Q, v_j, \dots, v_m \rangle$  where  $Q$  is a path  $\langle v_i, \dots, v_j \rangle$ . A cycle  $C = \langle v_0, v_1, \dots, v_m, v_0 \rangle$  is a sequence of adjacent vertices where the first vertex is the same as the last one. The length of a path  $P$  (a cycle  $C$  respectively) is the number of edges in  $P$  (in  $C$  respectively).

A cycle of  $G$  is a *hamiltonian cycle* if it traverses all the vertices exactly once. A graph  $G$  is called a *hamiltonian graph* if  $G$  contains a hamiltonian cycle. A path of  $G$  is a *hamiltonian path* if it contains all the vertices exactly once. A graph  $G$  is *hamiltonian connected* if there exists a hamiltonian path between any two different vertices of  $G$ . A graph  $G = (B \cup W, E)$  is *bipartite* if  $V(G)$  is the union of two disjoint sets  $B$  and  $W$  such that every edge joins  $B$  with  $W$ . It is easy to see that any bipartite graph with at least three vertices is

not hamiltonian connected. A bipartite graph  $G$  is *hamiltonian laceable* if there exists a hamiltonian path joining any two vertices from different partite sets. A graph  $G$  is *pancyclic* [2] if  $G$  includes cycles of all lengths. If these cycles are restricted to even length,  $G$  is called a *bipancyclic graph*. The distance from  $x$  to  $y$ , written  $d_G(x, y)$ , is the least length among all paths from  $x$  to  $y$  in  $G$ . A graph is *panconnected* if, for any two different vertices  $x$  and  $y$ , there exists a path of length  $l$  joining  $x$  and  $y$ , for every  $l$ ,  $d_G(x, y) \leq l \leq |V(G) - 1$ . The concept of panconnected graphs is proposed by Alavi and Williamson [1]. It is not hard to see that any bipartite graph with at least 3 vertices is not panconnected. Therefore, the concept of bipanconnected graphs is proposed. A bipartite graph is *bipanconnected* if, for any two different vertices  $x$  and  $y$ , there exists a path of length  $l$  joining from  $x$  to  $y$ , for every  $l$ ,  $d_G(x, y) \leq l \leq |V(G) - 1$  and  $(l - d_G(x, y))$  being even. There are many studies on bipanconnected graphs and bipancyclic graphs [3], [7], [9], [13].

We introduce some terms defined recently. Two paths  $P_1 = \langle u_1, u_2, \dots, u_m \rangle$  and  $P_2 = \langle v_1, v_2, \dots, v_m \rangle$  from  $a$  to  $b$  are *independent* [10] if  $u_1 = v_1 = a$ ,  $u_m = v_m = b$ , and  $u_i \neq v_i$  for  $2 \leq i \leq m - 1$ . Paths with equal length  $\{P_i\}_{i=1}^n$  from  $a$  to  $b$  are *mutually independent* [10] if they are pairwise independent. Two cycles  $C_1 = \langle u_1, u_2, \dots, u_m, u_1 \rangle$  and  $C_2 = \langle v_1, v_2, \dots, v_m, v_1 \rangle$  beginning at  $x$  are *independent* if  $u_1 = v_1 = x$  and  $u_i \neq v_i$  for  $2 \leq i \leq m$ . Cycles with equal length  $\{C_i\}_{i=1}^n$  beginning at  $x$  are *mutually independent* if every two cycles are independent. Two hamiltonian paths  $P_1 = \langle u_1, u_2, \dots, u_{|V(G)|} \rangle$  and  $P_2 = \langle v_1, v_2, \dots, v_{|V(G)|} \rangle$  are *independent beginning at  $x$*  [5] if  $u_1 = v_1 = x$  and  $u_i \neq v_i$  for  $2 \leq i \leq |V(G)|$ , denoted  $P_1 : x \rightarrow u_{|V(G)|}$  and  $P_2 : x \rightarrow v_{|V(G)|}$ . Hamiltonian paths  $\{P_i\}_{i=1}^n$  are *mutually independent hamiltonian paths beginning at  $x$*  [5] if any two of them are independent beginning at  $x$ .

An  $n$ -dimensional hypercube, denoted by  $Q_n$ , is a graph with  $2^n$  vertices, and each vertex  $u$  can be distinctly labeled by an  $n$ -bit binary string,  $u = u_{n-1}u_{n-2}\dots u_1u_0$ . There is an edge between two vertices if and only if their binary labels differ in exactly one bit position. Let  $(u, v)$  be an edge in  $Q_n$ . If the binary labels of  $u$  and  $v$  differ in  $i$ th position, then the edge between them is said to be in  $i$ th dimension and the edge  $(u, v)$  is called an  $i$ th dimension edge. We use  $Q_{n-1}^0$  to denote

the subgraph of  $Q_n$  induce by  $\{u \in V(Q_n) \mid u_i = 0\}$  and  $Q_{n-1}^1$  to denote the subgraph of  $Q_n$  induced by  $\{u \in V(Q_n) \mid u_i = 1\}$ .  $Q_{n-1}^0$  and  $Q_{n-1}^1$  are all isomorphic to  $Q_{n-1}$ .  $Q_n$  can be decomposed into  $Q_{n-1}^0$  and  $Q_{n-1}^1$  by dimension  $i$ , and  $Q_{n-1}^0$  and  $Q_{n-1}^1$  are  $(n-1)$ -dimensional subcubes of  $Q_n$  induced by the vertices with the  $i$ th bit position being 0 and 1 respectively. For each vertex  $u$  in  $Q_{n-1}^i$ ,  $i = \{0, 1\}$ , there is exactly one vertex in  $Q_{n-1}^{i-1}$ , denoted by  $\bar{u}$ , such that  $(u, \bar{u})$  is an edge in  $Q_n$ . There are many studies on the hypercubes [5], [9], [11], [12], [14], [15].

We now introduce a new concept. Let  $u$  and  $v$  be two distinct vertices of a bipartite graph  $G$  and let  $l$  be a positive integer length,  $d_G(u, v) \leq l \leq |V(G) - 1|$  with  $(l - d_G(u, v))$  being even. We say that the pair of vertices  $u, v$  is  $(m, l)$ -mutually independent bipanconnected if there exist  $m$  mutually independent paths  $\{P_i\}_{i=1}^m$  with length  $l$  from  $u$  to  $v$ . In this paper, we explore yet another strong property of the hypercubes. We prove that every pair of vertices  $u$  and  $v$  in the  $n$ -dimensional hypercube, with  $d_{Q_n}(u, v) \geq n - 1$ , is  $(n - 1, l)$ -mutually independent bipanconnected for every  $l$ ,  $d_{Q_n}(u, v) \leq l \leq |V(Q_n) - 1|$  with  $(l - d_{Q_n}(u, v))$  being even. As for  $d_{Q_n}(u, v) \leq n - 2$ , it is also  $(n - 1, l)$ -mutually independent bipanconnected if  $l \geq d_{Q_n}(u, v) + 2$ , and is only  $(l, l)$ -mutually independent bipanconnected if  $l = d_{Q_n}(u, v)$ . Our result strengthens a previous results of Sun et al. [14], and Li et al. [9]. Li et al. [9] proved that the hypercube  $Q_n$  is bipanconnected for  $n \geq 2$ . Sun et al. [14] proved that there are  $n - 1$  mutually independent hamiltonian paths in  $Q_n$  between every two vertices from different partite sets for  $n \geq 4$ . The number “ $n - 1$ ” in our result is tight as we have the following observation. Because each vertex of the hypercube  $Q_n$  has exactly  $n$  edges incident with it, we can expect at most  $n - 1$  mutually independent paths when the given two vertices are adjacent.

## II. PRELIMINARIES

In order to prove our claim, we need some previous results. The following results state that there exist  $n - 1$  mutually independent hamiltonian paths between two vertices. We shall strengthen the result by showing that there exist  $n - 1$  mutually independent paths of length  $l$  between two vertices, for every reasonable length  $l$ .

**Theorem 1.** [14] *Let  $x$  and  $y$  be two vertices from different partite sets of  $Q_n$ , for  $n \geq 4$ . Then there exist  $n - 1$  mutually independent hamiltonian paths joining  $x$  to  $y$ .*

**Theorem 2.** [14] *For  $n \geq 4$ , there are  $n$  mutually independent hamiltonian cycles beginning at any vertex  $x$  in  $Q_n$ .*

A hamiltonian laceable graph  $G$  is *hyper hamiltonian laceable* if for any vertex  $u$ , there is a hamiltonian path of  $G - \{u\}$  between every pair of vertices in the opposite partite set of  $u$ .

**Theorem 3.** [8] *For  $n \geq 2$ , the hypercube  $Q_n$  is hyper hamiltonian laceable.*

**Lemma 1.** [4] *Let  $F_v$  be a set of faulty vertices in  $Q_n$ . For  $n \geq 3$ , if  $|F_v| \leq n - 2$ , there exists a path of  $Q_n - F_v$  with*

*any odd length  $l$ ,  $3 \leq l \leq 2^n - 2|F_v| - 1$ , between any two adjacent vertices.*

**Lemma 2.** [14]  *$Q_n - \{x, y\}$  is hamiltonian laceable, if  $x$  and  $y$  are any two vertices from different partite sets of  $Q_n$  with  $n \geq 4$ .*

**Lemma 3.** [5] *In  $Q_n$ ,  $n \geq 2$ , let  $u$  be any vertex, and  $v_1, v_2, \dots, v_{n-1}$  be any  $n - 1$  vertices in the opposite partite set of  $u$ . There exist  $n - 1$  mutually independent hamiltonian paths beginning at  $u$  of  $Q_n$  such that  $\{P_i : u \rightarrow v_i\}_{i=1}^{n-1}$ .*

## III. MUTUALLY INDEPENDENT BIPANCONNECTED PROPERTY OF HYPERCUBE

**Lemma 4.** *Let  $x$  and  $y$  be two vertices from different partite sets of  $Q_n$  with  $n \geq 4$ . There exists a path of every odd length from 1 to  $2^n - 3$  joining any two adjacent fault-free vertices in  $Q_n - \{x, y\}$ .*

*Proof:* Let  $u, v$  be two adjacent fault-free vertices in  $Q_n - \{x, y\}$ . Because  $u$  and  $v$  are adjacent fault-free vertices, there exists a path of length 1 joining from  $u$  to  $v$  in  $Q_n - \{x, y\}$ . According to Lemma 1, there exists a path of every odd length from 3 to  $2^n - 2|2| - 1 (= 2^n - 5)$  joining  $u$  to  $v$  in  $Q_n - \{x, y\}$ . Then by Lemma 2, there exists a path of length  $2^n - 3$  joining  $u$  to  $v$  in  $Q_n - \{x, y\}$ . Therefore, the lemma holds. ■

Sun et al. [14] proved that any two hamiltonian path connecting 000 and 100 in  $Q_3$  are not independent, in other words, there do not exist 2 mutually independent hamiltonian paths in  $Q_3$  between 000 and 100. So, we will prove our theorem beginning from  $n \geq 4$  for  $Q_n$ . We found that there are only  $d$  mutually independent paths with length  $d$  if  $d_{Q_n}(u, v) = d$ . In order to see this, we have the following lemma.

**Lemma 5.** *Let  $u$  and  $v$  be two vertices of  $Q_n$  with  $d_{Q_n}(u, v) = d$ , there are  $d$  and at most  $d$  mutually independent paths with length  $d$  joining from  $u$  to  $v$ .*

*Proof:* By the symmetric property of the hypercubes, we may assume that  $u$  is the vertex with  $n$  bits containing  $n$  0's, and  $v$  is the vertex with  $n$  bits containing  $d$  1's. In order to see the basic idea, we first give an example  $n = 6$ . In  $Q_6$ . Let  $u = 000000$  and  $v = 001111$  then  $d_{Q_6}(u, v) = 4$ . We can construct 4 mutually independent paths with length 4 between  $u$  and  $v$ .

$$\begin{aligned} P_0 &= \langle u, 000001, 000011, 000111, v \rangle, \\ P_1 &= \langle u, 000010, 000110, 001110, v \rangle, \\ P_2 &= \langle u, 000100, 001100, 001101, v \rangle, \text{ and} \\ P_3 &= \langle u, 001000, 001001, 001011, v \rangle. \end{aligned}$$

For general  $n$ , let  $u = 0 \dots 0 = 0^n$  and  $v = 0 \dots 01 \dots 1 =$

$0^{n-d}1^d$ , then  $d_{Q_n}(u, v) = d$ .

$$\begin{aligned}
P_0 &= \langle 0^n, 0^{n-1}1, 0^{n-2}1^2, \dots, 0^{n-d+1}1^{d-1}, 0^{n-d}1^d \rangle, \\
P_1 &= \langle 0^n, 0^{n-2}10, 0^{n-3}1^20, \dots, 0^{n-d}1^{d-2}01, 0^{n-d}1^d \rangle, \\
P_2 &= \langle 0^n, 0^{n-3}10^2, 0^{n-4}1^20^2, \dots, 0^{n-d}1^{d-3}01^2, 0^{n-d}1^d \rangle, \\
P_3 &= \langle 0^n, 0^{n-4}10^3, 0^{n-5}1^20^3, \dots, 0^{n-d}1^{d-4}01^3, 0^{n-d}1^d \rangle, \\
&\vdots \\
P_{d-2} &= \langle 0^n, 0^{n-d-1}10^{d-2}, 0^{n-d}1^20^{d-2}, 0^{n-d}1^20^{d-3}1, \\
&\quad 0^{n-d}1^20^{d-4}1^2, \dots, 0^{n-d}1^201^{d-3}, 0^{n-d}1^d \rangle, \\
P_{d-1} &= \langle 0^n, 0^{n-d}10^{d-1}, 0^{n-d}10^{d-2}1, 0^{n-d}10^{d-3}1^2, \\
&\quad 0^{n-d}10^{d-4}1^3, \dots, 0^{n-d}101^{d-2}, 0^{n-d}1^d \rangle.
\end{aligned}$$

$\{P_0, P_1, \dots, P_{d-1}\}$  form  $d$  mutually independent paths with length  $d$  joining  $u$  to  $v$ . If there exists a  $(d+1)$ th path  $P'$  with length  $d$  between  $u$  and  $v$  such that  $P'$  is mutually independent to the first  $d$  paths. So the first vertex after the beginning vertex  $u$  of  $P'$  has to be different from all those of  $P_i$   $i = 0$  to  $d-1$ . Without loss of generality, assume that the first vertex after the beginning vertex  $u$  of  $P'$  is  $(x)^i = 0^i10^{n-i-1}$  for  $0 \leq i \leq n-d-1$ . It is easy to see that  $d_{Q_n}((x)^i, v) = d+1$ , since there are  $d+1$  distinct bits between  $(x)^i$  and  $v$ . Therefore, it is impossible to find out a  $(d+1)$ th path with length  $d$  between  $u$  and  $v$  which is independent to  $P_0, P_1, \dots, P_{d-1}$ . ■

We now show our main result Theorem 5 below. Our proof is by induction on  $n$ , for  $Q_n$ . The base case is  $n = 4$ .

**Theorem 4.** *Let  $u$  and  $v$  be a pair of vertices of  $Q_4$ . If  $d_{Q_4}(u, v) \geq 3$ ,  $Q_4$  is  $(3, l)$ -mutually independent bipanconnected for every  $l$ ,  $d_{Q_4}(u, v) \leq l \leq 2^4 - 1$  with  $(l - d_{Q_4}(u, v))$  being even. As for  $d_{Q_4}(u, v) \leq 2$ , it is also  $(3, l)$ -mutually independent bipanconnected if  $l \geq d_{Q_4}(u, v) + 2$ , and is only  $(l, l)$ -mutually independent bipanconnected if  $l = d_{Q_4}(u, v)$ .*

We will use the notation  $P_i^k$  or  $R_i^k$  to denote a path  $i$  with length  $k$ .

**Lemma 6.** *Let  $u$  and  $v$  be two adjacent vertices of  $Q_n$  for  $n \geq 4$ . There exist  $n-1$  mutually independent paths  $\{P_i^l\}_{i=1}^{n-1}$  of  $Q_n$  with any odd length  $l$ ,  $3 \leq l \leq 2^n - 1$ , joining from  $u$  to  $v$ .*

*Proof:* We choose a dimension to divide the hypercube  $Q_n$  into two subcubes  $Q_{n-1}^0$  and  $Q_{n-1}^1$  such that  $u$  is a black vertex in  $Q_{n-1}^0$  and  $v$  a white vertex in  $Q_{n-1}^1$ . Notice that  $\bar{u} = v$ . According to Theorem 2, there exist  $n-1$  mutually independent hamiltonian cycles  $\{C_i\}_{i=1}^{n-1}$  in  $Q_{n-1}^0$  beginning at  $u$ . For each  $k$ ,  $1 \leq k \leq 2^{n-1} - 1$ , let  $C_i = \langle u, R_i^k, x_{i,k}, x_{i,k+1}, \dots, x_{i,2^{n-1}-1}, u \rangle$  for  $1 \leq i \leq n-1$ , where  $R_i^k = \langle u, x_{i,1}, x_{i,2}, \dots, x_{i,k} \rangle$  and  $|R_i^k| = k$ . Let  $S_i^k = \langle \bar{x}_{i,k}, \dots, \bar{x}_{i,2}, \bar{x}_{i,1}, \bar{u} \rangle$  for  $1 \leq i \leq n-1$ . Combine  $R_i^k$  and  $S_i^k$ , we let  $P_i^{2k+1} = \langle u, R_i^k, x_{i,k}, \bar{x}_{i,k}, S_i^k, \bar{u} = v \rangle$ ,  $1 \leq k \leq 2^{n-1} - 1$ , for  $1 \leq i \leq n-1$ . Then  $P_i^{2k+1}$  is a path joining  $u$  to  $v$  with length  $2k+1$ . Since  $1 \leq k \leq 2^{n-1} - 1$  so  $3 \leq 2k+1 \leq 2^n - 1$ . Therefore, there exist  $n-1$

mutually independent paths  $\{P_i^l\}_{i=1}^{n-1}$  with any odd length  $l$ ,  $3 \leq l \leq 2^n - 1$ , joining from  $u$  to  $v$ . ■

**Lemma 7.** *Let  $u$  and  $v$  be two vertices from the same partite set of  $Q_n$  for  $n \geq 4$ . There exist  $n-1$  mutually independent paths  $\{P_i^l\}_{i=1}^{n-1}$  of  $Q_n$  with any even length  $l$ ,  $d_{Q_n}(u, v) + 2 \leq l \leq 2^n - 2$ , joining from  $u$  to  $v$ .*

*Proof:* We prove the statement by induction on  $n$ . By Theorem 4, the statement holds for  $n = 4$ . Suppose that the result holds for  $Q_{n-1}$ ,  $n \geq 5$ . Without loss of generality, let  $u$  and  $v$  be two black vertices of  $Q_n$ . We may choose a dimension to divide the hypercube  $Q_n$  into two subcubes  $Q_{n-1}^0$  and  $Q_{n-1}^1$  such that  $u \in Q_{n-1}^0$  and  $v \in Q_{n-1}^1$ . Therefore,  $\bar{u}$  and  $\bar{v}$  are two white vertices in  $Q_{n-1}^1$  and  $Q_{n-1}^0$ , respectively. Assume that  $d_{Q_n}(u, v) = d$  and  $d$  is even, then it is easy to see that  $d_{Q_n}(u, \bar{v}) = d_{Q_n}(u, v) - 1 = d - 1$ . According to the length  $l$  of the paths, we divide the proof into the following three cases. In each case, the length  $l$  is assumed to be an even number. We shall find  $n-1$  mutually independent paths with length  $l$  joining from  $u$  to  $v$ .

**Case 1.** For even length  $l$  and  $d+2 \leq l \leq 2^{n-1}$ .

By induction hypothesis, there exist  $n-2$  mutually independent paths  $\{R_i^k\}_{i=1}^{n-2}$  of  $Q_{n-1}^0$  with odd length  $k$ ,  $d+1 \leq k \leq 2^{n-1} - 1$ , between  $u$  and  $\bar{v}$ . For  $1 \leq i \leq n-2$ , we let  $R_i^k = \langle u, x_{i,1}, x_{i,2}, \dots, x_{i,k-1}, \bar{v} \rangle$ . Now, for each  $l$  between  $d+2$  and  $2^{n-1}$ , we show how to construct the  $n-1$  mutually independent paths with length  $l$ . Let  $P_1^{k+1} = \langle u, x_{1,1}, x_{1,2}, \dots, x_{1,k-1}, \bar{v}, v \rangle$ ,  $P_i^{k+1} = \langle u, x_{i,1}, x_{i,2}, \dots, x_{i,k-1}, \bar{x}_{i,k-1}, v \rangle$  for  $2 \leq i \leq n-2$ , and  $P_{n-1}^{k+1} = \langle u, \bar{u}, \bar{x}_{1,1}, \bar{x}_{1,2}, \dots, \bar{x}_{1,k-1}, v \rangle$ ,  $d+2 \leq k+1 \leq 2^{n-1}$ . Set  $l = k+1$ . So,  $\{P_i^l\}_{i=1}^{n-1}$  form  $n-1$  mutually independent paths with each even length  $l$ ,  $d+2 \leq l \leq 2^{n-1}$ , joining from  $u$  to  $v$ .

**Case 2.** For even length  $l$  and  $2^{n-1} + 2 \leq l \leq d + 2^{n-1} - 2$ . According to induction hypothesis, there exist  $n-2$  mutually independent paths  $\{R_i\}_{i=1}^{n-2}$  of  $Q_{n-1}^0$  with odd length  $d-1$  between  $u$  and  $\bar{v}$ . Without loss of generality, we write  $R_i = \langle u, x_{i,1}, x_{i,2}, \dots, x_{i,d-2}, \bar{v} \rangle$  for  $1 \leq i \leq n-2$ . For each  $m$ ,  $2 \leq m \leq d-2$  and  $m$  is even, by Lemma 3, there exist  $n-2$  mutually independent hamiltonian paths  $\{S_i\}_{i=1}^{n-2}$  of  $Q_{n-1}^1$  beginning at  $v$  such that  $\{S_i : v \rightarrow \bar{x}_{i,m}\}_{i=1}^{n-2}$ . For  $1 \leq i \leq n-2$ , let  $P_i^{m+2^{n-1}} = \langle u, x_{i,1}, x_{i,2}, \dots, x_{i,m}, \bar{x}_{i,m}, (S_i)^{-1}, v \rangle$ ,  $2^{n-1} + 2 \leq m + 2^{n-1} \leq d + 2^{n-1} - 2$ . Set  $l = m + 2^{n-1}$ .

We have the first  $n-2$  mutually independent paths with each even length  $l$ ,  $2^{n-1} + 2 \leq l \leq d + 2^{n-1} - 2$  joining  $u$  to  $v$ . Finally, we construct the  $(n-1)$ th path joining  $u$  to  $v$ . Let  $z$  be any white vertex in  $Q_{n-1}^1$ . By Lemma 4, there exists a path  $T^k$  of  $Q_{n-1}^1 - \{v, z\}$  with any odd length  $k$ ,  $1 \leq k \leq d-3$ , joining  $\bar{u}$  to  $\bar{x}_{n-1,1}$ , and by Theorem 3, there exists a hamiltonian path  $U$  of  $Q_{n-1}^0 - \{u\}$  between  $x_{n-1,1}$  to  $\bar{v}$ . Let  $P_{n-1}^{k+2^{n-1}+1} = \langle u, \bar{u}, T^k, \bar{x}_{n-1,1}, x_{n-1,1}, U, \bar{v}, v \rangle$ ,  $2^{n-1} + 2 \leq k + 2^{n-1} + 1 \leq d + 2^{n-1} - 2$ . Set  $l = k + 2^{n-1} + 1$ . So,  $\{P_i^l\}_{i=1}^{n-1}$  form  $n-1$  mutually independent paths with each even length  $l$ ,  $2^{n-1} + 2 \leq l \leq d + 2^{n-1} - 2$ , joining from  $u$  to  $v$ .

**Case 3.** For even length  $l$  and  $d + 2^{n-1} - 2 \leq l \leq 2^n - 2$ . Again, by induction hypothesis, there exist  $n - 2$  mutually independent paths  $\{R_i^m\}_{i=1}^{n-2}$  between  $u$  and  $\bar{v}$  in  $Q_{n-1}^0$  with odd length  $m$ ,  $d - 1 \leq m \leq 2^{n-1} - 1$ . Let  $R_i^m = \langle u, x_{i,1}, x_{i,2}, \dots, x_{i,m-1}, \bar{v} \rangle$  for  $1 \leq i \leq n - 2$ . By Lemma 3, there exist  $n - 2$  mutually independent hamiltonian paths  $\{S_i\}_{i=1}^{n-2}$  of  $Q_{n-1}^1$  beginning at  $v$  such that  $\{S_i : v \rightarrow \bar{x}_{i,m-1}\}_{i=1}^{n-2}$ . Let  $P_i^{m+2^{n-1}-1} = \langle u, x_{i,1}, x_{i,2}, \dots, x_{i,m-1}, \bar{x}_{i,m-1}, (S_i)^{-1}, v \rangle$  for  $1 \leq i \leq n - 2$ ,  $d + 2^{n-1} - 2 \leq m + 2^{n-1} - 1 \leq 2^n - 2$ . Set  $l = m + 2^{n-1} - 1$ . We have the first  $n - 2$  mutually independent paths with each even length  $l$ ,  $d + 2^{n-1} - 2 \leq l \leq 2^n - 2$  joining  $u$  to  $v$ . Finally, we construct the  $(n - 1)$ th path joining  $u$  to  $v$ . Assume that  $z$  is any white vertex in  $Q_{n-1}^1$ . According to Lemma 4, there exists a path  $T^k$  of  $Q_{n-1}^1 - \{v, z\}$  with any odd length  $k$ ,  $d - 3 \leq k \leq 2^{n-1} - 3$ , joining  $\bar{u}$  to  $\bar{x}_{n-1,1}$ , and by Theorem 3, there exists a hamiltonian path  $U$  of  $Q_{n-1}^0 - \{u\}$  between  $x_{n-1,1}$  to  $\bar{v}$ . Let  $P_{n-1}^{k+2^{n-1}+1} = \langle u, \bar{u}, T^k, \bar{x}_{n-1,1}, x_{n-1,1}, U, \bar{v}, v \rangle$ ,  $d + 2^{n-1} - 2 \leq k + 2^{n-1} + 1 \leq 2^n - 2$ . Set  $l = k + 2^{n-1} + 1$ . So,  $\{P_i^l\}_{i=1}^{n-1}$  form  $n - 1$  mutually independent paths with each even length  $l$ ,  $d + 2^{n-1} - 2 \leq l \leq 2^n - 2$ , joining from  $u$  to  $v$ . ■

**Lemma 8.** Let  $u$  and  $v$  be two nonadjacent vertices from different partite sets of  $Q_n$  for  $n \geq 4$ . There exist  $n - 1$  mutually independent paths  $\{P_i^l\}_{i=1}^{n-1}$  of  $Q_n$  with any odd length  $l$ ,  $d_{Q_n}(u, v) + 2 \leq l \leq 2^n - 1$ , joining from  $u$  to  $v$ .

By Theorem 4, Lemma 5, Lemma 6, Lemma 7, and Lemma 8, we have the following theorem.

**Theorem 5.** Let  $u$  and  $v$  be any pair of vertices of  $Q_n$ . For  $d_{Q_n}(u, v) \geq n - 1$ ,  $Q_n$  is  $(n - 1, l)$ -mutually independent bipanconnected for every  $l$ ,  $d_{Q_n}(u, v) \leq l \leq 2^n - 1$  with  $(l - d_{Q_n}(u, v))$  being even. As for  $d_{Q_n}(u, v) \leq n - 2$ , it is also  $(n - 1, l)$ -mutually independent bipanconnected if  $l \geq d_{Q_n}(u, v) + 2$ , and is only  $(l, l)$ -mutually independent bipanconnected if  $l = d_{Q_n}(u, v)$ .

#### IV. CONCLUSION

In this paper, we explore yet another strong property of the hypercubes. We prove that every pair of vertices  $u$  and  $v$  in the  $n$ -dimensional hypercube, with  $d_{Q_n}(u, v) \geq n - 1$ , is  $(n - 1, l)$ -mutually independent bipanconnected for every  $l$ ,  $d_{Q_n}(u, v) \leq l \leq |V(Q_n) - 1|$  with  $(l - d_{Q_n}(u, v))$  being even. As for  $d_{Q_n}(u, v) \leq n - 2$ , it is also  $(n - 1, l)$ -mutually independent bipanconnected if  $l \geq d_{Q_n}(u, v) + 2$ , and is only  $(l, l)$ -mutually independent bipanconnected if  $l = d_{Q_n}(u, v)$ . Our result strengthens a previous results of Sun et al. [14], and Li et al. [9]. Li et al. [9] proved that the hypercube  $Q_n$  is bipanconnected for  $n \geq 2$ . Sun et al. [14] proved that there are  $n - 1$  mutually independent hamiltonian paths in  $Q_n$  between every two vertices from different partite sets for  $n \geq 4$ . The number “ $n - 1$ ” in our result is tight as we have the following observation. Because each vertex of the hypercube  $Q_n$  has exactly  $n$  edges incident with it, we can expect at most  $n - 1$

mutually independent paths when the given two vertices are adjacent.

#### ACKNOWLEDGMENT

This research was partially supported by the National Science Council of the Republic of China under contract NSC 96-2221-E-009-137-MY3, and the Aiming for the Top University and Elite Research Center Development Plan.

#### REFERENCES

- [1] Y. Alavi and J. E. Williamson, Panconnected Graph, *Studia Scientiarum Mathematicarum Hungarica*, **10** (1975) 19–22.
- [2] J. A. Bondy, Pancyclic Graphs, I, *Journal of Combinatorial Theory, Series B*, **11** (1971) 80–84.
- [3] S.-Y. Hsieh and T.-J. Lin, Path and Cycle Embedding of 3-ary  $N$ -cubes, *Proceedings of the Second International Conference on Innovative Computing, Informatio and Control, (ICICIC 2007)*, (2007) 233–236.
- [4] S.-Y. Hsieh and T.-H. Shen, Edge-bipancyclicity of a Hypercube with Faulty Vertices and Edges, *Discrete Applied Mathematics*, **156** (2008) 1802–1808.
- [5] S.-Y. Hsieh and Y.-F. Weng, Fault-Tolerant Embedding of Pairwise Independent Hamiltonian Paths on a Faulty Hypercube with Edge Faults, *Theory of Computing Systems*. Online First.
- [6] L.-H. Hsu and C.-K. Lin, *Graph Theory and Interconnection Networks*, CRC Press, New York, 2008.
- [7] Y. Kikuchi and T. Araki, Edge-bipancyclicity and Edge-fault-tolerant Bipancyclicity of Bubble-sort Graphs, *Information Processing Letters*, **100** (2006) 52–59.
- [8] M. Lewinter and W. Widulski, Hyper-Hamilton Laceable and Caterpillar-Spannable Product Graphs, *Computer and Mathematics with Applications*, **34** (1997) 99–104.
- [9] T.-K. Li, C.-H. Tsai, J. J. M. Tan, and L.-H. Hsu, Bipanconnectivity and Edge-fault-tolerant Bipancyclic of Hypercubes, *Information Processing Letters*, **21** (2003) 107–110.
- [10] C.-K. Lin, H.-M. Huang, L.-H. Hsu, and S. Bau, Mutually Hamiltonian Paths in Star Networks, *Networks*, **46** (2005) 110–117.
- [11] G. Simmons, Almost all  $n$ -dimensional Rectangular Lattices are Hamiltonian Laceable, *Congressus Numerantium*, **21** (1978) 103–108.
- [12] Y. Saad and M. H. Schultz, Topological Properties of Hypercubes, *IEEE Transactions on Computers*, **37** (1988) 867–872.
- [13] L. A. Stewart and Y. Xiang, Bipanconnectivity and Bipancyclicity in  $k$ -ary  $n$ -cubes, *IEEE Transactions on Parallel and Distributed Systems*. Online First.
- [14] C.-M. Sun, C.-K. Lin, H.-M. Huang, and L.-H. Hsu, Mutually Independent Hamiltonian Paths and Cycles in Hypercubes, *Journal of Interconnection Networks*, **7** (2006) 235–255.
- [15] C. H. Tsai, Linear array and ring embeddings in conditional faulty hypercubes, *Theoretical Computer Science*, **314** (2004) 431–443.