

RESEARCH ARTICLE

Open Access

Role of the small RNA RyhB in the Fur regulon in mediating the capsular polysaccharide biosynthesis and iron acquisition systems in *Klebsiella pneumoniae*

Su-Hua Huang^{1†}, Chien-Kuo Wang^{1†}, Hwei-Ling Peng², Chien-Chen Wu², Ying-Tsong Chen^{3,4,5}, Yi-Ming Hong⁶ and Ching-Ting Lin^{6*}

Abstract

Background: The capsular polysaccharide (CPS) and iron acquisition systems are important determinants of *Klebsiella pneumoniae* infections, and we have previously reported that the ferric uptake repressor (Fur) can play dual role in iron acquisition and CPS biosynthesis. In many bacteria, Fur negatively controls the transcription of the small non-coding RNA RyhB to modulate cellular functions and virulence. However, in *K. pneumoniae*, the role played by RyhB in the Fur regulon has not been characterised. This study investigated Fur regulation of *ryhB* transcription and the functional role of RyhB in *K. pneumoniae*.

Results: Deletion of *fur* from *K. pneumoniae* increased the transcription of *ryhB*; the electric mobility shift assay and the Fur-titration assay revealed that Fur could bind to the promoter region of *ryhB*, suggesting that Fur directly represses *ryhB* transcription. Additionally, in a Δfur strain with elevated CPS production, deletion of *ryhB* obviously reduced CPS production. The following promoter-reporter assay and quantitative real-time PCR of *cps* genes verified that RyhB activated *orf1* and *orf16* transcription to elevate CPS production. However, deletion of *ryhB* did not affect the mRNA levels of *rcaA*, *rmpA*, or *rmpA2*. These results imply that Fur represses the transcription of *ryhB* to mediate the biosynthesis of CPS, which is independent of RcsA, RmpA, and RmpA2. In addition, the Δfur strain's high level of serum resistance was attenuated by the deletion of *ryhB*, indicating that RyhB plays a positive role in protecting the bacterium from serum killing. Finally, deletion of *ryhB* in Δfur reduced the expression of several genes corresponding to 3 iron acquisition systems in *K. pneumoniae*, and resulted in reduced siderophore production.

Conclusions: The regulation and functional role of RyhB in *K. pneumoniae* is characterized in this study. RyhB participates in Fur regulon to modulate the bacterial CPS biosynthesis and iron acquisition systems in *K. pneumoniae*.

Keywords: RyhB, Fur, Capsular polysaccharide, Iron acquisition system, *Klebsiella pneumoniae*

* Correspondence: gingting@mail.cmu.edu.tw

†Equal contributors

⁶School of Chinese Medicine, China Medical University, Taichung 40402, Taiwan

Full list of author information is available at the end of the article

Background

Klebsiella pneumoniae, a member of *Enterobacteriaceae*, is a rod-shaped gram-negative opportunistic pathogen. A common cause of nosocomial infection, it is also found in various community-acquired infections, including bacteraemia, septicaemia, and urinary tract and respiratory infections, particularly in immunocompromised patients [1-4]. In Asian countries, especially Taiwan and Korea, *K. pneumoniae* is the predominant pathogen found in pyogenic liver abscess in diabetic patients [2,3,5]. The rapid development of antimicrobial resistance in *K. pneumoniae* has further troubled the clinical choices for treatments [6,7]. Studies of the pathogenic mechanisms of *K. pneumoniae* are, therefore, essential in identifying new targets for the development of antibacterial agents.

Multiple virulence factors have been identified to be involved in *K. pneumoniae* infection, which include capsular polysaccharide (CPS), lipopolysaccharides, fimbriae, iron-acquisition system, and antibiotic resistance. Among these factors, CPS is probably considered the major determinants of pathogenesis. The pyogenic liver abscess isolates often carry heavy CPS that could protect the bacteria from phagocytosis and killing by serum factors [8,9]. Apart from the antiphagocytic function, *Klebsiella* CPS also helps the bacterial colonization and biofilm formation at the infection sites [10-12]. The capsular serotypes of *K. pneumoniae* have been classified as more than 77 recognized capsular antigens [13,14]. In Taiwan, a high prevalence of K1 and K2 serotypes of *K. pneumoniae* was documented in liver abscess of diabetes mellitus patients [15]. The *cps* gene clusters that are responsible for the synthesis of different serotypes of CPS have been determined [16]. The K2 *cps* gene cluster of *K. pneumoniae* Chedid contains a total number of 19 open reading frames (ORFs) organized into three transcription units, *orf1-2*, *orf3-15*, and *orf16-17* [16]. In the previous studies, numerous regulatory systems were demonstrated to control the biosynthesis of CPS via regulating the *cps* transcriptions in *K. pneumoniae*, such as the Rcs system, RmpA, RmpA2, KvhR, KvgAS, and KvhAS [17-20]. Among these, ferric uptake regulator (Fur) represses the gene expression of *rcaA*, *rmpA*, and *rmpA2* to decrease CPS biosynthesis [21,22]. Therefore, overlapping regulons governed the regulation of these assorted virulence genes in response to numerous stress conditions.

Bacterial cells are constantly challenged by various environmental stresses from their natural habitats. Similar to many gastrointestinal (GI) pathogens, *K. pneumoniae* faces several challenges during infection and colonisation of the human body. These include gastric acid, the immune system, and a limited supply of oxygen and nutrients [23,24]. Among these, the concentration of

iron in the environment is critical for the control of cellular metabolism. Limitation of iron abolishes bacterial growth, but high intracellular concentrations of iron may damage bacteria because of the formation of undesired reactive oxygen species (ROS). Iron homeostasis maintained by the transport, storage, and metabolism of iron is tightly controlled by Fur in many gram-negative bacteria [25-27]. To regulate gene transcription, Fur protein functions as a dimer with Fe²⁺ as a cofactor to bind to a 19-bp consensus sequence, called the Fur box (GATAATGATwATCATTATC; w = A or T), in the promoters of downstream genes [28]. In several gram-negative pathogens, Fur represses the expression of genes involved in iron homeostasis and in the regulation of multiple cellular functions such as oxidative stress, energy metabolism, acid tolerance, and virulence gene production [29-32]. In *K. pneumoniae*, Fur plays a dual role in controlling CPS biosynthesis and iron acquisition [21]. Recently, we also found that type 3 fimbriae expression and bacterial biofilm formation were also controlled by Fur and iron availability [33]. Therefore, the regulatory mechanism of Fur in control of multiple cellular function and virulence factors in *K. pneumoniae* needs to be further investigated.

Although Fur typically acts as a repressor, it also functions as a transcriptional activator for the gene expression such as *acnA*, *fumA*, and *sdhCDAB* (tricarboxylic acid [TCA] cycle enzymes), *bfr* and *ftnA* (iron storage), and *sodB* (iron superoxide dismutase [FeSOD]) [34-38]. However, positive regulation by Fur is often indirect, mediated by Fur-dependent repression of a small non-coding RNA (sRNA), RyhB [39]. RyhB negatively regulates gene expression by base pairing with mRNAs to trigger their degradation via RNase E and RNase III [40]. In many bacteria, RyhB participates in Fur-mediated positive regulation of various important cellular functions, including TCA cycle activity, resistance to oxidative stress, and iron homeostasis in *Escherichia coli* and *Vibrio cholerae* [35,39,41-43]; biofilm formation in *V. cholerae* [44]; and virulence in *Shigella dysenteriae* [45]. In *E. coli*, RyhB has been demonstrated to directly regulate more than 18 transcripts, encoding a total of 56 proteins, most of them involved in iron metabolism [35]. Although the significance of RyhB has been demonstrated in different species, to date, the regulatory relationship of RyhB and Fur, and functionality of RyhB in *K. pneumoniae* has not been studied.

In this study, the regulatory role of Fur in *ryhB* expression in *K. pneumoniae* was investigated. A *ryhB*-deletion mutant in wild type (WT) and Δfur strains and the induced expression of *ryhB* in WT were generated to demonstrate the role of RyhB in mediating CPS biosynthesis and iron acquisition systems.

Results

Fur directly represses *ryhB* expression in *K. pneumoniae*

To determine whether *K. pneumoniae ryhB* is regulated by Fur, a LacZ reporter system was used. The *ryhB* promoter was cloned into the upstream region of a promoterless *lacZ* gene in *placZ15*. The resulting plasmid pRyhB15 was then introduced into *K. pneumoniae CG43S3 ΔlacZ* and *ΔlacZΔfur*. The bacterial β-galactosidase activity was measured to assess the expression level of *ryhB*. As shown in Figure 1A, the expression of *ryhB* was higher in *ΔlacZΔfur* than *ΔlacZ*. Introduction of the complement plasmid *pfur*, but not the empty vector control (pRK415),

into *ΔlacZΔfur* restored the Fur-deletion effect. Moreover, addition of the iron chelator 2, 2-dipyridyl (Dip) to the growth medium increased *ryhB* promoter activity, suggesting that a Fur-Fe(II) complex influences *ryhB* expression. To verify that Fur directly regulates the expression of *ryhB*, an electrophoretic mobility shift assay (EMSA) was performed. As shown in Figure 1B, purified recombinant His₆-Fur protein was able to bind the upstream region of *ryhB* (*P_{ryhB}*), but not the *P_{ryhB}** fragment, whose putative Fur-box was deleted. In addition, the binding ability was abolished by the addition of 200 μM EDTA to the reaction mixture (data not shown). Furthermore, *E. coli* H1717,




Figure 1 Fur directly represses the expression of *ryhB*. (A) The β-galactosidase activities of the *K. pneumoniae CG43S3ΔlacZ* strain and the isogenic *fur* deletion mutant carrying pRyhB15 (*P_{ryhB}::lacZ*) were determined from overnight cultures grown in LB with or without Dip. The plasmids pRK415 (vector control) and *pfur* were introduced into *Δfur* to observe the complement effect. The average of triplicate experiments is shown. Error bars indicate standard deviations. (B) EMSA of the recombinant His₆::Fur and the *ryhB* promoter regions, as indicated in the margin. DNA was incubated with an increasing amount of His₆::Fur for 30 min, and then loaded onto a 5% non-denaturing polyacrylamide gel. The gel was stained with SYBR Green EMSA stain and photographed. *P_{ryhB}** indicates deletion of the *fur* box in *P_{ryhB}*. (C) Assessment of the binding of Fur to the *ryhB* promoter by using the FURTA. *E. coli* H1717 strains carrying the vector control, pT7-7, or the P1 region harboured on pT7-7 are indicated. A red colony (Lac⁺) is considered to have a FURTA-positive phenotype.

when harbouring a plasmid containing *K. pneumoniae* P_{ryhB} , also showed a Fur titration assay (FURTA)-positive phenotype (Figure 1C). The results suggest that, in an iron dependent manner, Fur suppresses *ryhB* promoter activity in *K. pneumoniae* by direct interaction with the Fur-box region upstream of *ryhB*.

RyhB activates CPS biosynthesis

In *K. pneumoniae* CG43, we found that the deletion of *fur* resulted in elevated CPS production [21,22]. To investigate if RyhB participates in Fur-regulated CPS biosynthesis, the CPS amount was assessed using measuring glucuronic acid content, which served as an indicator for *Klebsiella* K2 CPS [46], in *K. pneumoniae* strains, including WT, $\Delta ryhB$, Δfur , and $\Delta fur\Delta ryhB$, was quantified. As shown in Figure 2A, although the deletion of *ryhB* alone did not change on the amount of K2 CPS production, the elevated CPS amount in Δfur cells was abolished by the deletion of *ryhB* when the bacteria were grown in LB medium. The result indicates that Fur regulates the expression of RyhB to repress CPS biosynthesis. To confirm the RyhB expression could activate the CPS biosynthesis, the effect of RyhB induction on CPS amount was determined using an IPTG-inducible vector, pETQ. As shown in Figure 2B, the induced expression of *ryhB* in *K. pneumoniae* CG43 increased CPS production, which confirms that RyhB positively regulates CPS biosynthesis.

RyhB increased the transcriptional level of the K2 *cps* gene cluster

To investigate whether RyhB affects the expression of the three *cps* gene clusters, the mRNA levels of *orf1*, *orf3*, and *orf16* in Δfur and $\Delta fur\Delta ryhB$ strains were measured by quantitative real-time PCR (qRT-PCR). As

shown in Figure 3A, compared to the mRNA levels in the Δfur strain, the mRNA levels of *orf1* and *orf16* were apparent decreased in the $\Delta fur\Delta ryhB$ strain, and that of *orf3* also had a slight reduction in the $\Delta fur\Delta ryhB$ strain. The result suggests that overexpression of RyhB activated the *cps* gene expression. To confirm our hypothesis, the effect of *ryhB* induction on the mRNA levels of *orf1*, *orf3*, and *orf16* was tested using an IPTG-inducible vector, pETQ. As shown in Figure 3B, the mRNA levels of *orf1* and *orf16* were higher in the pETQ-*ryhB* strain with IPTG induction than the pETQ mock strain, while no significant difference in *orf3* expression was observed. To further investigate whether RyhB acts as a transcriptional activator for the promoter activity of *orf1*, *orf3*, and *orf16*, the reporter plasmids pOrf12 ($P_{orf1-2}::lacZ$), pOrf315 ($P_{orf3-15}::lacZ$), and pOrf1617 ($P_{orf16-17}::lacZ$), each carrying a *lacZ* reporter gene transcriptionally fused to the putative promoter region of the K2 *cps* gene cluster [17], were used to transform the *K. pneumoniae* strains CG43S3 $\Delta lacZ\Delta fur$ and $\Delta lacZ\Delta fur\Delta ryhB$. The promoter activity measurements shown in Figure 3C revealed that the deletion of *ryhB* in $\Delta lacZ\Delta fur$ reduced activity of $P_{orf1-2}::lacZ$ by at least 50%, while no obvious change was detected in the activity of $P_{orf3-15}::lacZ$. The activity of $P_{orf16-17}::lacZ$ was reduced by more than 75% in $\Delta lacZ\Delta fur\Delta ryhB$ as compared to the $\Delta lacZ\Delta fur$ strain. These results imply that RyhB enhances CPS biosynthesis in *K. pneumoniae* by boosting the transcriptional level of the *orf1* and *orf16* gene clusters.

RyhB does not affect the *rcaA*, *rmpA2*, and *rmpA* mRNA expression level

In previous studies, *K. pneumoniae* Fur was found to repress the expression of genes encoding the *cps*




Figure 2 RyhB activates CPS biosynthesis. (A) Comparison of CPS levels in WT, $\Delta ryhB$, Δfur , and $\Delta fur\Delta ryhB$ strains. Bacterial strains were grown in LB medium at 37°C with agitation. After 16 h of growth, the bacterial glucuronic acid content was determined. *, $P < 0.001$ compared with WT. (B) WT strains carrying the vector control (pETQ) or pETQ-*ryhB* were grown in LB with 100 μM IPTG to induce *ryhB* expression. *, $P < 0.001$ compared with WT strains carrying pETQ.




Figure 3 RyhB activates the transcriptional level of the *orf1* and *orf16*. (A) qRT-PCR analyses of the expression of the K2 *cps* genes (*orf1*, *orf3*, and *orf16*) were measured in Δfur and $\Delta fur\Delta ryhB$ strains. (B) WT strain carrying the IPTG inducible vector pETQ and pETQ-ryhB in response to 100 μ M IPTG. (C) The β -galactosidase activities of *K. pneumoniae* CG4353 $\Delta lacZ\Delta fur$ and $\Delta lacZ\Delta fur\Delta ryhB$ carrying the reporter plasmid pOrf12 ($P_{orf1-2}::lacZ$), pOrf315 ($P_{orf3-15}::lacZ$) or pOrf1617 ($P_{orf16-17}::lacZ$) were determined using log-phased cultures grown in LB broth. The results shown are an average of triplicate samples. Error bars indicate standard deviations.

regulatory proteins RcsA, RmpA, and RmpA2 [21,22]. To investigate whether RyhB affects the expression of *rcaA*, *rmpA*, and *rmpA2* to increase the *orf1* and *orf16* transcripts, the mRNA levels were measured by qRT-PCR after inducing the expression of *ryhB* in WT. However, qRT-PCR results did not show a significant effect of *ryhB* on the mRNA levels of *rmpA*, *rmpA2*, and *rcaA* (Data not shown), suggesting that the activation of RyhB on the *orf1* and *orf16* expression is not via RmpA, RmpA2, and RcsA.

Deletion of *ryhB* attenuated the higher serum resistance in Δfur strain

In addition to the roles played by RyhB and Fur in regulating the CPS amount, we suggest that RyhB and Fur may also affect the ability of the strain to resist the bactericidal effects of serum. In a human serum resistance assay, we found that the deletion of *fur* in WT increased the survival rate in treatment with 75% normal human serum from 63.3% to 87.9% (Figure 4). However, the deletion of *ryhB* in WT had no apparent effect on the survival rate on treatment with 75% serum, and the higher serum resistance in Δfur cells was abolished by the deletion of *ryhB*. This result indicates that RyhB may participate with Fur in regulating serum resistance in *K. pneumoniae*.

The regulatory role of RyhB in iron-acquisition systems

To assess whether RyhB affects iron-acquisition in *K. pneumoniae*, the Chrome azurol S (CAS) assay was used to measure siderophore secretions in Δfur and $\Delta fur\Delta ryhB$ strains (Figure 5). When bacteria were grown in M9 minimal medium ($\sim 2 \mu$ M iron) to mimic iron-limited condition, the deletion of *ryhB* in Δfur reduced the formation of the orange halo. However, this change

was not observed when bacteria were grown in LB medium ($\sim 18 \mu$ M iron). Compared to M9 minimal medium contains $\sim 2 \mu$ M iron, LB medium is considered an iron-repletion medium. Under iron-repletion, Fur is able to exert its repression on *ryhB* transcription. Thus, *ryhB*-deletion effect is difficult to observed under the growth condition that *ryhB* is poorly expressed. Our results suggest that in the regulation of iron-acquisition systems, RyhB plays a role downstream of Fur in *K. pneumoniae* under iron-limiting conditions.

To investigate the effects on downstream targets of RyhB in iron-acquisition regulons, the expression of genes corresponding to the eight putative iron-acquisition systems in *K. pneumoniae* CG43 was measured in Δfur and $\Delta fur\Delta ryhB$ by qRT-PCR (Table 1). In




Figure 4 Effect of Fur and RyhB on susceptibility to normal human serum. Survival percentage of WT, $\Delta ryhB$, Δfur , $\Delta fur\Delta ryhB$, and $\Delta galU$ (negative control) strains on treatment with 75% healthy human serum was determined, respectively. The results shown are an average of triplicate samples. Error bars indicate standard deviations.




Figure 5 Deletion of *ryhB* decreases *K. pneumoniae* Δfur siderophore production assessed using CAS assay. Each of the strains, Δfur and $\Delta fur\Delta ryhB$, was grown overnight in LB medium or M9 minimal medium, and then 5 μ l each of cultures respectively was added onto a CAS agar plate. The orange halos formed around the colonies correspond to the iron-chelating activity of the siderophores in bacteria.

M9 minimal medium, the expression of genes (*iucA*, *fepA*, *fepB*, *entC*, *fecA*, and *fecE*) corresponding to three iron-acquisition systems (aerobactin, enterobactin, and ferric citrate) was decreased by half in the $\Delta fur\Delta ryhB$ strain ($\Delta fur\Delta ryhB/\Delta fur < 0.5$). However, the expression of *fhuA* and *sitA* was significantly increased more than two-fold ($\Delta fur\Delta ryhB/\Delta fur > 2.0$). These results imply

Table 1 qRT-PCR analyses of the expression of iron-acquisition genes in *K. pneumoniae* Δfur $\Delta ryhB$ and Δfur strains

Systems	Gene	RNA expression ratio ^a $\Delta fur\Delta ryhB/\Delta fur$
Fe ³⁺		
Ferrichrome	<i>fhuA</i>	2.62 ± 0.07
Aerobactin	<i>iucA</i>	0.19 ± 0.06
Enterobactin	<i>fepA</i>	0.36 ± 0.01
	<i>fepB</i>	0.33 ± 0.05
	<i>entC</i>	0.46 ± 0.02
Ferric citrate	<i>fecA</i>	0.19 ± 0.02
	<i>fecE</i>	0.34 ± 0.03
Salmochelin	<i>iroB</i>	0.52 ± 0.05
Heme	<i>hmuR</i>	0.69 ± 0.01
Fe ²⁺		
Ferrous iron	<i>feoB</i>	0.55 ± 0.18
	<i>sitA</i>	2.81 ± 0.08

^a Mean expression ratio (±SD) of $\Delta fur\Delta ryhB$ relative to Δfur .

that RyhB activates the expression of *iucA*, *fepA*, *fepB*, *entC*, *fecA*, and *fecE*, but represses the expression of *fhuA* and *sitA*.

Discussion

In this study, we provide an initial characterisation of *K. pneumoniae* RyhB. In *K. pneumoniae*, sequence comparison indicated that the nucleotide sequence of the *ryhB* gene (91 bp) is 92.3% identical to the *E. coli* version (90 bp). However, the promoter sequence of *K. pneumoniae* *ryhB* is only 72.4% identical to that of *E. coli*. In this study, we found that the expression of *ryhB* in *K. pneumoniae* is directly repressed by Fur-Fe(II), as is the case in *E. coli* (Figure 1).

In addition, structure of the genomic neighbourhood of *ryhB* differs between the 2 species. In the *E. coli* genome, *ryhB* is found between *yhhX* and *yhhY*. In the *K. pneumoniae* genome, *ryhB* is flanked by *yhhY* and a hypothetical ORF. By Pfam search, the hypothetical ORF was found to contain a bactofilin domain (E-value = 3.7 e-24), which belongs to a new class of polymer-forming proteins that serve as versatile molecular scaffolds in a variety of cellular pathways [47]. Even though the function of this hypothetical protein in *K. pneumoniae* has not yet been investigated, we found that RyhB could strongly repress the expression of this hypothetical protein (unpublished data). This result suggests that RyhB could participate in a variety of cellular pathways in *K. pneumoniae*.

We previously showed in *K. pneumoniae*, Fur represses CPS biosynthesis via regulation of RmpA, RmpA2, and RcsA. In addition to these 3 regulators, one or more regulators may be involved in the Fur-mediated control of *cps* transcription [21]. In this study, we found that RyhB also participates in Fur-regulated CPS biosynthesis via activation of *orf1* and *orf16* transcription and is independent of the 3 regulators, RmpA, RmpA2, and RcsA (Figure 2 and 3). We want to further analyse whether any potential transcriptional regulator-binding motifs exist in the promoter sequences of *orf1* and *orf16*. We noted that a binding site typical of IscR, a transcriptional repressor that controls Fe-S biosynthesis [48], was located 172 bp upstream of the translation start site of GalF (encoded by *orf1*, 5'-ATAACCTGAACGAAAA TAAGATTAT-3'). The predication indicated that IscR could participate in control of *orf1* expression. Furthermore, a previous study reported that RyhB promotes the degradation of *iscSUA* transcripts, resulting in an increase in the ratio of apo-IscR/holo-IscR [48]. Whether RyhB activates CPS biosynthesis via regulation of the ratio of apo-IscR/holo-IscR in *K. pneumoniae* awaits further analysis. However, the regulatory mechanism of *cps* transcription is more complex than expected; whether another unknown transcriptional regulator is involved in

activation of RyhB's effect on *orf16* transcription needs to be investigated. In addition, CPS is considered the major determinant that can protect the bacteria from phagocytosis and killing by serum factors [8,9]. In this study, higher serum resistance was found in Δfur , but this higher serum resistance was attenuated by further deletion of *ryhB* (Figure 4). We suggest the protective role of RyhB against serum killing is due to the activation of CPS biosynthesis.

In *E. coli*, RyhB plays a positive role in control of the intracellular iron concentration via the degradation of nonessential iron-using proteins or an increase in siderophore production [49-51]. In this study, we also found the deletion of *ryhB* in Δfur decreased siderophore production on the CAS plate under iron-limiting condition (Figure 5). Consistent with *E. coli* [51], RyhB in *K. pneumoniae* regulates siderophore production by activating the expression of enterobactin system genes (*entC*, *fepA*, and *fepB*). In addition, we found that RyhB may activate *iucA* and *fecA* expression. Since sRNA may positively regulate its target mRNAs via an anti-antisense mechanism to disrupt an intrinsic inhibitory structure in the 5' mRNA region that sequesters the ribosome-binding site and the first translation codon [52,53], the 5'-untranslated regions of the *iuc* and *fec* operons were analysed for sequences complementary to RyhB by prediction with the bioinformatics application RNAhybrid [54] (<http://bibiserv.techfak.uni-bielefeld.de/rnahybrid/submission.html>). However, no apparent base pairing was found in the 5'-untranslated region of the *iuc* or *fec* operons, suggesting that the activation of *iucA* and *fecA* by RyhB is not a result of direct interaction. Furthermore, RyhB was found to repress the expression of *fhuA* and *sitA* in *K. pneumoniae*. In *E. coli*, RyhB represses the expression of *fhuA*, which also corresponds to our results [35]. A possible pairing between RyhB with the adjacent sequence of translational start site of *fhuA* and *sitA* was also predicted by the RNAhybrid algorithm. Alignment of the protected residues predicts that RyhB forms a 7+4+4 bp RNA duplex with the *sitA* mRNA (Additional file 1: Figure S1), but no apparent base pairing was found between RyhB and *fhuA*. However, the direct interaction of RyhB with the *sitA* mRNA remains to be confirmed. In *E. coli*, RyhB has been shown to repress several genes that are involved in iron-binding, which may increase the intracellular iron concentration, thereby allowing a better usage of iron and more complete Fur repression of these genes [35,55]. Nevertheless, this possibility in *K. pneumoniae* needs to be proven by careful experiments. In this study, the coordinated action of Fur and RyhB was found to regulate the expression of the iron acquisition systems for maintaining intracellular iron homeostasis in *K. pneumoniae*.

Conclusions

In this study, we provide an initial characterisation of *K. pneumoniae* RyhB. Our results suggest that RyhB plays an important role in the Fur regulon, which modulates the CPS biosynthesis and iron acquisition systems in *K. pneumoniae*, both of which contribute to the infectivity and survival of the bacterium.

Methods

Bacterial strains, plasmids, and media

Bacterial strains and plasmids used in this study are listed in Table 2. Primers used in this study are list in Additional file 2: Table S1. Bacteria were routinely cultured at 37°C in Luria-Bertani (LB) medium or M9 minimal medium supplemented with appropriate antibiotics. The antibiotics used include ampicillin (100 µg/ml), kanamycin (25 µg/ml), streptomycin (500 µg/ml), and tetracycline (12.5 µg/ml).

Construction of the gene-deletion mutants

Specific gene deletion was introduced into *K. pneumoniae* CG43S3 using an allelic exchange strategy as previously described [57]. The pKAS46 system was used in the selection of the mutants [59], and the mutations were respectively confirmed by PCR and Southern hybridization (data not shown).

Measurement of promoter activity

The promoter region of *ryhB* was PCR-amplified with primer pair pGT44/pGT45, and the amplicons were then cloned into placZ15 [63]. The promoter-reporter plasmids, pRyhB15, pOrf12, pOrf315, and pOrf1617, were individually mobilized into *K. pneumoniae* strains by conjugation from *E. coli* S17-1 λpir . The bacteria were grown to logarithmic phase in LB broth with or without 200 µM Dip (OD₆₀₀ of 0.7), and the β -galactosidase activity was measured as previously described [63].

EMSA

Recombinant *K. pneumoniae* Fur protein was expressed in *E. coli* and purified as previously described [22]. DNA fragment of the putative promoter region of *ryhB* was respectively PCR amplified by using specific primer sets (Table 2). The purified His₆-Fur was incubated with 10-ng DNA in a 15 µl solution containing 50 mM Tris-HCl (pH 7.5), 100 mM NaCl, 100 mM dithiothreitol, 200 µM MnCl₂, and 1 µg/µl BSA at room temperature for 20 min. The samples were then loaded onto a native gel of 5% nondenaturing polyacrylamide containing 5% glycerol in 0.5× TB buffer (45 mM Tris-HCl, pH 8.0, 45 mM boric acid). Gels were electrophoresed with a 20-mA current at 4°C and then stained with SABR safe Gel stain (Invitrogen).

Table 2 Bacterial strains and plasmids used in this study

Strains or plasmids	Descriptions	Reference or source
<i>K. pneumoniae</i>		
CG43S3	CG43 Sm ^r	[56]
$\Delta lacZ$	CG43S3 $\Delta lacZ$	[17]
Δfur	CG43S3 Δfur	[22]
$\Delta lacZ\Delta fur$	CG43S3 $\Delta lacZ\Delta fur$	[22]
$\Delta ryhB$	CG43S3 $\Delta ryhB$	This study
$\Delta fur\Delta ryhB$	CG43S3 $\Delta fur\Delta ryhB$	This study
$\Delta lacZ\Delta fur\Delta ryhB$	CG43S3 $\Delta lacZ\Delta fur\Delta ryhB$	This study
$\Delta galU$	CG43S3 $\Delta galU$	[57]
<i>E. coli</i>		
DH5a	<i>supE44</i> $\Delta lacU169$ (<i>f80 lacZ</i> $\Delta\mu$ 15) <i>hsdR17 recA1 endA1 gyrA96 thi-1 relA1</i>	[58]
BL21-RIL	<i>F ompT hsdS_B[r_gm_B]gal dcm</i> [DE3]	Laboratory stock
S17-1 λ pir H1717	<i>hsdR recA pro</i> RP4-2 [Tc::Mu; Km::Trn7] [λ pir] <i>araD139</i> $\Delta lacU169$ <i>rpsL150 relA1 flbB5301 deoC1 ptsF25 rbsR aroB fhuF::λ placMu</i>	[59,60]
Plasmids		
pKAS46	Positive selection suicide vector, <i>rpsL</i> Ap ^r Km ^r	[59]
yT&A	TA cloning vector	Yeastern
pRK415	Broad-host-range IncP cloning vector, Tc ^r	[61]
pT7-7	Cloning vector, Ap ^r	[62]
pETQ	Km ^r , protein expression vector	[61]
placZ15	Cm ^r , promoter selection vector, <i>lacZ</i> ⁺	[17]
pfur	Tc ^r , 0.8-kb fragment containing a <i>fur</i> allele cloned into pRK415	[22]
pET30c-Fur	Km ^r , 450-bp fragment encoding full-length Fur cloned into pET30c	[22]
pRyhB04	2.0 kb fragment containing an internal ~70-bp deletion in <i>ryhB</i> cloned into pKAS46	This study
pRyhB15	Cm ^r , 178-bp fragment containing the region upstream of <i>ryhB</i> cloned into placZ15	This study
pOrf12	Cm ^r , 500-bp fragment containing the region upstream of <i>Klebsiella K2 cps orf1-orf2</i> cloned into placZ15	[17]
pOrf315	Cm ^r , 900-bp fragment containing the region upstream of <i>Klebsiella K2 cps orf3-orf15</i> cloned into placZ15	[17]
pOrf1617	Cm ^r , 300-bp fragment containing the region upstream of <i>Klebsiella K2 cps orf16-orf17</i> cloned into placZ15	[17]
pT7-7- <i>pryhB</i>	178-bp fragment containing the putative <i>ryhB</i> promoter, cloned into pT7-7	This study
pETQ- <i>ryhB</i>	Km ^r , 326-bp fragment containing the promoter and coding region of <i>ryhB</i> cloned into pETQ	This study

FURTA

FURTA was performed according to the method described by Stojiljkovic *et al.* [64]. DNA sequences containing a putative Fur box were PCR amplified with specific primer sets and then cloned into pT7-7. The resulting plasmids were introduced into the *E. coli* strain H1717, and the transformants were plated onto MacConkey-lactose plates containing 100 μ g/ml ampicillin and 30 μ M Fe(NH₄)₂(SO₄)₂. The indicator strain H1717 contained a chromosomal *fhuF::lacZ* fusion, and a low affinity Fur box has been demonstrated in the *fhuF* promoter. The introduction of pT7-7 derived plasmids carrying Fur-binding sequences could thus cause the removal of Fur from the *fhuF* Fur box [60]. H1717 harboring pT7-7 was used as a negative control. Colony

phenotype was observed after incubation at 37°C for 10 h. Red colony (Lac+) denoted a FURTA-positive phenotype and indicated the binding of Fur to the DNA sequence cloned into the pT7-7 plasmid.

Extraction and quantification of CPS

CPS was extracted and quantified as previously described [65]. The glucuronic acid content, represents the amount of *K. pneumoniae* K2 CPS, was determined from a standard curve of glucuronic acid (Sigma-Aldrich) and expressed as micrograms per 10⁹ CFU [46].

qRT-PCR

Total RNAs were isolated from early-exponential-phase grown bacteria cells by use of the RNeasy midi-column

(QIAGEN) according to the manufacturer's instructions. RNA was DNase-treated with RNase-free DNase I (MoBioPlus) to eliminate DNA contamination. RNA of 100 ng was reverse-transcribed with the Transcriptor First Strand cDNA Synthesis Kit (Roche) using random primers. qRT-PCR was performed in a Roche LightCycler[®] 1.5 Instrument using LightCycler TaqMan Master (Roche). Primers and probes were designed for selected target sequences using Universal ProbeLibrary Assay Design Center (Roche-applied science) and listed in Additional file 2: Table S1. Data were analyzed using the real time PCR software of Roche LightCycler[®] 1.5 Instrument. Relative gene expressions were quantified using the comparative threshold cycle $2^{-\Delta\Delta CT}$ method with 23S rRNA as the endogenous reference.

Bacterial survival in serum

Normal human serum, pooled from healthy volunteers, was divided into equal volumes and stored at -70°C before use. Bacterial survival in serum was determined with minor modifications [57]. First, The bacteria were grown to log phase in LB broth and the viable bacterial concentration was adjusted to 1×10^6 colony forming units/ml. 1 ml of the cultures was washed twice by using phosphate-buffered saline (PBS) and resuspended in 1 ml PBS. The mixture containing 250 μl of the cell suspension and 750 μl of pooled human serum was incubated at 37°C for 60 min. The number of viable bacteria was then determined by plate counting. The survival rate was expressed as the number of viable bacteria treated with human serum compared to the number of pre-treatment. The assay was performed triple, each with triplicate samples. The data from one of the representative experiments are shown and expressed as the mean and standard deviation from the three samples. The 0% survival of *K. pneumoniae* CG43S3 $\Delta galU$ served as a negative control.

CAS assay

The CAS assay was performed according to the method described by Schwyn and Neilands [66]. Each of the bacterial strain was grown overnight in M9 minimal medium, and then 5 μl of culture was added onto a CAS agar plate. After 24 hr incubation at 37°C , effects of the bacterial siderophore production could be observed. Siderophore production was apparent as an orange halo around the colonies; absence of a halo indicated the inability to produce siderophores.

Statistical method

An unpaired *t*-test was used to determine the statistical significance and values of $P < 0.001$ were considered significant. The results of CPS quantification and qRT-PCR analysis were derived from a single experiment

representative of three independent experiments. Each sample was assayed in triplicate and the mean activity and standard deviation are presented.

Additional files

Additional file 1: Figure S1. RyhB pairs with *sitA*. The file contains supplemental figure S1 that the potential base pairing in RyhB/*sitA* mRNA in this study.

Additional file 2: Table S1. Primers used in this study. The file contains supplemental Table S1 that the detailed information of primer sets used in this study.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

The work is supported by the grants from National Science Council (NSC 97-2314-B-039-042-MY2 and NSC 99-2320-B-039-002-MY3) and China Medical University (CMU98-ASIA-01 and CMU99-ASIA-07).

Author details

¹Department of Biotechnology, Asia University, Taichung 41354, Taiwan. ²Department of Biological Science and Technology, National Chiao Tung University, Hsin Chu 30068, Taiwan. ³Institute of Genomics and Bioinformatics, National Chung Hsing University, Tai Chung City 40227, Taiwan. ⁴Biotechnology Center, National Chung Hsing University, Tai Chung City 40227, Taiwan. ⁵Institute of Molecular and Genomic Medicine, National Health Research Institutes, Miaoli County 35053, Taiwan. ⁶School of Chinese Medicine, China Medical University, Taichung 40402, Taiwan.

Authors' contributions

SHH, CKW, HLP, and CTL made substantial contributions to design and conduct the experiments. YMH performed qRT-PCR and growth experiments. SHH and CKW performed the bioinformatics analyses and interpretation of data. CCW, YTC, and HLP contributed to the writing and editing of the manuscript. CTL coordinated the study and performed manuscript editing. All authors have read and approved this work.

Received: 15 May 2012 Accepted: 9 July 2012

Published: 24 July 2012

References

1. Chou FF, Kou HK: Endogenous endophthalmitis associated with pyogenic hepatic abscess. *J Am Coll Surg* 1996, **182**(1):33–36.
2. Han SH: Review of hepatic abscess from *Klebsiella pneumoniae*. An association with diabetes mellitus and septic endophthalmitis. *West J Med* 1995, **162**(3):220–224.
3. Lau YJ, Hu BS, Wu WL, Lin YH, Chang HY, Shi ZY: Identification of a major cluster of *Klebsiella pneumoniae* isolates from patients with liver abscess in Taiwan. *J Clin Microbiol* 2000, **38**(1):412–414.
4. Peng HL, Wang PY, Wu JL, Chiu CT, Chang HY: Molecular epidemiology of *Klebsiella pneumoniae*. *Zhonghua Min Guo Wei Sheng Wu Ji Mian Yi Xue Za Zhi* 1991, **24**(3):264–271.
5. Yang YS, Siu LK, Yeh KM, Fung CP, Huang SJ, Hung HC, Lin JC, Chang FY: Recurrent *Klebsiella pneumoniae* liver abscess: clinical and microbiological characteristics. *J Clin Microbiol* 2009, **47**(10):3336–3339.
6. Paterson DL: Resistance in gram-negative bacteria: enterobacteriaceae. *Am J Med* 2006, **119**(6 Suppl 1):S20–S28. discussion S62–70.
7. Lagamayo EN: Antimicrobial resistance in major pathogens of hospital-acquired pneumonia in Asian countries. *Am J Infect Control* 2008, **36**(4 Suppl):S101–S108.
8. Sahly H, Podschun R, Oelschlaeger TA, Greiwe M, Parolis H, Hasty D, Kekow J, Ullmann U, Ofek I, Sela S: Capsule impedes adhesion to and invasion of epithelial cells by *Klebsiella pneumoniae*. *Infect Immun* 2000, **68**(12):6744–6749.
9. Lin JC, Chang FY, Fung CP, Xu JZ, Cheng HP, Wang JJ, Huang LY, Siu LK: High prevalence of phagocytic-resistant capsular serotypes of

- Klebsiella pneumoniae* in liver abscess. *Microbes Infect* 2004, 6(13):1191–1198.
10. Boddicker JD, Anderson RA, Jagnow J, Clegg S: Signature-tagged mutagenesis of *Klebsiella pneumoniae* to identify genes that influence biofilm formation on extracellular matrix material. *Infect Immun* 2006, 74(8):4590–4597.
 11. Moranta D, Regueiro V, March C, Llobet E, Margareto J, Larrate E, Garmendia J, Bengoechea JA: *Klebsiella pneumoniae* capsule polysaccharide impedes the expression of beta-defensins by airway epithelial cells. *Infect Immun* 2010, 78(3):1135–1146.
 12. Favre-Bonte S, Joly B, Forestier C: Consequences of reduction of *Klebsiella pneumoniae* capsule expression on interactions of this bacterium with epithelial cells. *Infect Immun* 1999, 67(2):554–561.
 13. Fung CP, Hu BS, Chang FY, Lee SC, Kuo BI, Ho M, Siu LK, Liu CY: A 5-year study of the seroepidemiology of *Klebsiella pneumoniae*: high prevalence of capsular serotype K1 in Taiwan and implication for vaccine efficacy. *J Infect Dis* 2000, 181(6):2075–2079.
 14. Pan YJ, Fang HC, Yang HC, Lin TL, Hsieh PF, Tsai FC, Keynan Y, Wang JT: Capsular polysaccharide synthesis regions in *Klebsiella pneumoniae* serotype K57 and a new capsular serotype. *J Clin Microbiol* 2008, 46(7):2231–2240.
 15. Fung CP, Chang FY, Lee SC, Hu BS, Kuo BI, Liu CY, Ho M, Siu LK: A global emerging disease of *Klebsiella pneumoniae* liver abscess: is serotype K1 an important factor for complicated endophthalmitis? *Gut* 2002, 50(3):420–424.
 16. Arakawa Y, Wacharotayankun R, Nagatsuka T, Ito H, Kato N, Ohta M: Genomic organization of the *Klebsiella pneumoniae* cps region responsible for serotype K2 capsular polysaccharide synthesis in the virulent strain Chedid. *J Bacteriol* 1995, 177(7):1788–1796.
 17. Lin CT, Huang TY, Liang WC, Peng HL: Homologous response regulators KvgA, KvhA and KvhR regulate the synthesis of capsular polysaccharide in *Klebsiella pneumoniae* CG43 in a coordinated manner. *J Biochem (Tokyo)* 2006, 140(3):429–438.
 18. Majdalani N, Gottesman S: The Rcs phosphorelay: a complex signal transduction system. *Annu Rev Microbiol* 2005, 59:379–405.
 19. Gottesman S, Stout V: Regulation of capsular polysaccharide synthesis in *Escherichia coli* K12. *Mol Microbiol* 1991, 5(7):1599–1606.
 20. Stout V: Regulation of capsule synthesis includes interactions of the RcsC/RcsB regulatory pair. *Res Microbiol* 1994, 145(5–6):389–392.
 21. Lin CT, Wu CC, Chen YS, Lai YC, Chi C, Lin JC, Chen Y, Peng HL: Fur regulation of the capsular polysaccharide biosynthesis and iron-acquisition systems in *Klebsiella pneumoniae* CG43. *Microbiology* 2011, 157(Pt 2):419–429.
 22. Cheng HY, Chen YS, Wu CY, Chang HY, Lai YC, Peng HL: RmpA regulation of capsular polysaccharide biosynthesis in *Klebsiella pneumoniae* CG43. *J Bacteriol* 2010, 192(12):3144–3158.
 23. De Champs C, Sauvart MP, Chanal C, Sirot D, Gazuy N, Malhuret R, Baguet JC, Sirot J: Prospective survey of colonization and infection caused by expanded-spectrum-beta-lactamase-producing members of the family Enterobacteriaceae in an intensive care unit. *J Clin Microbiol* 1989, 27(12):2887–2890.
 24. Markowitz SM, Veazey JM Jr, Macrina FL, Mayhall CG, Lamb VA: Sequential outbreaks of infection due to *Klebsiella pneumoniae* in a neonatal intensive care unit: implication of a conjugative R plasmid. *J Infect Dis* 1980, 142(1):106–112.
 25. Ernst JF, Bennett RL, Rothfield LI: Constitutive expression of the iron-enterochelin and ferrichrome uptake systems in a mutant strain of *Salmonella typhimurium*. *J Bacteriol* 1978, 135(3):928–934.
 26. Hantke K: Regulation of ferric iron transport in *Escherichia coli* K12: isolation of a constitutive mutant. *Mol Gen Genet* 1981, 182(2):288–292.
 27. Achenbach LA, Yang W: The fur gene from *Klebsiella pneumoniae*: characterization, genomic organization and phylogenetic analysis. *Gene* 1997, 185(2):201–207.
 28. Griggs DW, Konisky J: Mechanism for iron-regulated transcription of the *Escherichia coli* cir gene: metal-dependent binding of fur protein to the promoters. *J Bacteriol* 1989, 171(2):1048–1054.
 29. Hassett DJ, Sokol PA, Howell ML, Ma JF, Schweizer HT, Ochsner U, Vasil ML: Ferric uptake regulator (Fur) mutants of *Pseudomonas aeruginosa* demonstrate defective siderophore-mediated iron uptake, altered aerobic growth, and decreased superoxide dismutase and catalase activities. *J Bacteriol* 1996, 178(14):3996–4003.
 30. Ochsner UA, Vasil ML: Gene repression by the ferric uptake regulator in *Pseudomonas aeruginosa*: cycle selection of iron-regulated genes. *Proc Natl Acad Sci USA* 1996, 93(9):4409–4414.
 31. Bijlsma JJ, Waidner B, Vliet AH, Hughes NJ, Hag S, Bereswill S, Kelly DJ, Vandenbroucke-Grauls CM, Kist M, Kusters JG: The *Helicobacter pylori* homologue of the ferric uptake regulator is involved in acid resistance. *Infect Immun* 2002, 70(2):606–611.
 32. van Vliet AH, Stoof J, Poppelaars SW, Bereswill S, Homuth G, Kist M, Kuipers EJ, Kusters JG: Differential regulation of amidase- and formamidase-mediated ammonia production by the *Helicobacter pylori* fur repressor. *J Biol Chem* 2003, 278(11):9052–9057.
 33. Wu CC, Lin CT, Cheng WY, Huang CJ, Wang ZC, Peng HL: Fur-dependent MrkHI regulation of type 3 fimbriae in *Klebsiella pneumoniae* CG43. *Microbiology* 2012, 158(Pt 4):1045–1056.
 34. Hantke K: Iron and metal regulation in bacteria. *Curr Opin Microbiol* 2001, 4(2):172–177.
 35. Masse E, Vanderpool CK, Gottesman S: Effect of RyhB small RNA on global iron use in *Escherichia coli*. *J Bacteriol* 2005, 187(20):6962–6971.
 36. Andrews SC, Harrison PM, Guest JR: Cloning, sequencing, and mapping of the bacterioferritin gene (*bfr*) of *Escherichia coli* K-12. *J Bacteriol* 1989, 171(7):3940–3947.
 37. Gruer MJ, Guest JR: Two genetically-distinct and differentially-regulated aconitases (*AcnA* and *AcnB*) in *Escherichia coli*. *Microbiology* 1994, 140(Pt 10):2531–2541.
 38. Niederhoffer EC, Naranjo CM, Bradley KL, Fee JA: Control of *Escherichia coli* superoxide dismutase (*sodA* and *sodB*) genes by the ferric uptake regulation (*fur*) locus. *J Bacteriol* 1990, 172(4):1930–1938.
 39. Masse E, Gottesman S: A small RNA regulates the expression of genes involved in iron metabolism in *Escherichia coli*. *Proc Natl Acad Sci USA* 2002, 99(7):4620–4625.
 40. Masse E, Escorcia FE, Gottesman S: Coupled degradation of a small regulatory RNA and its mRNA targets in *Escherichia coli*. *Genes Dev* 2003, 17(19):2374–2383.
 41. Dubrac S, Touati D: Fur positive regulation of iron superoxide dismutase in *Escherichia coli*: functional analysis of the *sodB* promoter. *J Bacteriol* 2000, 182(13):3802–3808.
 42. Davis BM, Quinones M, Pratt J, Ding Y, Waldor MK: Characterization of the small untranslated RNA RyhB and its regulon in *Vibrio cholerae*. *J Bacteriol* 2005, 187(12):4005–4014.
 43. Argaman L, Elgrably-Weiss M, Hershko T, Vogel J, Altuvia S: RelA protein stimulates the activity of RyhB small RNA by acting on RNA-binding protein Hfq. *Proc Natl Acad Sci USA* 2012, 109(12):4621–4626.
 44. Mey AR, Craig SA, Payne SM: Characterization of *Vibrio cholerae* RyhB: the RyhB regulon and role of *ryhB* in biofilm formation. *Infect Immun* 2005, 73(9):5706–5719.
 45. Murphy ER, Payne SM: RyhB, an iron-responsive small RNA molecule, regulates *Shigella dysenteriae* virulence. *Infect Immun* 2007, 75(7):3470–3477.
 46. Blumenkrantz N, Asboe-Hansen G: New method for quantitative determination of uronic acids. *Anal Biochem* 1973, 54(2):484–489.
 47. Kuhn J, Briegel A, Morschel E, Kahnt J, Leser K, Wick S, Jensen GJ, Thanbichler M: Bactofilins, a ubiquitous class of cytoskeletal proteins mediating polar localization of a cell wall synthase in *Caulobacter crescentus*. *EMBO J* 2010, 29(2):327–339.
 48. Desnoyers G, Morissette A, Prevost K, Masse E: Small RNA-induced differential degradation of the polycistronic mRNA *iscRSUA*. *EMBO J* 2009, 28(11):1551–1561.
 49. Masse E, Salvail H, Desnoyers G, Arguin M: Small RNAs controlling iron metabolism. *Curr Opin Microbiol* 2007, 10(2):140–145.
 50. Jacques JF, Jang S, Prevost K, Desnoyers G, Desmarais M, Imlay J, Masse E: RyhB small RNA modulates the free intracellular iron pool and is essential for normal growth during iron limitation in *Escherichia coli*. *Mol Microbiol* 2006, 62(4):1181–1190.
 51. Salvail H, Lanthier-Bourbonnais P, Sobota JM, Caza M, Benjamin JA, Mendieta ME, Lepine F, Dozois CM, Imlay J, Masse E: A small RNA promotes siderophore production through transcriptional and metabolic remodeling. *Proc Natl Acad Sci USA* 2010, 107(34):15223–15228.
 52. Frohlich KS, Vogel J: Activation of gene expression by small RNA. *Curr Opin Microbiol* 2009, 12(6):674–682.
 53. Prevost K, Salvail H, Desnoyers G, Jacques JF, Phaneuf E, Masse E: The small RNA RyhB activates the translation of *shiA* mRNA encoding a permease

- of shikimate, a compound involved in siderophore synthesis. *Mol Microbiol* 2007, **64**(5):1260–1273.
54. Rehmsmeier M, Steffen P, Hochsmann M, Giegerich R: **Fast and effective prediction of microRNA/target duplexes.** *RNA* 2004, **10**(10):1507–1517.
 55. Salvail H, Masse E: **Regulating iron storage and metabolism with RNA: an overview of posttranscriptional controls of intracellular iron homeostasis.** *Wiley Interdiscip Rev RNA* 2012, **3**(1):26–36.
 56. Lai YC, Peng HL, Chang HY: **Identification of genes induced in vivo during *Klebsiella pneumoniae* CG43 infection.** *Infect Immun* 2001, **69**(11):7140–7145.
 57. Lai YC, Peng HL, Chang HY: **RmpA2, an activator of capsule biosynthesis in *Klebsiella pneumoniae* CG43, regulates K2 cps gene expression at the transcriptional level.** *J Bacteriol* 2003, **185**(3):788–800.
 58. Hanahan D: **Studies on transformation of *Escherichia coli* with plasmids.** *J Mol Biol* 1983, **166**(4):557–580.
 59. Skorupski K, Taylor RK: **Positive selection vectors for allelic exchange.** *Gene* 1996, **169**(1):47–52.
 60. Hantke K: **Selection procedure for deregulated iron transport mutants (*fur*) in *Escherichia coli* K 12: *fur* not only affects iron metabolism.** *Mol Gen Genet* 1987, **210**(1):135–139.
 61. Keen NT, Tamaki S, Kobayashi D, Trollinger D: **Improved broad-host-range plasmids for DNA cloning in gram-negative bacteria.** *Gene* 1988, **70**(1):191–197.
 62. Tabor S, Richardson CC: **A bacteriophage T7 RNA polymerase/promoter system for controlled exclusive expression of specific genes.** *Proc Natl Acad Sci USA* 1985, **82**(4):1074–1078.
 63. Lin CT, Huang TY, Liang WC, Peng HL: **Homologous response regulators KvgA, KvhA and KvhR regulate the synthesis of capsular polysaccharide in *Klebsiella pneumoniae* CG43 in a coordinated manner.** *J Biochem* 2006, **140**(3):429–438.
 64. Stojiljkovic I, Baumber AJ, Hantke K: **Fur regulon in gram-negative bacteria. Identification and characterization of new iron-regulated *Escherichia coli* genes by a *fur* titration assay.** *J Mol Biol* 1994, **236**(2):531–545.
 65. Domenico P, Schwartz S, Cunha BA: **Reduction of capsular polysaccharide production in *Klebsiella pneumoniae* by sodium salicylate.** *Infect Immun* 1989, **57**(12):3778–3782.
 66. Schwyn B, Neilands JB: **Universal chemical assay for the detection and determination of siderophores.** *Anal Biochem* 1987, **160**(1):47–56.

doi:10.1186/1471-2180-12-148

Cite this article as: Huang et al.: Role of the small RNA RyhB in the Fur regulon in mediating the capsular polysaccharide biosynthesis and iron acquisition systems in *Klebsiella pneumoniae*. *BMC Microbiology* 2012 **12**:148.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

