

A New Trans-to-Cis Photoisomerization Mechanism of Azobenzene on the $S_1(n,\pi^*)$ Surface

Eric Wei-Guang Diau

Department of Applied Chemistry and Center for Interdisciplinary Molecular Science,
National Chiao Tung University, Hsinchu, Taiwan 30050, Republic of China

Received: October 9, 2003


The rotation-inversion controversy for the photoisomerization mechanism of azobenzene has been addressed on the basis of the CASSCF calculations for the excited-state relaxed surface scan along the CNN bending (inversion), the CNNC torsion (rotation), and the concerted CNN bending (concerted inversion) reaction pathways. According to our calculated results, the inversion channel involves a substantial energy barrier and a large S_0 – S_1 energy gap so that it is a highly unfavorable channel to be considered for an efficient electronic relaxation. The rotation channel is essentially barrierless with a conical intersection close to the midpoint of the pathway to reasonably account for the subpicosecond to picosecond relaxation times observed in recent ultrafast experiments. Along the concerted inversion reaction path, a sloped conical intersection (S_0/S_1 CI_inv) was found to have the geometry close to a linear CNNC configuration. When the photoexcitation occurs in the S_2 state, the concerted inversion channel may be open and the S_0/S_1 CI_inv would become energetically accessible to produce more trans- S_0 isomers. A new mechanism is proposed to rationalize the early experimental results for the following two cases: (1) the trans-to-cis photoisomerization quantum yields on S_2 excitation are much lower than the yields on S_1 excitation; (2) the yields from both the S_2 and the S_1 excitations become essentially equal when the rotation channel is blocked by chemical modification or in an inclusion complex.

1. Introduction

The mechanism of the photoisomerization of *trans*-azobenzene has been a fundamental subject for many years not only for its potential applications in industry^{1–7} but also for its long-standing debate on the isomerization process occurring along either the inversional or the rotational pathway (Scheme 1). In solutions, the UV–visible spectra of the molecule are featured with two major absorption bands corresponding to the $S_0 \rightarrow S_1$ and the $S_0 \rightarrow S_2$ transitions. The former transition relates to a symmetry-forbidden $n-\pi^*$ transition with the maximum intensity near 440 nm whereas the latter transition corresponds to a symmetry-allowed $\pi-\pi^*$ transition with the maximum intensity near 320 nm.^{7,8} Early steady-state measurements show that the quantum yields of the trans-to-cis isomerization are wavelength-dependent: The yield measured in hexane solvent is 0.20–0.27 upon excitation to the $S_1(n,\pi^*)$ state but it drops dramatically to 0.09–0.12 for the $S_2(\pi,\pi^*)$ excitation.^{7–9} However, the quantum yields become almost wavelength-independent when rotation about the NN double bond is restricted by blocking the rotation with a cyclophane structure,¹⁰ with a crown ether,¹¹ or with the inclusion in the cyclodextrin cavity.¹² These observations imply that two different isomerization mechanisms are operated on excitation into these two electronic excited states; i.e., visible excitation to the $S_1(n,\pi^*)$ state leads to the isomerization via inversion around one nitrogen atom in the same molecular plane whereas UV excitation to the $S_2(\pi,\pi^*)$ state results in the isomerization via the rotation around the NN double bond.⁷ This conclusion is supported by early ab initio work of Monti et al.¹³ based on minimal basis set CI calculations and has been widely adopted in recent time-resolved studies using the technique of femtosecond transient absorption spectroscopy.^{14–18}

Moreover, recent time-resolved Raman study on the $S_2(\pi,\pi^*)$ excitation has demonstrated the similarity of the NN stretch

SCHEME 1



frequency between the S_1 and the S_0 states and suggested that the NN bonding retains a double bond character and thus the S_1 species has a planar structure.¹⁹ This evidence partly supports the previous conclusion that the isomerization of azobenzene on the S_1 surface is via an inversion mechanism. However, the subsequent time-resolved fluorescence studies^{20,21} of the same laboratory have made a conclusion that the rotational isomerization mechanism starting directly from the S_2 state *does not exist* and the isomerization of azobenzene takes place exclusively on the S_1 surface by an in-plane inversion mechanism regardless of the initial excitations. Apparently, such a consecutive model ($S_2 \rightarrow S_1 \rightarrow S_0$ via inversion only) cannot account for the early spectroscopic data. To rationalize for the smaller isomerization quantum yield previously observed from the S_2 excitation, a new relaxation channel in the vibrationally hot S_1 state that is open to form only the trans isomer has been proposed.^{20,21}

On the other hand, the aforementioned inversion photoisomerization mechanisms have been challenged by recent high-

level theoretical calculations. Based on the multiconfiguration self-consistent field (MCSCF) approach using the complete active space (CAS) SCF wave functions, both recent ab initio results^{22,23} have shown that the S_1 potential energy surface (PES) of *trans*-azobenzene involves substantial energy barrier along the inversion coordinate, but the surface is essentially barrierless along the rotation coordinate. Furthermore, the results of Ishikawa et al.²³ also indicate that a conical intersection (CI) between the S_0 and the S_1 states may be located near the midpoint of the rotation pathway, suggesting that the photoisomerization of azobenzene on the S_1 surface may favor a rotation channel. The rotation mechanism reported for azobenzene is also consistent with the topological feature of azomethane on the S_1 surface.²⁴

In a previous study,²⁵ we reported on the femtosecond fluorescence dynamics of *trans*-azobenzene in hexane with excitation directly into the $S_1(n,\pi^*)$ state. The observed fluorescence temporal profiles feature two components and our assignments are based on the rotation mechanism supported by recent ab initio calculations.^{22,23} The fast-decay component is due to a structural relaxation from the Franck–Condon region toward the rotational reaction path and the slow-decay component is arisen from the other nuclear motions along the rotation coordinate to search for the S_0/S_1 CI on the multidimensional PES. However, such an energy-favorable rotational isomerization channel in azobenzene is not feasible for the rotation-blocked azobenzene derivatives and a new relaxation channel must be open to account for the observed quantum yield discrepancies previously reported.⁷ To shed light on the issue for the rotation-inversion controversy, we have carried out CASSCF calculations for azobenzene to follow the minimum-energy pathways on the S_1 surface along three different reaction coordinates—the CNN bending coordinate, the CNNC torsional coordinate, and the concerted CNN bending coordinate. Our results on the CNN bending angle scan and the CNNC torsional angle scan are similar to those of recent studies,^{22,23} but we have found a new conical intersection along the concerted CNN bending coordinate responsible for the new relaxation channel to rationalize the old quantum yield results. A new mechanism is presented in this paper to provide a solution for the long-standing controversy on the subject of photoisomerization of azobenzene.

2. Method of Calculations

Our strategy for the CASSCF calculations is aimed at finding a proper active space that is capable of describing the trans-to-cis isomerization processes of azobenzene on the S_1 PES. According to the previous calculations of azomethane,²⁴ the active space should at least include six electrons distributed among four orbitals ($n-$, π , $n+$, and π^*), abbreviated by CAS-(6,4), to reasonably describe the PES along the rotational isomerization coordinate. Because there are some delocalized π^* and σ^* orbitals that are relevant to better describe the PES of azobenzene along the three coordinates of interest, we have included two more virtual orbitals (either two π^* for the rotation channel or $\pi^* + \sigma^*$ for the inversion channels) in the active space for the calculations at the CAS(6,6) level. The selected orbitals in the active space are $3a + 3b$ for the CNNC torsional angle scan (C_2), $3a' + 3a''$ for the CNN bending angle scan (C_s), and $a_g + a_u + 2b_g + 2b_u$ for the concerted CNN bending angle scan (C_{2h}).

The geometry of each point along three different reaction coordinates was optimized according to a relaxed surface scan procedure at the state-specific CAS(6,6)/4-31G level whereas

the geometry of a conical intersection between the S_0 and the S_1 states was fully optimized at the state-averaged CAS(6,6)/4-31G level. For all the stationary points optimized on the potential energy surfaces, force constant calculations were performed at the CAS(6,6)/4-31G level, the same as geometry optimization, and single point energy calculations were further carried out at the CAS(6,6)/cc-pVDZ level of theory. All the electronic structure calculations were performed using the Gaussian software package.²⁶

Because *trans*-azobenzene does not have a permanent dipole moment, the effect of the solvents affecting the PES is not expected to be significant. In fact, we have checked such an effect on the basis of the time-dependent density functional theory (TDDFT) calculations with the polarizable continuum model (PCM) implemented in Gaussian 03. In comparison with the isolated molecule in the gas phase, the vertical excitation energy of azobenzene is lower by only 0.01 and 0.05 eV in hexane and DMSO, respectively, predicted at the TD-B3LYP/6-311++G(d,p) level of theory. The solvation energies of *trans*-azobenzene (<0.05 eV) were computed to be much smaller than the uncertainties of the excitation energies (typically ~ 0.2 eV; see below) from the theoretical calculations currently employed. Therefore, all the surface-scan calculations reported in this article were carried out in the gas-phase environment without considering the negligible solvent effect.

3. Results and Discussions

Three minimum-energy pathways on the S_1 PES of azobenzene were characterized by optimizing the geometries along the CNN bending coordinate (the inversion channel), the CNNC torsional coordinate (the rotation channel), and the concerted CNN bending coordinate (the concerted inversion channel); the corresponding potential energy curves are shown in Figures 1–3, respectively. The key structural parameters and the relative energies of the relevant stationary points and conical intersections are summarized in Table 1. There are two major points worth mentioning to justify the accuracy of our calculations. First, we found that the calculated S_1 excitation energy at the Franck–Condon geometry (S_1 -FC) is 59.6 kcal mol⁻¹ at the CAS(6,6)/4-31G level, and it slightly increases to 63.4 kcal mol⁻¹ with further single-point energy calculation at the CAS-(6,6)/cc-pVDZ level of theory. Note that the observed absorption spectrum of *trans*-azobenzene has the maximum intensity at 450 nm (=63.5 kcal mol⁻¹ in energy), which is in excellent agreement with our theoretical prediction at the CAS(6,6)/cc-pVDZ level. Second, the S_1 minimum, $S_1(C_2)$, was calculated to be 47.7 kcal mol⁻¹ in energy above the S_0 minimum at the level of geometry optimization.²⁷ Further single-point energy calculation at the CAS(6,6)/cc-pVDZ level gives the relative energy of $S_1(C_2)$ with respect to $S_0(C_{2h})$ being 50.1 kcal mol⁻¹, which is again in excellent agreement with the experimental observation that the red edge of the S_1 absorption band ends at ~ 560 nm (~ 51 kcal mol⁻¹ in energy). Therefore, we expect that the energetic uncertainties for the results of the surface-scan calculations shown in Figures 1–3 are within ~ 4 kcal mol⁻¹ (or ~ 0.2 eV). The results and their dynamical significance for the photoisomerization of azobenzene are discussed in the following.

3.1. Inversion Channel. The potential energy curves shown in Figure 1 were obtained from scanning one of the CNN bending reaction coordinate (RC) when optimizing the other structural parameters on the S_1 PES with C_s symmetry. The S_1 potential energy significantly increases along the CNN bending RC toward the smaller bending angle direction. On the other




Figure 1. Minimum-energy pathway of azobenzene along the CNN bending coordinate optimized for the second root (the S_1 state) at the state-specific CAS(6,6)/4-31G level of theory. On top, the corresponding optimized structures are shown from left to right for the bending angles of 90° , 130° , 180° , and 226° , respectively.




Figure 2. Minimum-energy pathway of azobenzene along the CNNC torsional coordinate optimized for the second root (the S_1 state) at the state-specific CAS(6,6)/4-31G level of theory. On top, the corresponding optimized structures are shown from left to right for the torsional angles of 180° , 90° , and 0° , respectively. The optimized structure of the S_0/S_1 CI_rot is shown in the inset.

hand, following the in-plane CNN bending RC toward the larger bending angle direction, the S_1 potential energy curve also goes uphill and reaches a plateau area at the RC greater than 180° . The results are similar to those of Cattaneo and Persico²² and Ishikawa et al.²³ and this topological picture may provide a theoretical description for the inversion mechanism shown in Scheme 1.

The minimum-energy point of the S_1 curve, $S_1(C_s)$, is located above the ground-state minimum by $45.0 \text{ kcal mol}^{-1}$ (with ZPE corrections) at 129.4° ;²⁷ the other CNN bending angle is 127.8° .

Following the in-plane CNN bending reaction path from $S_1(C_s)$ toward the larger bending angle direction, the S_1 and the S_0 potential energy curves do come closer but there are two factors affecting the electronic relaxation through this channel. First, the energy difference between the $S_1(C_s)$ and the S_1 species at 180° , $S_1\text{-}180(C_s)$, is close to $\sim 20 \text{ kcal mol}^{-1}$, which is a substantial energy barrier to overcome along this reaction path. Second, the energy gap between the two states is larger than 10 kcal mol^{-1} at the bending angles greater than 180° (toward the cis isomer). In fact, a state-averaged calculation at the CAS-




Figure 3. Minimum-energy pathway of azobenzene along the concerted CNN bending coordinate optimized for the second root (the S_1 state) at the state-specific CAS(6,6)/4-31G level of theory. On top, the corresponding optimized structures are shown from left to right for the two equivalent bending angles equal to 92° , 128° , and 180° , respectively. The optimized structure of the S_0/S_1 CI_inv is shown in the inset. The dashed curves are the corresponding state-averaged results for the S_0 and the S_1 states.

TABLE 1: CASSCF-Optimized Structural Parameters and High-Correlated Energies of Relevant Stationary Points on the S_0 and the S_1 PESs of Azobenzene^a

species	$\nu_{\text{imaginary}}/$ $i \text{ cm}^{-1}$	ZPE/ kcal mol^{-1}	distance/ \AA		angle/deg			relative energy ^b / kcal mol^{-1}		
			$r(\text{NN})$	$r(\text{CN})$	$\theta(\text{CNN})$	$\delta(\text{CNCC})$	$\phi(\text{NNCC})$	$\Delta E(4-31\text{G})^c$	$\Delta E(\text{pVDZ})^d$	$\Delta E(\text{exp})^e$
$S_0(C_{2h}), ^1A_g$	f	130.1	1.253	1.428	116.8	180.0	0.0	0.0	0.0	0.0
S_1_FC	g	130.1 ^h	1.253	1.428	116.8	180.0	0.0	59.6	63.4	63.5
$S_1(C_{2h}), ^1B_g$	$50(b_u), 22(a_u)$	128.2	1.270	1.384	128.7	180.0	0.0	47.1	51.1	
$S_1(C_2), ^1B$	f	129.1	1.263	1.398	127.9	128.5	-5.7	47.7	50.1	~ 50
$S_1(C_s), ^1A''$	$21(a'')$	129.1	1.269	1.387	129.4	180.0	0.0	45.0	50.7	
				1.390	127.8		0.0			
$S_1_180(C_s), ^1A''$	$290(a''), 227(a'), 17(a'')$	127.7	1.249	1.400	127.3	i	0.0	61.0	68.1	
				1.362	180.0		i			
S_0/S_1 CI_rot	g	129.1 ^j	1.291	1.391	129.2	92.1	-0.4	42.2	47.8	
				1.421	120.2		0.9			
S_0/S_1 CI_inv	g	128.2 ^k	1.211	1.349	157.3	180.0	0.0	68.9	77.5	
				1.346	156.0		0.0			

^a Geometries and zero-point energies (ZPEs) were determined at the CAS(6,6)/4-31G level of theory. ^b Values are relative to the ground-state minimum, $S_0(C_{2h}), ^1A_g$, including the ZPE corrections. ^c CAS(6,6)/4-31G; the reference total energy is -568.30919 hartree. ^d CAS(6,6)/cc-pVDZ; the reference total energy is -569.18400 hartree. ^e Based on the absorption spectrum shown in ref 16 that the maximum S_1 absorption band locates at 450 nm and the tail of the absorption at ~ 560 nm. ^f All vibrational frequencies are positive values. ^g Vibrational frequency calculations were not performed. ^h ZPE is taken from the $S_0(C_{2h})$ species. ⁱ $\delta(\text{CNCC})$ or $\phi(\text{NNCC})$ is not defined for a linear CNN configuration. ^j ZPE is taken from the $S_1(C_2)$ species. ^k ZPE is taken from the $S_1(C_{2h})$ species.

(6,6)/cc-pVDZ level indicates that the energy difference between the two states is $21.2 \text{ kcal mol}^{-1}$ at this semilinear geometry. Such a large energy gap between the S_0 and the S_1 states is inconsistent with recent real-time observations that the electronic relaxation of azobenzene on the S_1 surface occurs on the subpicosecond to picosecond time scale.^{14–21,25} Furthermore, the force constant calculation at the optimized semilinear geometry (Table 1) gives three imaginary frequencies, indicating that the semilinear azobenzene corresponds to a high-order saddle point on the S_1 surface along the inversional pathway. This is another theoretical evidence for the photoisomerization of azobenzene following the in-plane inversion channel to be highly unfavorable.

Fujino and Tahara have recently reported on the NN stretch frequency observed in the transient Raman spectra to be close

to that of the ground state (1428 cm^{-1} vs 1440 cm^{-1}).¹⁹ This observation leads to a conclusion for the excited-state species having a double bond character that supports an inversion mechanism for photoisomerization of azobenzene. However, the NN bond order throughout the entire inversional pathway was calculated to be ~ 1.0 ,²³ which is inconsistent with the conclusion of the time-resolved Raman study.¹⁹ Our results support the argument of Ishikawa et al.²³ but disagree with the conclusion of Tahara and co-workers.^{19–21}

3.2. Rotation Channel. The potential energy curves shown in Figure 2 were obtained from scanning the CNCC torsional angle when the other structural parameters were optimized on the S_1 PES with C_2 symmetry. Once again, the results are similar to those of Cattaneo and Persico²² and Ishikawa et al.²³ and this topological picture may provide a theoretical description




Figure 4. Structure and the corresponding gradient difference vector (\mathbf{x}_1) and the derivatives coupling vector (\mathbf{x}_2) of the S_0/S_1 CI_inv obtained from the optimization at the state-averaged CAS(6,6)/4-31G level of theory. The right panel shows the plausible reaction paths on the S_0 PES after funneling through the S_0/S_1 CI_inv: Following the gradient difference vector \mathbf{x}_1 either a hot *trans*-azobenzene or the phenyl radical pair/ N_2 is produced; following the derivative coupling vector \mathbf{x}_2 the twisted *trans* isomer is formed.

for the rotation mechanism shown in Scheme 1. The molecule is planar with C_{2h} symmetry at the torsional angle of 180° , but the rotational motion along the CNNC torsional coordinate reduces the molecular symmetry to C_2 . Along the rotational reaction path, a local minimum, $S_1(C_2)$, is found at the torsional angle of 128.5° . According to our calculations, the S_1 surface along the rotational pathway is rather flat (Figure 2) and $S_1(C_2)$ is a true S_1 minimum with all the vibrational frequencies being positive; the frequency corresponding to the CNNC torsional motion is 15 cm^{-1} . Following the CNNC out-of-plane torsional reaction path from $S_1(C_2)$ toward the smaller torsional angle direction, the S_1 curve still remains flat but the S_0 curve increases drastically so that a surface touching is expected at a perpendicularly twisted configuration (torsional angle $\sim 90^\circ$) as in the case of azomethane recently reported.²⁴ According to the two-dimensional surface-scan calculation reported by Ishikawa et al.,²³ a conical intersection was evident at the location with the CNNC torsional angle of 88° and the CNN bending angle of 130° . In the present study, this conical intersection (S_0/S_1 CI_rot) has been fully optimized to locate at the CNNC torsional angle of 92.1° with two CNN bending angles equal to 129.2° and 120.2° , respectively; the optimized structure is shown in the inset of Figure 2 and its key structural parameters are summarized in Table 1. The structure of CI_rot has C_1 symmetry and the energy relative to the S_0 minimum is $47.8\text{ kcal mol}^{-1}$ predicted at the CAS(6,6)/cc-pVDZ level with the ZPE corrections according to the $S_1(C_2)$ species.

According to our calculated results, there are two important factors affecting the electronic relaxation through this rotation channel. First, there is no energy barrier involved along the rotational pathway from the Franck–Condon region (torsional angle = 180°) down to the $S_1(C_2)$ minimum area (torsional angle $\sim 130^\circ$). This feature provides an energy-favorable reaction path for the excited molecules to follow upon initial excitation to the $S_1(n,\pi^*)$ state. Second, the S_0/S_1 CI_rot can be easily reached at the middle point of the rotational pathway without involving a significant energy barrier. Funneling through the CI, different reaction pathways on the ground-state surface may be predicted by following the gradient difference vector (\mathbf{x}_1) or the derivative coupling vector (\mathbf{x}_2) direction.²⁸ The calculated results of azobenzene are similar to those of azomethane:²⁴ The \mathbf{x}_1 direction gives the guidance for a combined motion involving

the NN stretch, the asymmetric CN stretch and the asymmetric CNN bending modes whereas the \mathbf{x}_2 direction is mainly related to the CNNC torsional mode that gives the *trans* and the *cis* isomers as final products on the S_0 PES. Therefore, the rotational reaction pathway is not only energetically but also dynamically a favorable electronic relaxation channel for the photoisomerization of azobenzene to occur from the S_1 surface going through the S_0/S_1 CI_rot and eventually forming both the *trans*- S_0 and the *cis*- S_0 species via the \mathbf{x}_2 direction of the CI. When this channel is blocked by structural^{10,11} or environmental¹² modification, another relaxation pathway must be considered.

3.3. Concerted Inversion Channel. The potential energy curves shown in Figure 3 were obtained from synchronously scanning the two CNN bending angles when the other structural parameters on the S_1 PES with C_{2h} symmetry were optimized. The minimum-energy point of the S_1 curve, $S_1(C_{2h})$, is located above the ground-state minimum by $51.1\text{ kcal mol}^{-1}$ (including ZPE corrections) predicted at the CAS(6,6)/cc-pVDZ level with the two equivalent CNN bending angles equal to 128.7° (Table 1). Following the concerted CNN bending reaction path from $S_1(C_{2h})$ toward the larger bending angle direction, the S_1 and the S_0 potential energy curves come closer quickly and the two states tend to touch at bending angles greater than 148° . Further state-averaged CASSCF calculations have indicated that the two states interact strongly at larger angles and a surface crossing occurs at $\sim 157^\circ$; the results are shown as dashed curves in Figure 3. The attempt to optimize the geometry for the conical intersection (S_0/S_1 CI_inv) with C_{2h} symmetry was unsuccessful, but the geometry was successfully optimized without imposing a symmetry constraint. The optimized structure of S_0/S_1 CI_inv has C_s symmetry with two different CNN bending angles equal to 157.3° and 156.0° , respectively. The energy of S_0/S_1 CI_inv with respect to $S_1(C_2)$ is $21.2\text{ kcal mol}^{-1}$ calculated at the CAS(6,6)/4-31G level, but the relative energy increases to $27.4\text{ kcal mol}^{-1}$ when calculated at the CAS(6,6)/cc-pVDZ level of theory (Table 1). This theoretical finding suggests that another photochemical funnel (S_0/S_1 CI_inv) may be accessible along the concerted CNN bending coordinate when sufficient internal energy is provided upon excitation to the hot S_1 state or higher electronic excited states.

According to the results demonstrated in Figure 4, funneling through the S_0/S_1 CI_inv may lead to two different reaction

paths to be followed on the ground-state surface via either the gradient difference vector (\mathbf{x}_1) or the derivative coupling vector (\mathbf{x}_2) direction.²⁸ Basically, the \mathbf{x}_1 vector corresponds to a combined in-plane motion whereas the \mathbf{x}_2 vector relates to an out-of-plane bending motion. Following the \mathbf{x}_1 vector in the forward direction, the NN bond is stretched but the two CN bonds and the two CNN angles are simultaneously compressed and bent. Because breaking the NN double bond on the S_0 PES requires overcoming a substantial energy barrier (112.6 kcal mol⁻¹),²⁹ we expect the molecule would become vibrationally hot at the trans- S_0 configuration. If the reverse \mathbf{x}_1 vector direction is followed, the NN bond is compressed but the two CN bonds are stretched together with the two CNN bending angles bent in a concerted manner. This motion allows a concerted CN bond-breaking process to occur on the S_0 PES if the available internal energy is sufficient (65.8 kcal mol⁻¹).²⁹ On the other hand, following both forward and reverse directions of \mathbf{x}_2 , the concerted CNN out-of-plane bending motion becomes active and the two N atoms are moving in the opposite direction perpendicular to the original molecular plane to form a hot twisted trans- S_0 species. Even though this hot trans- S_0 species has the CNNC torsional angle of $\sim 180^\circ$, its two phenyl rings are perpendicularly twisted with respect to the new molecular plane; i.e., the two NNCC torsional angles are $\sim 90^\circ$. Therefore, we expect that eventually a planar trans- S_0 isomer would be formed via twisting the two phenyl rings on the ground-state surface because the twisted species is unstable and it corresponds to a second-order saddle point on the S_0 PES with the energy higher than the trans- S_0 minimum by 12.2 kcal mol⁻¹.²⁹

Early experimental measurements indicate that the trans-to-cis photoisomerization quantum yield is ~ 0.25 on S_1 excitation but it drops to only ~ 0.1 on S_2 excitation.⁷ The quantum yields are independent of the wavelengths on both excitations with a critical transition point occurring at ~ 370 nm.⁷ The reduction of the quantum yield on S_2 excitation has been proposed to be due to the opening of a new relaxation channel that produces only the trans- S_0 isomer.^{19–21} On the basis of our theoretical analysis aforementioned, following the concerted CNN bending reaction path (concerted inversion channel) to funnel through the S_0/S_1 CI_{inv} would give either a phenyl radical pair and a N₂ molecule or a hot *trans*-azobenzene on the ground-state surface. Because the production of a cis isomer is less favorable for the electronic relaxation along this channel, the trans-to-cis quantum yield of azobenzene is expected to be lower if this channel is open upon excitation. Furthermore, the calculated relative energy of the S_0/S_1 CI_{inv} is 77.5 kcal mol⁻¹ (Table 1), which is in excellent agreement with the transition point being observed at ~ 370 nm (~ 77 kcal mol⁻¹ above S_0 minimum). These theoretical findings suggest that the new electronic relaxation channel may be the concerted CNN bending reaction path with a photochemical funnel (S_0/S_1 CI_{inv}) located at the geometry close to a linear CNNC configuration. Therefore, our results may provide a theoretical foundation to rationalize the early steady-state observations: The rotation channel is dominant on S_1 excitation to give a higher trans-to-cis quantum yields whereas the concerted inversion channel is involved for lower values being observed when this channel is open upon excitations with sufficient energy (i.e., $\lambda < 370$ nm).

When the rotation channel is blocked by putting the molecule in a cyclodextrin cavity, only this new relaxation channel is feasible to give more trans isomers so that the values of the photoisomerization yields obtained from both the S_2 and the S_1 excitations become equally small (~ 0.13).¹² The rotation-


blocked azobenzene derivatives via chemical modification also show equal photoisomerization yields from the two excitations but with higher values (~ 0.25).^{10,11} Higher trans-to-cis quantum yields means more cis isomers were produced via the new relaxation channel. For a rotation-blocked azobenzene derivative, the two phenyl rings are structurally restricted so that they cannot twist freely to form more stable trans- S_0 isomers when following both forward and reverse \mathbf{x}_2 directions of the S_0/S_1 CI_{inv} along the new relaxation pathway. Thus, relatively more cis- S_0 isomers will be produced in the case of rotation-blocked azobenzene derivatives^{10,11} than in the case of azobenzene–cyclodextrin inclusion complexes.¹²

Further experimental evidence for the involvement of the S_0/S_1 CI_{inv} was found in recent time-resolved studies, where the azobenzene derivative (methyl orange) in different cyclodextrin environments was investigated using the ultrafast transient lens technique.^{30,31} The trans-to-cis isomerization quantum yield of methyl orange on S_2 excitation ($\lambda = 400$ nm)³² was determined to be 0.1 in pure water, and the yield remains the same in the 1:1 methyl orange/ α -cyclodextrin inclusion complex. This is because only the dimethylaniline moiety of the molecule is included in the nanocavity and the azo functional group is outside the cavity so that it can still move freely; either via the CNNC torsional motion or through the concerted CNN bending motion. However, the quantum yield only reduces to 0.07 in the 1:2 complex where the molecule was completely included in two α -cyclodextrins and the rotational motion about the NN double bond was completely inhibited. This experimental evidence indicates that the rotational reaction pathway is only a minor photoisomerization channel to be considered upon S_2 excitation of an azo compound. The observed subpicosecond to picosecond relaxation times in the inclusion complexes support the mechanism that the electronic relaxation of the azobenzene derivative mainly proceeds through the concerted inversion channel with the involvement of the S_0/S_1 CI_{inv} as an efficient photochemical funnel. The observed quantum yield of 0.07 in the 1:2 inclusion complex may be rationalized by the fact that the azo group inside the nanocavity still has room to adjust its position to produce some cis isomers on the hot ground-state PES, and this is the reason the yield in 1:1 complex is very similar to the yield (0.1) in 1:1 complex and in aqueous solution. On the other hand, in a 2:2 complex when two molecules were included in parallel in two γ -cyclodextrins, not only the observed time constants were remarkably elongated but also the quantum yield was further reduced to zero. The interaction between the two monomers should be considered in this case and this could effectively hinder the motion of the azo group along the concerted inversion channel. If this channel is only partly followed, the electronic relaxation may occur at the early stage without the involvement of the photochemical funnel (S_0/S_1 CI_{inv}), which is the result for the much slower relaxation rates being observed. Moreover, the two CNN bending angles do not have to change from the Franck–Condon structure too much, which eventually leads to the production of only the trans isomers as final products being observed.

4. Conclusion

Our results for the photoisomerization mechanism of *trans*-azobenzene from the S_1 state down to the S_0 surface are summarized in Scheme 2. According to the present study, the widely accepted inversion channel has been *ruled out* because a substantial energy barrier is involved and the energy gap between the two electronic states is too large to allow an efficient

SCHEME 2



electronic relaxation to occur. On the other hand, a photochemical funnel (S_0/S_1 CI_rot) is located on the midpoint of the rotational pathway that produces both trans and cis isomers on the S_0 PES. Therefore, the conventional rotation channel is responsible for a cold S_1 species to proceed with zero or little available internal energy. Upon excitation to the S_2 state, a hot S_1 species is produced so that the new relaxation channel can be open via the concerted CNN bending motion to reach another photochemical funnel (S_0/S_1 CI_inv). Funneling through the S_0/S_1 CI_inv gives either a trans isomer or the other bond-breaking side products, which may be the reason for the observed trans-to-cis photoisomerization quantum yields on S_2 excitation being much lower than the S_1 excitation. When the rotation channel is blocked by chemical modification^{10,11} or an inclusion complex,¹² only the new relaxation channel is feasible so that the values of the photoisomerization yields obtained from both the S_2 and the S_1 excitations become equal. The present study provides solid theoretical evidence to address the issue for the long-standing controversy on the photoisomerization mechanism of azobenzene. The new relaxation channel, hitherto unreported, is intrinsically an inversion channel involving the concerted CNN bending motion in the same molecular plane to approach a linear CNNC configuration to reach the conical intersection for an efficient electronic relaxation. We notice that the semilinear (inversion), the skew (rotation), and the linear (concerted inversion) isomerization channels are three typical photochemical processes being considered for azomethane since the 1970s.³³ For azobenzene, the photochemical behavior of the concerted inversion channel has not been discussed until now, which indeed has led to the rotation-inversion controversy on the photoisomerization mechanism of azobenzene and its derivatives for more than 20 years.

Acknowledgment. This work was supported by the National Science Council of Republic of China with the project contract number 90-2119-M-009-001. I thank Prof. M. C. Lin, Prof. Wen-Sheng Chung, Prof. Robert S. H. Liu, and Prof. Maurizio Persico for many helpful discussions. The suggestions and comments made by the reviewers are much appreciated.

References and Notes

(1) Berg, R. H.; Hvilsted, S.; Ramanujam, P. S. *Nature* **1996**, *383*, 505–508.

- (2) Ikeda, T.; Tsutsumi, O. *Science* **1995**, *168*, 1873–1875.
- (3) Willner, I.; Rubin, S.; Riklin, A. *J. Am. Chem. Soc.* **1991**, *113*, 3321–3325.
- (4) Liu, Z. F.; Hashimoto, K.; Hujishima, A. *Nature* **1990**, *347*, 658–660.
- (5) Sudesh, G.; Neckers, D. C. *Chem. Rev.* **1989**, *89*, 1915–1925.
- (6) Tamai, N.; Miyasaka, H. *Chem. Rev.* **2000**, *100*, 1875–1890.
- (7) Rau, H. In *Photochromism: Molecules and Systems*; Durr, H., Bouas-Laurent, H., Eds.; Elsevier: Amsterdam, 1990; pp 165–192 and references therein.
- (8) Zimmerman, G.; Chow, L.-Y.; Paik, U.-J. *J. Chem. Phys.* **1958**, *80*, 3528–3531.
- (9) Bortollus, P.; Monti, S. *J. Phys. Chem.* **1979**, *83*, 648–652.
- (10) Rau, H.; Lüddecke, E. *J. Am. Chem. Soc.* **1982**, *104*, 1616–1620.
- (11) Rau, H. *J. Photochem.* **1984**, *26*, 221–225.
- (12) Bortollus, P.; Monti, S. *J. Phys. Chem.* **1987**, *91*, 5046–5050.
- (13) Monti, S.; Orlandi, G.; Palmieri, P. *Chem. Phys.* **1982**, *71*, 87–99.
- (14) Lednev, I. K.; Ye, T.-Q.; Hester, R. E.; Moore, J. N. *J. Phys. Chem.* **1996**, *100*, 13338–13341.
- (15) Nägele, T.; Hoche, R.; Zinth W.; Wachtveitl, J. *Chem. Phys. Lett.* **1997**, *272*, 489–495.
- (16) Lednev, I. K.; Ye, T.-Q.; Matousek, P.; Towrie, M.; Fogg, P.; Neuwahl, F. V. R.; Umapathy, S.; Hester, R. E.; Moore, J. N. *Chem. Phys. Lett.* **1998**, *290*, 68–74.
- (17) Lednev, I. K.; Ye, T.-Q.; Abbott, L. C.; Hester, R. E.; Moore, J. N. *J. Phys. Chem. A* **1998**, *102*, 9161–9166.
- (18) Hirose, Y.; Yui, H.; Sawada, T. *J. Phys. Chem. A* **2002**, *106*, 3067–3071.
- (19) Fujino, T.; Tahara, T. *J. Phys. Chem. A* **2000**, *104*, 4203–4210.
- (20) Fujino, T.; Arzhantsev, S. Y.; Tahara, T. *J. Phys. Chem. A* **2001**, *105*, 8123–8129.
- (21) Fujino, T.; Arzhantsev, S. Y.; Tahara, T. *Bull. Chem. Soc. Jpn.* **2002**, *75*, 1031–1040.
- (22) Cattaneo, P.; Persico, M. *Phys. Chem. Chem. Phys.* **1999**, *1*, 4739–4743.
- (23) Ishikawa, T.; Noro, T.; Shoda, T. *J. Chem. Phys.* **2001**, *115*, 7503–7512.
- (24) Diao, E. W.-G.; Zewail, A. H. *Chem. Phys. Chem.* **2003**, *4*, 445–456 and references therein.
- (25) Lu, Y.-C.; Chang, C.-W.; Diao, E. W.-G. *J. Chin. Chem. Soc.* **2002**, *49*, 693–701.
- (26) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; et al. *Gaussian98*, revision A.11; Gaussian, Inc.: Pittsburgh, PA, 1998.
- (27) Based on the similar CAS orbitals, the total energies of the $S_1(C_s)$ and the $S_1(C_{2h})$ species are -568.23581 and -568.23121 hartree, respectively, calculated at the CAS(6,6)/4-31G level of theory. However, due to the convergence problem for describing the entire rotational reaction path, the $S_1(C_2)$ species was calculated at the same level using the slightly different CAS orbitals, which gives -568.23152 hartree. Force-constant calculations indicate that both the $S_1(C_s)$ and the $S_1(C_{2h})$ species involve imaginary frequencies whereas the $S_1(C_2)$ species is a true local minimum with all frequencies being positive (Table 1).
- (28) Bernardi, F.; Olivucci, M.; Robb, M. A. *Chem. Soc. Rev.* **1996**, *25*, 321–328, and references therein.
- (29) Based on the results calculated at the UB3LYP/6-31G(d) level of theory with zero-point energy correction at the same level.
- (30) Takei, M.; Yui, H.; Hirose, Y.; Sawada, T. *J. Phys. Chem. A* **2001**, *105*, 11395–11399.
- (31) Yui, H.; Takei, M.; Hirose, Y.; Sawada, T. *Rev. Sci. Instrum.* **2003**, *74*, 907–909.
- (32) Deffering from the case of *trans*-azobenzene, the S_2 state of methyl orange is hit by the 400 nm excitation due to a substantial red shift of the S_2 band in absorption spectrum upon substitution of azobenzene with the dimethyl amino group.
- (33) Camp, R. N.; Epstein, I. R.; Steel, C. *J. Am. Chem. Soc.* **1977**, *99*, 2453–2459.