

Table of contents

Chapter 1 General Introduction	1
1.1 Luminescence	1
1.2 Fluorescence lifetime	2
1.3 Phase-modulation and pulse excitation method.....	3
1.4 Pulse excitation techniques	4
1.5 Summary	6
Chapter 2 Experimental setup	10
2.1 Introduction.....	10
2.2 Ultrashort pulse formation: principle of mode locking.....	10
2.3 Active mode-locking and passive mode-locking	14
2.4 Birefringent filter	16
2.5 Femtosecond Ti:sapphire laser: Mira 900-D.....	17
2.6 Autocorrelator (Mini, Inrad)	17
2.7 Fluorescence optically gated system.....	19
2.8 Time-Correlated Single Photon Counting (TCSPC) system.....	20
2.8.1 Double subtractive monochromator.....	22
2.8.2 Constant fractional discriminator (CFD)	23
2.8.3 Time-to-amplitude converter (TAC)	24
2.8.4 Analog-to-digital converter (ADC)	24
2.8.5 Multi-channel analyzer (MCA).....	25
2.8.6 Variable delay box.....	25
2.8.7 Microchannel-plate photomultiplier Tube (MCP PMT).....	25
Chapter 3 Relaxation Dynamics of PPB nanoparticles.....	26
3.1 Introduction.....	26
3.2 Experimental Section	29
3.3 Results and discussion	31
3.3.1 Formation of PPB nanoparticles	31
3.3.2 Steady-state absorption/emission spectra	33
3.3.3 Picosecond relaxation dynamics of PPB nanoparticles	36
3.3.4 Molecular structures of PPB in solid state and in solution	43
3.3.5 Femtosecond relaxation dynamics of free PPB and DSB molecules in THF	48
3.3.6 Femtosecond relaxation dynamic of PPB and DSB nanoparticles	51
3.4 Concluding Remarks.....	52
Chapter 4 Relaxation Dynamics of CNDSB Nanobelts	54
4.1 Introduction.....	54

4.2 Experimental Section	55
4.3 Results and Discussion	56
4.3.1 Formation of CN-DSB Nanostructures.....	56
4.3.2 Steady-state Absorption/Emission Spectra	59
4.3.3 Picosecond Relaxation Dynamics of CNDSB nanobelts.....	61
4.3.4 Femtosecond Relaxation dynamics of CNDSB in THF solution and nanobelts	65
4.4 Concluding Remarks.....	71
Chapter 5 Hydration Dynamics of BMVC in organic solvent.....	73
5.1 Introduction.....	73
5.2 Experimental section.....	75
5.3 Results and Discussion	76
5.3.1 Steady state spectra	76
5.3.2 Excited state dynamics of BMVC in tris-buffer	77
5.3.3 Solvation dynamics of BMVC in tris-buffer	81
5.3.4 Excited state dynamics and solvation dynamics of BMVC in glycerol	84
5.4 Concluding Remarks.....	88
Chapter 6 Hydration dynamics of BMVC in duplex and quadruplex DNA.....	89
6.1 Introduction.....	89
6.2 Experimental section.....	91
6.3 Results and discussion	92
6.3.1 Steady state spectra	92
6.3.2 Time-resolved fluorescence measurement.....	93
6.3.3 Fluorescence anisotropy measurement	95
6.3.4 Hydration dynamics of BMVC/LD complex.....	97
6.3.5 Hydration dynamics of BMVC/Hum 24 complex	100
6.4 Concluding Remarks.....	102

Table of Figures

Figure 1.1: Possible non-radiative processes that compete with the fluorescence emission	2
Figure 1.2: The principle of time-domain method for measuring fluorescence lifetime	4
Figure 1.3: Scheme layout of the principle of fluorescence up-conversion	6
Figure 2.1: The scheme diagram of a simple laser cavity.....	10
Figure 2.2: The imaginary rectangular spectra in laser cavity.	11
Figure 2.3: The illustration of the laser intensity change versus time for N=4.	13
Figure 2.4: The scheme diagram for cross-sectional view of CW wave and a mode-locked pulse. (A) Slit is open; the loss of a CW wave and a mode-locked pulse is the same. (B) Slit is closed; the loss of a CW wave increases, and only the mode-locked pulse can survive	15
Figure 2.5: The optical scheme for Coherent Mira-900 D oscillator.	18
Figure 2.6: The optical layout of femtosecond autocorrelator.....	18
Figure 2.7 :The optical scheme for femtosecond fluorescence optical gated system. M1~M7: dielectric mirror (700~1000 nm), BS:dichroic beamsplitter, B:berck polarization compensator, P1 and P2:parabolic mirror, S: rotating sample cell.	20
Figure 2.8: The optical layout of TCSPC system	21
Figure 2.9: Data acquisition and timing diagram of TCSPC module	22
Figure 2.10: Basic function of CFD, a small portion of the input signal is inverted, and the arriving time is determined by the zero cross point.	23
Figure 3.1: SEM images of PPB nanocrystals showing the sizes in average diameters of (A) 50, (B) 150, (C) 300 nm and (D) $\sim \mu$ m obtained from nanoparticle's suspension containing 65, 70, 75 and 80% volume fractions of water in THF, respectively.	32
Figure 3.2: (A) UV-Visible absorption spectra changes of PPB depend on the water fractions in THF. (B) PL spectra changes of PPB depend on the water fractions in THF.	34
Figure 3.3: (A) Picosecond fluorescence transients of PPB shown in log scale are obtained from the excitation at $\lambda_{ex} = 310$ nm and the probe at $\lambda_{fl} = 420$ nm at various water/THF mixtures as indicated. (B) Picosecond fluorescence transients of PPB are shown in linear scale at 0, 65, 70, 75 and 80% solutions. The inset shows the corresponding transients in the 1 ns scale.	37
Figure 3.4: Picosecond fluorescence transients of PPB at 75% solution with $\lambda_{ex} = 310$ nm and $\lambda_{fl}/nm =$ (A) 390, (B) 450 and (C) 510. The data are fitted according to the dual-emissive model (Scheme 2). The solid black curves are theoretical fits	

with residues shown as the green traces. The τ_1 , τ_2 and τ_3 correspond to the A and B, and A' in Scheme 2, respectively.	38
Figure 3.5: Three fitted temporal coefficients of PPB, (A) τ_1 , (B) τ_2 , and (C) τ_3 . the corresponding fitted parameters are summarized in Table 1. The dashed curve shown in (c) represents the wavelength-dependent single-exponential-decay coefficient of the PPB single crystal.	41
Figure 3.6: (A) Powder XRD pattern of the PPB bulk material. (B) Powder XRD patterns of the colloid PPB samples from various water/THF mixtures as indicated.	44
Figure 3.7: (A) Powder XRD pattern of the PPB bulk material. (B) Powder XRD patterns of the colloid PPB samples from various water/THF mixtures as indicated.	45
Figure 3.8: The ground state structures of (A) PPB and (B) DSB optimized at the B3LYP/6-31G(d) level of theory. The relevant torsional angles are indicated.	47
Figure 3.9: Femtosecond fluorescence transients of (A) PPB and (B) DSB in pure THF obtained from $\lambda_{ex} = 355$ nm and $\lambda_{fl} = 450$ nm. The transient of PPB was fitted according to a combined consecutive kinetic model, $A' \xrightarrow{\tau_1} B'$ and $A \xrightarrow{\tau_2} B \xrightarrow{\tau_3} C$, while that of DSB by a simple consecutive model, $A \xrightarrow{\tau_1} B \xrightarrow{\tau_2} C$. The solid black curves are theoretical fits with residues shown as green traces; the blue and magenta curves under each transient are the deconvoluted components corresponding to A and B, respectively; dark yellow curve shown under the transient of PPB represents A'.	49
Figure 3.10: Femtosecond transients of (A) PPB and (B) DSB nanoparticles in 75% solution. The transient was obtained from $\lambda_{ex}=360$ nm and $\lambda_{em}=450$ nm. Both transient can be fitted by a bi-exponential function. The fast component (blue) decays ~ 45 ps, and the slow component (magenta) persists in 0~100 ps region.	51
Figure 4.1: SEM images showing the evolution of CN-DSB nanostructures obtained at different water additions and at concentrations of 3.6×10^{-5} M viz. (A) 70, (B) 80, (C) 90% to THF, (D) 1.8×10^{-5} M for 80% solution, (E) 7.2×10^{-5} M for 80%, and for the (F) CN-DSB bulk crystals, respectively.	58
Figure 4.2: (A) UV-visible spectra of CNDSB in various water/THF mixtures, the absorbance of PMMA and single crystal film were also indicated as red and olive symbols, respectively. (B) PL spectra of CNDSB in various percentages of water/THF solutions, PMMA thin film, and single crystal film.	59
Figure 4.3: Picosecond time-resolved spectra of CNDSB in various water/THF mixtures, PMMA thin film, and single crystal obtained at $\lambda_{ex}=375$ nm, $\lambda_{em}=500$ nm. The inset was the corresponding transients in shorter time scale.	62

- Figure 4.4: Picosecond transients of CNDSB in (A) 0%, (B) 70%, (C) 80%, (D) 90%, (E) single crystal, (F) PMMA film, each transients can be fitted with a bi-exponential function. The solid black curves are theoretical fits with residues shown as green traces; the short component was indicated as magenta line, and long component was indicated as dark cyan. The blue trace in 70% solution indicates the contribution from nonaggregate CNDSB molecules. 64
- Figure 4.5: Femtosecond time-resolved spectra of CNDSB in (A) THF and (B) PMMA thin film. In each transient, the solid black curves are theoretical fits and the residues were shown as green traces. For THF solution, the transient was obtained at $\lambda_{ex}=375$ nm, $\lambda_{em}=480$ nm, and deconvoluted with bi-exponential function. The short component and long component were indicated as blue and dark cyan, respectively. For PMMA films, the transient was obtained at $\lambda_{ex}=375$ nm and $\lambda_{em}=460$ nm, and deconvoluted with bi-exponential function. The long component (dark cyan line) was fixed at the value obtaining from TCSPC technique. 67
- Figure 4.6: Femtosecond transients of CNDSB in 80% Water/THF solution, the excitation wavelength was fixed at 375 nm and probe at (A) 460 nm, (B) 500nm, (C) 550 nm. For each transient, at least three components were required to get the satisfactory fitting. The longest component (magenta line) was fixed at the value obtained from TCSPC result. 68
- Figure 4.7: (a) The blank test of CNDSB at 80% water/THF solution: The solid black line indicated the signal of CNDSB nanobelts, and the red line was the transient of 80% water/THF solution. (b) $I_{//}$ and I_{\perp} indicates the relative polarization of the collection fluorescence is parallel or perpendicular relative to the polarization of excitation beam. The anisotropy decay $r(t)$ was calculated using the following equation: $r(t) = \frac{I_{//} - I_{\perp}}{I_{//} + 2I_{\perp}}$ 70
- Figure 5.1: The steady state UV and PL spectra of BMVC in tris-buffer and glycerol. The concentration of each sample was fixed at $5 \times 10^{-6} M$, and the excitation wavelength was fixed at 430 nm. 75
- Figure 5.2: Time resolved fluorescence spectra of BMVC dissolved in tris-buffer and glycerol. The TCSPC result is shown in the main panel, and the up-conversion results in the inset. The fitting time-coefficients of each transient are shown as indicated. Figure A and B are the transients of BMVC in tris-buffer; Figure C and D are the transients of BMVC in glycerol. 78
- Figure 5.3: (A) Femtosecond time resolved spectra of BMVC in tris-buffer with $\lambda_{ex}=430$ nm, and probing wavelength varied from 510 nm to 640 nm. (B) Femtosecond time resolved spectra of BMVC in glycerol with $\lambda_{ex}=430$ nm, and

probing wavelength varied from 530 nm to 640 nm.....	83
Figure 5.4: (A) The reconstructed time-resolved emission spectra of BMVC in tris-buffer. The row data at each delay time were shown as scatter and the fitted curves as solid curves. Steady state spectrum was indicated at dash line. (B) The C(t) value of BMVC in tris-buffer. Time coefficients were obtained by fitted the curve with bi-exponential model. The two components were indicated as blue and dark cyan curves, and the residual was indicated as green line.	84
Figure 5.5: (a) The reconstructed time-resolved emission spectra of BMVC in glycerol. The row data at each delay time and the fitted curves were shown as scatter and solid curves, respectively. Steady state spectrum was indicated at dash line. (b) The C (t) value of BMVC in glycerol. The time coefficient was obtained by fitted the curve with bi-exponential model. The two components were indicated as blue and dark cyan curves, and the residual was indicated as green line.....	87
Figure 6.1: Steady state spectrum of BMVC and BMVC interacted with LD and Hum24 DNA. The concentration is fixed at 5×10^{-6} M, and the excitation wavelength was indicated as gray dash line.	93
Figure 6.2: The femtosecond time-resolved spectra of BMVC in tris-buffer, and interacted with LD and Hum24 DNA. The excitation was fixed at 430 nm, and the probing wavelength was fixed at 590 nm	94
Figure 6.3: Fluorescence anisotropy decay of BMVC at various environments. The anisotropy was reconstructed using the TCSPC results, and the instrument response function is indicated as blue line. (A) BMVC in tris-buffer; (B) BMVC+Hum24; (C) BMVC+LD.....	96
Figure 6.4: Femtosecond time-resolved anisotropy decay of (A)BMVC in tris-buffer, (B) BMVC/Hum24 and (C)BMVC/LD complex.	98
Figure 6.5: (A) The fluorescence transient of BMVC/ LD complex probing at 540 nm, 600nm, and 640 nm. (B) The constructed time-resolved emission spectrum of BMVC/ LD complex at various delay time. (C) The corresponding solvent relaxation spectra of BMVC/ LD complex.....	99
Figure 6.6: (A) The fluorescence transient of BMVC/ Hum24 complex probing at 540 nm, 600nm, and 640 nm. (B) The constructed time-resolved emission spectrum of BMVC/ Hum24 complex at various delay time. (C) The corresponding solvent relaxation spectra of BMVC/ Hum 24 complex.....	101