

List of Figures

Figure 1.1 Electron and hole mobilities versus temperature in ultrapure naphthalene.....	04
Figure 1.2 The temperature dependence of the field effect mobility of pentacene and tetracene molecular crystals.....	05
Figure 1.3 The transport and luminescence characteristics of the first organic electroluminescent device.....	06
Figure 1.4 Tang and Van Slyke's organic light emitting device.....	08
Figure 1.5 Tang's photovoltaic cell is built on a glass substrate coated with transparent indium tin oxide.....	09
Figure 1.6 The electroluminescent device structure used to demonstrate emission from the organic dyes shown above.....	10
Figure 1.7 Organic active matrix displays, courtesy of Steven Van Slyke and Kodak.....	11
Figure 2.1 The molecule makes a transition from the ground vibrational state to the state with a vibrational wavefunction with the highest overlap with the initial vibrational wavefunction.....	25
Figure 2.2 Dynamics of exciton formation.....	27
Figure 2.3 Energy level diagram of an emissive molecule with its first coordination sphere of matrix molecules.....	30
Figure 2.4 The diagram explains the effect of triplet harvesting.....	33
Figure 2.5 The relative positions of the first excited singlet and triplet levels in a typical molecule.....	35
Figure 3.1. Comparison of the UV-Vis absorption spectra of five cyclometalated	

iridium complexes in CH ₂ Cl ₂	59
Figure 3.2 Comparison of solution PL spectra for six triplet Ir (III)-based complexes in CH ₂ Cl ₂	60
Figure 3.3 ORTEP drawing of (CF ₃ -bo) ₂ Ir(acac).....	63
Figure 3.4 EL device structures with (a) CF ₃ -, (b) Me- substituted, and (c) parent complexes as a dopant.....	65
Figure 3.5 EL performance of three (x/y-bo) ₂ Ir(acac) doped EL devices.....	68
Figure 4.1 Comparison of solution PL spectra for (L' ^X) ₂ Ir(acac) complexes at room temperature in CH ₂ Cl ₂	80
Figure 4.2 Temperature dependences of PL lifetimes for (ppl) ₂ Ir(acac).....	84
Figure 4.3 EL device structures with (ppl) ₂ Ir(acac), (4OMeppl) ₂ Ir(acac), and Ir(4Fppl) ₃ complexes as dopants.....	86
Figure 4.4 Comparison of EL spectra for devices.....	86
Figure 5.1 EL device structures with Ir(4-CF ₃ bt) ₂ (acac):CBP and Ir(2-phq) ₂ acac :CBP complexes as dopants.....	94
Figure 5.2 The absorption bands of [Ir(4-CF ₃ bt) ₂ (acac)] complex.....	98
Figure 5.3 ORTEP drawing of (CF ₃ -bt) ₂ Ir(acac).....	99
Figure 5.4 PL quantum efficiency η_{PL} vs dopant concentration in thin film samples of Ir(4-CF ₃ bt) ₂ (acac):CBP and Ir(2-phq) ₂ acac :CBP.....	102
Figure 5.5 Dopant concentration dependence of vs yield and EQE of Ir(4- CF ₃ bt) ₂ (acac): CBP film.....	103
Figure 5.6 Temperature dependence of decay lifetime and PL intensity for Ir(4-CF ₃ bt) ₂ (acac).....	104
Figure 5.7 EL spectra of devices a, b, c, d and e fabricated with different dopant contents of Ir(4-CF ₃ bt) ₂ (acac) complex.....	106

Figure 6.1 The device structure for Device-1.....	117
Figure 6.2 EL spectra and performance of devices 1-11.....	120
Figure 6.3 The PL solution for complexes Ir(2-phq) ₃ and Ir(4-CF ₃ bt) ₂ (acac)...	122
Figure 6.4 External quantum efficiency versus current density for WOLEDs.	123
Figure 6.5 External quantum efficiency versus current density for WOLEDs.	124
Figure 7.1 Comparison of the UV-Vis absorption spectra of (G1-Ir-bt ₃) and Ir(bt) ₃ complexes in CH ₂ Cl ₂	144
Figure 7.2 Comparison of solution PL spectra for (G1-Ir- bt ₃) and Ir(bt) ₃ complexes in CH ₂ Cl ₂	144
Figure 7.3 The PLE spectra of Ir-G1-bt ₃ dendrimer in CH ₂ Cl ₂ at room Tempature.....	146
Figure 7.4 EL device structure with the (G1-bt ₃ Ir) and (bt) ₃ Ir as a dopant.....	147
Figure 7.5 Currentdensity dependance of luminance (a), luminance efficiency (b), and EQE (c) for (Ir-G1-bt ₃) and Ir(bt) ₃	148
Figure 7.6 Comparison of EL spectra for devices based on (G1-bt ₃ Ir) and (bt) ₃ Ir complexes.....	149
Figure 7.7 Comparison of EL spectra for (a) G1 -Ir(ppy) ₃ and Ir(ppy) ₃ complexes and (b) G1 -Ir(Fppy) ₃ and Ir(Fppy) ₃ complexes.....	152
Figure 7.8 Comparison of (a) Efficiency versus luminance, showing the relatively efficiencies, (b) EQE vs current density, and (c) voltage vs current density for Ir(ppy) ₃ and G1 -Ir(ppy) ₃	154
Figure 7.9 (a) voltage vs current density (b) EQE vs current density (c) luminance vs current density (d) luminance vs current density about G1 -Ir(Fppy) ₃	156

List of Tables

Table 3.1. Spectroscopic, redox and photophysical properties for all Ir complexes synthesized in this research.....	61
Table 3.2 Selected Bond lengths [Å] for (CF ₃ -bo) ₂ Ir(acac).....	63
Table 3.3 Crystal data and structure refinement for (CF ₃ bo) ₂ Ir(acac).....	64
Table 3.4 Comparative study of the electroluminescent performance of three EL devices based on (bo) ₂ Ir(acac), (bo-Me) ₂ Ir(acac) an(bo-CF ₃) ₂ Ir(acac) as a dopant.....	67
Table 4.1 Crystal data and structure refinement for Ir(III) 2-(4-fluorophenyl)-1-pyrroline complex.....	81
Table 4.2 Comparison of selected bond distances (Å) and bond angles(deg) for Ir(III) 2-(4-fluorophenyl)-1-pyrroline complex.....	82
Table 4.3 EL performance of devices D-a, D-b, and D-c fabricated with (4-x-ppl) ₂ Ir(acac) complexes (x = H, OMe, F).....	87
Table 5.1 Crystal Data and Structure Refinement for (CF ₃ bt) ₂ Ir(acac).....	100
Table 5.2 Selected Bond Distances [Å] and Angles [°] for (CF ₃ bt) ₂ Ir(acac)..	101
Table 5.3 EL Performance of devices a , b , c , d and e fabricated with different content of Ir(4-CF ₃ bt) ₂ (acac) complex.....	105
Table 6.1 All devices structure of WOLEDs.....	116
Table 6.2 Comparison of EL performance for white light devices 1-11.....	121
Table 7.1 Comparative study of the electroluminescent properties of three EL devices based on (G1-bt ₃ Ir) and (bt) ₃ Ir as a dopant. Device structure: ITO/PEDOT(35 nm)/ dopant /TPBI (30 nm)/MgAg (100 nm).....	150
Table 7.2 Comparative study of the electroluminescent properties of three EL devices based on Ir(ppy) ₃ and G1 -Ir(ppy) ₃ as a dopant. Device	

structure: ITO/PEDOT(35 nm)/ dopant /TPBI (30 nm)/MgAg (100 nm).....155

Table 7.3 Comparative study of the EL properties of two devices based on

Ir(Fppy)₃ and **G1**-Ir(Fppy)₃ as a dopant. Device structure:

ITO/PEDOT(35 nm)/ dopant /TPBI (30 nm)/MgAg (100 nm).....157



List of Acronyms

Alq₃ tris(8-hydroxyquinoline) aluminum(III)

BCP 9-dimethyl-4,7-diphenyl-1, 10-phenanthroline

CBP 4, 4'-bis(9-carbazolyl)-1,1'-biphenyl

CDBP 4,4'-bis(9-carbazolyl)-2,2'-dimethyl-biphenyl

CF color filter

CIE Commission Internationale d'Eclairage

CRTs cathode ray tubes

CuPc copper phthalocyanine

CV cyclicvoltametric

DI deionized

DMCT dopant-to-matrix charge transfer

EL electroluminescent

EML emitting material Layer

ETL electron transport layer

EQE external quantum efficiency

FIr6 bis[4',6'-difluoro- phenylpyridinato)tetrakis(1-pyrazolyl)borate]

FWHM full width at half maximum

HBL hole and exciton blocking layer

HOMO highest occupied molecular orbital

HTL hole transport layer

IQE internal quantum efficiency

Ir-complex iridium (III) complex

ISC intersystem crossing



ITO indium-tin-oxide

I-V current-voltage

LCDs liquid crystal displays

LED light emitting device

LUMO lowest unoccupied molecular orbital

mCP 1,3-Bis(carbazol-9-yl)benzene

MLCT metal to ligand charge transfer

NMP N-Methyl-2-Pyrrolidone

NPB 4,4'-bis[(1-naphthyl)(phenyl)amino]-1, 1'-biphenyl

α -NPD 4,4'-bis[N-(1-naphthyl)-N-phenyl-amino]biphenyl

OLED organic light emitting devices

PDAs potential applications

PEDOT polyethylene dioxythiophene

PL Photoluminescence

PLQY photoluminescence quantum yield

PVK polyvinylcarbazole

QE quantum efficiency

QY quantum yield

RGB red, green and blue

SOC spin-orbit coupling

TBP Tri-n-butylphosphine

TCTA 4,4',4''-tris(N-carbazolyl)triphenylamine

TPD N, N'-diphenyl-N, N'-di(3-methylphenyl)-1,1'-biphenyl-4,4'-diamine

UV ultra-violet

UV-VIS ultraviolet-visible

WOLEDs white organic light-emitting devices

