

References

- Ayad A, Boot-Handford RP, Humphries MJ, Kadler KE, Shuttleworth CA, 1994. The extracellular matrix facts book. San Diego, CA: Academic Press, Inc. p 104.
- Beanes SR, Dang C, Soo C, Ting K, 2003. Skin repair and scar formation: the central role of TGF- β . *Exp Rev Mol Med* 5:1-22.
- Bertone C, Sgro LC, 1982. Clinical data on topical application in gynaecology of polydeoxiribonucleotide of human placenta. *Int J Tissue React* 4:165-167.
- Bigliardi P, 1982. Treatment of acute radiodermatitis of first and second degree with semi-greasy placenta ointment. *Int J Tiss Reac* 4: 153-154.
- Birkedal-Hansen H, Taylor RE, 1982. Detergent-activation of latent collagenase and resolution of its component molecules. *Biochem Biophys Res Commun* 107:1173-1178.
- Breier G, Damert A, Plate KH, Risau W, 1997. Angiogenesis in embryos and ischemic diseases. *Thromb Haemost* 78:678-683.
- Carmeliet P, 2003. Angiogenesis in health and disease. *Nat Med* 9:653-660.
- Cass DL, Meuli M, Adzick NS, 1997. Scar wars: Implications of fetal wound healing for the pediatric burn patient. *Pediatr Surg Intern* 12:484-489.
- Chomczynski P, Sacchi N, 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate–phenol–chloroform extraction. *Anal Biochem* 162:156-159.
- Clark RAF ed. The Molecular and Cellular Biology of Wound Repair. 2nd ed, London, Plenum Press: 1996.
- Creemers EE, Cleutjens JP, Smits JF, Daemen MJ, 2001. Matrix metalloproteinase inhibition after myocardial infarction: a new approach to prevent heart failure? *Circ Res* 89:201-210.
- Dang CM, Beanies SR, Lee H, Zhang X, Soo C, Ting K, 2003. Scarless fetal wounds are associated with an increased matrix metalloproteinase-to-tissue-derived inhibitor of metalloproteinase ratio. *Plast Reconstr Surg* 111:2273-85.

Deitch EA, Wheelaan TM, Rose MP, Clothier J, Cotter J, 1983. Hypertrophic burn scars: analysis of variables. *J Trauma* 23:895-898.

Diegelmann RF, Evans MC, 2004. Wound healing: an overview of acute, fibrotic and delayed healing. *Front Biosci* 9:283-289.

Eickelberg O, 2001. Endless healing: TGF-beta, SMADs, and fibrosis. *FEBS Lett* 506:11-14.

Falanga V, 1992. Growth factors and chronic wounds: the need to understand the microenvironment. *J Dermatol* 19:667-672.

Falanga V, Grinnell F, Gilchrest B, Maddox YT, Moshell A, 1994. Workshop on the pathogenesis of chronic wounds. *J Invest Dermatol* 102, 125-127.

Feugate JE, Wong L, Li QJ, Martins-Green M, 2002. The CXC chemokine cCAF stimulates precocious deposition of ECM molecules by wound fibroblasts, accelerating development of granulation tissue. *BMC Cell Biology* 3:13-27.

Florkiewicz RZ and Sommer A, 1989. Human basic fibroblast growth factor gene encodes four polypeptides: three initiate translation from non-AUG codons. *Proc Natl Acad Sci USA* 86:3978-3981.

Folkman J, 1991. What is the evidence that tumors are angiogenesis-dependent? *J Natl Cancer Inst* 82:4-6.

Folkman J, 1992. The role of angiogenesis in tumor growth. *Semin Cancer Biology* 3: 65-71.

Folkman J, and M Klagsbrun, 1987. Angiogenic factors. *Science* 235:442-447.

Folkman J, Shing T, 1992. Angiogenesis. *J Biol Chem* 267:10931-10934.

Fukai F, Ohtaki M, Fujii N, Yajima H, Ishii T, Nishizawa Y, Miyazaki K, Katayama T, 1995. Release of biological activities from quiescent fibronectin by a conformational change and limited proteolysis by matrix metalloproteinases. *Biochemistry* 34:11453-11459.

Gailit J, Clark RA, 1994. Wound repair in the context of extracellular matrix. *Curr Opin Cell Biol* 6:717-725.

Grazul-Bilska AT, Johnson ML, Bilski JJ, Redmer DA, Reynolds LP, Abdullah A, Abdullah KM, 2003. Wound healing: the role of growth factors. *Drugs Today (Barc)*. 2003 39:787-800.

Griffith LG, 2000. Polymeric biomaterials. *Acta Mater* 48:263-277.

Hauser GA., 1982. Placental extract injections in the treatment of loss of hair in women. *Int J Tissue React* 4:159-163.

Higley HR, Ksander GA, Gerhardt CO, Falanga V, 1995. Extravasation of macromolecules and possible trapping of transforming growth factor-beta in venous ulceration. *Br J Dermatol* 132:79-85.

Holvoet S, Vincent C, Schmitt D, Serres M, 2003. The inhibition of MAPK pathway is correlated with down-regulation of MMP-9 secretion induced by TNF- α in human keratinocytes. *Experimental Cell Research* 290:108-119.

Liota LA, Stetler-Stevenson WG, 1990. *Cancer Biology* 1:96-106.

Lorenz HP, Adzick NS, 1993. Scarless skin wound repair in the fetus. *West J Med* 159:350-355.

Mallya SK, Mookhtiar KA, Gao Y, Brew K, Dioszegi M, Birkedal-Hansen H, Van Wart HE, 1990. Characterization of 58-kilodalton human neutrophil collagenase: Comparison with human fibroblast collagenase. *Biochemistry* 29: 10628-10634.

Marchese C, Chedid M, Dirsch OR, Csaky KG, Santanelli F, Latini C, LaRochelle WJ, Torrisi MR, Aaronson SA, 1995. Modulation of keratinocyte growth factor and its receptor in reepithelializing human skin. *J Exp Med* 182:1369-1376.

Martin P, 1997. Wound healing: aiming for perfect skin regeneration. *Science* 276:75-81.

Marx M, Perlmutter RA, Madri JA, 1994. Modulation of PDGF-receptor expression in microvascular endothelial cells during in vitro angiogenesis. *J Clin Invest* 93:131-139.

Matrisian LM, 1992. The matrix-degrading metalloproteinases. *Bioassays* 14:455-463.

Mauviel A, 1993. Cytokine regulation of metalloproteinase gene expression. *Cell Biochem* 53:288-295.

Muralidhar RV and Panda T, 1999. Useful products from human placenta. *Bioprocess Enginerring* 20:23-25

Muratore O, Pesce Schito A, Cattarini G, Tonoli EL, Gianoglio S, Schiappacasse S, Felli L, Picchetta F, Schito GC, 1997. Evaluation of the trophic effect of human placental polydeoxyribonucleotide on human knee skin fibroblasts in primary culture. *Cell Mol Life Sci* 53:279-285.

Nagase H, 1997. Activation mechanisms of matrix metalloproteinases. *Biol Chem* 378: 151-160.

Nanney LB, 1990. Epidermal and dermal effects of epidermal growth factor during wound repair. *J Invest Dermatol* 94:624-629.



Neely AN, Clendening CE, Gardner J, Greenhalgh DG, Warden GD, 1999. Gelatinase activity in keloids and hypertrophic scars. *Wound Repair Regen* 7:166-171.

Odorisio T, Schietroma C, Zaccaria ML, Cianfarani F, Tiveron C, Tatangelo L, Failla CM, Zambruno G, 2002. Mice overexpression placenta growth factor exhibit increased vascularization and vessel permeability. *J Cell Sci* 115:2559-2567.

Orecchia G, Donadini G and Boschetti L, 1984. Estratto di placenta: attualita' del loro impiego in dermatologia. *Chron Derm* 5: 687-698.

Postlethwaite AE, Keski-Oja J, Moses HL, Kang AH, 1987. Stimulation of the chemotactic migration of human fibroblasts by transforming growth factor-beta. *J Exp Med* 165: 251-256.

Priestley GC, 1993. An introduction to the skin and its diseases. In Molecular Aspects of Dermatology. Priestley GC, editor. John Wiley and Sons Ltd., Chichester, UK. 1-17.

Raghav R, 1994. The role of extracellular matrix in postinflammatory wound healing and fibrosis. *FASEB J* 8: 823-831.

Ravanti L, Kähäri VM, 2000. Matrix metalloproteinases in wound repair. *Int J Mol Med* 6: 391-407.

Rees RS, Robson MC, Smiell JM, Perry BH, 1997. Pressure Ulcer Study Group. Bepacupermin gel in the treatment of pressure ulcers. *Wound Repair Regen* 7:141-147.

Reynolds LP, Redmer DA, 1995. Utero-placental vascular development and placental fuction. *J Anim Sci* 73:1839-1851.

Risau W, 1997. Mechanisms of angiogenesis. *Nature* 386:671-674.

Rosenthal M, 1982. The application of an extract of human placenta in the treatment of rheumatic affections. *Int J Tissue React* 4:147-51.

Samuels P, Tan AK, 1999. Fetal scarless wound healing. *J Otolaryngol* 28: 296-302.

Seghezzi G, Patel S, Ren CJ, Gualandris A, Pintucci G, Robbins ES, Shapiro RL, Galloway AC, Rifkin DB, Mignatti P, 1998. Fibroblast growth factor-2 (FGF-2) induces vascular endothelial growth factor (VEGF) expression in the endothelial cells of forming capillaries: an autocrine mechanism contributing to angiogenesis. *J Cell Biol* 141:1659-1673.

Stocum DL. Wound repair, regeneration and articial tissues. Georgetown, TX: R. G. Landes, 1995

Wahl SM, Hunt DA, Wakefield LM, McCartney-Francis N, Wahl LM, Roberts AB, Sporn MB, 1987. Transforming growth factor-beta (TGF-beta) induces monocyte chemotaxis and growthfactor production. *Proc Natl Acad Sci USA* 84: 5788-5792.

Watson AL, Palmer ME, Burton GJ, 1996. An in vitro model for the study of wound healing in first trimester human placenta. *Cell Tissue Res* 286:431-438.

Saunders WB, 1992. Wound healing: Biochemical and chemical aspects. Philadelphia: 96-112.

Wells A, 2000. Tumor invasion: role of growth factor-induced cell moility. *Adv Cancer Res* 78:31-101.

Werb Z, 1989. Proteinases and matrix degradation. In: Kelly WN, Harris ED, Ruddy S, Sledge CB, editors. Textbook of rheumatology, 3rd, ed. Philadelphia: WB Saunders 300-321.

Woessner JF Jr, 1995. Quantification of matrix metalloproteinases in tissue sample. *Meth Enzymol* 248:510-528.

Wu CH, Chang GY, Chang WC, Hsu CT, Chen RS, 2003. Wound healing effects of Porcine placental extracts on rats with thermal injury. *British J Dermatol* 148:236-245.

Vigano` M, Klinger M, Colonna M, Blandini D, De Angelis A and Crabai P, 1990. Utilizzo di un Polidesossiribonucleotide (PDRN) nelle ustioni di grado intermedio. *Giorn Ital Ric Clin Terap* 11: 5-9

JP patent number: JP58177918, 1983. Inhibitor for melanin formation by placental-decomposition product.

JP patent number: JP6234798, 1994. Cell metabolism activator extracted from bovine placenta.

