



ELSEVIER

Contents lists available at ScienceDirect

Journal of Differential Equations

www.elsevier.com/locate/jde



Traveling wavefronts for time-delayed reaction–diffusion equation: (I) Local nonlinearity

Ming Mei^{a,b,*}, Chi-Kun Lin^c, Chi-Tien Lin^d, Joseph W.-H. So^e

^a Department of Mathematics, Champlain College Saint-Lambert, Saint-Lambert, QC, J4P 3P2, Canada

^b Department of Mathematics and Statistics, McGill University, Montreal, QC, H3G 1M8, Canada

^c Department of Applied Mathematics, National Chiao Tung University, Hsinchu 30010, Taiwan, ROC

^d Department of Applied Mathematics, Providence University, Taichung 43301, Taiwan, ROC

^e Department of Mathematical and Statistical Sciences, University of Alberta, Edmonton, AB, T6G 2G1, Canada

ARTICLE INFO

Article history:

Received 26 October 2008

Revised 16 December 2008

Available online 5 February 2009

MSC:

35K57

34K20

92D25

Keywords:

Reaction–diffusion equation

Time-delay

Traveling waves

Stability

ABSTRACT

In this paper, we study a class of time-delayed reaction–diffusion equation with local nonlinearity for the birth rate. For all wavefronts with the speed $c > c_*$, where $c_* > 0$ is the critical wave speed, we prove that these wavefronts are asymptotically stable, when the initial perturbation around the traveling waves decays exponentially as $x \rightarrow -\infty$, but the initial perturbation can be arbitrarily large in other locations. This essentially improves the stability results obtained by Mei, So, Li and Shen [M. Mei, J.W.-H. So, M.Y. Li, S.S.P. Shen, Asymptotic stability of traveling waves for the Nicholson's blowflies equation with diffusion, Proc. Roy. Soc. Edinburgh Sect. A 134 (2004) 579–594] for the speed $c > 2\sqrt{D_m(\varepsilon p - d_m)}$ with small initial perturbation and by Lin and Mei [C.-K. Lin, M. Mei, On travelling wavefronts of the Nicholson's blowflies equations with diffusion, submitted for publication] for $c > c_*$ with sufficiently small delay time $r \approx 0$. The approach adopted in this paper is the technical weighted energy method used in [M. Mei, J.W.-H. So, M.Y. Li, S.S.P. Shen, Asymptotic stability of traveling waves for the Nicholson's blowflies equation with diffusion, Proc. Roy. Soc. Edinburgh Sect. A 134 (2004) 579–594], but inspired by Gourley [S.A. Gourley, Linear stability of travelling fronts in an age-structured reaction–diffusion population model, Quart. J. Mech. Appl. Math. 58 (2005) 257–268] and based on the property of the critical wavefronts, the weight function is carefully

* Corresponding author at: Department of Mathematics, Champlain College Saint-Lambert, Saint-Lambert, QC, J4P 3P2, Canada.

E-mail addresses: mmei@champlaincollege.qc.ca, mei@math.mcgill.ca (M. Mei), cklin@math.nctu.edu.tw (C.-K. Lin), ctlm@pu.edu.tw (C.-T. Lin), joseph.so@ualberta.ca (J.W.-H. So).

selected and it plays a key role in proving the stability for any $c > c_*$ and for an arbitrary time-delay $r > 0$.

© 2009 Elsevier Inc. All rights reserved.

1. Introduction

In this series of study, we consider a class of time-delayed reaction–diffusion equation arising from the population dynamics (see, for example [27])

$$\frac{\partial v}{\partial t} - D_m \frac{\partial^2 v}{\partial x^2} + d_m v = \varepsilon \int_{-\infty}^{\infty} b(v(t-r, x-y)) f_{\alpha}(y) dy, \quad t > 0, \quad x \in R, \tag{1.1}$$

with the initial condition

$$v(s, x) = v_0(s, x), \quad s \in [-r, 0], \quad x \in R, \tag{1.2}$$

which describes the population distribution of single species with age-structure and spatial diffusion. Here, $v(t, x)$ denotes the total mature population at time t and location x . $D_m > 0$ is the diffusion rate of the mature population in space, $d_m > 0$ is the death rate of the mature population, and $r > 0$ denotes the maturation age. Let $d_i(z)$ and $D_i(z)$ be the death rate and the spatial diffusion rate for the immature population for age $z \in [0, r]$, respectively. Define $\varepsilon > 0$ and $\alpha \geq 0$ as follows:

$$\varepsilon = e^{-\int_0^r d_i(z) dz} \quad \text{and} \quad \alpha = \int_0^r D_i(z) dz. \tag{1.3}$$

They represent the impact of the death rate of the immature and the effect of the dispersal rate of the immature, respectively. If the mature population move more effectively than the immature, i.e. $D_i(z) \leq D_m$, then, according to (1.3) for α , we have

$$\alpha \leq rD_m. \tag{1.4}$$

The function $b(v(t, x))$ is the birth function. As we know, the reaction term in a reaction–diffusion equation, which is some what like the birth function in our equation (1.1), plays an important role to depict the character of the equation. One interesting type of such a function is the Fisher–KPP’s reaction function $b(v) = v(1 - v)$ with bi-stable nodes $v = 0$ and $v = 1$ as the equilibria (see [4, 10,24,25]). The related nonlinear wave stability for the time-delayed reaction–diffusion equation was studied by Smith and Zhao [25] by the upper–lower solution method. The other interesting type is the reaction term with two nodes as the equilibria, but one is stable and the other one is unstable, for example, the Nicholson’s blowflies birth rate $b(v) = pve^{-av}$ (see [6,8,11,14,15,20,21,26–29,33,34,37], etc.). The related wave stability was then investigated in [15,20,21] by the technical weighted energy method, because, as mentioned therein, the upper–lower solution method is defective in this case. For such kind birth functions with one stable node and one unstable node, in this paper, as considered in [14], we take the birth function as

$$b_1(v) = pve^{-av^q} \quad \text{or} \quad b_2(v) = \frac{pv}{1 + av^q}, \tag{1.5}$$

where $p > 0$, $q > 0$ and $a > 0$ are constants. Here, $b_1(v)$, as an exponential type, is a generalization of the Nicholson’s blowflies birth rate as considered before, and $b_2(v)$ is the kind of important rational-type function. Note that, for $b_1(v)$ with $q = 1$, it is the well-known Nicholson’s blowflies

birth function, and the corresponding equation (1.1) is the so-called Nicholson's blowflies equation with diffusion. Lastly, $f_\alpha(y)$ is the heat kernel

$$f_\alpha(y) = \frac{1}{\sqrt{4\pi\alpha}} e^{-\frac{y^2}{2\alpha}} \quad \text{with} \quad \int_{-\infty}^{\infty} f_\alpha(y) dy = 1 \quad \text{for all } \alpha > 0. \tag{1.6}$$

When $\alpha \rightarrow 0$, namely $D_i(z) \approx 0$, which means that the immature population is almost non-mobile, by the property of the heat kernel we have

$$\lim_{\alpha \rightarrow 0} \int_{-\infty}^{\infty} b(v(t-r, x-y)) f_\alpha(y) dy = b(v(t-r, x)).$$

In this case, the nonlocal equation (1.1) is reduced to a local time-delayed reaction-diffusion equation for $\alpha = 0$:

$$\frac{\partial v}{\partial t} - D_m \frac{\partial^2 v}{\partial x^2} + d_m v = \varepsilon b(v(t-r, x)). \tag{1.7}$$

The models (1.1) and (1.7) as well as related equations have been extensively studied recently, see [1–31] and [33–37] and the references therein. The existence of traveling wavefronts for the local equation (1.7) and the nonlocal equation (1.1) were obtained by So and Zou [29], So, Wu and Zou [26] and Liang and Wu [14] using the method of upper-lower solutions. The boundedness and the asymptotic behavior for the critical wave speed with respect to the delay time r as well as the diffusion rate D_m were analyzed by Wu, Wei and Mei in [33,34]. Furthermore, the stability of the traveling wavefronts was showed by Mei, So, Li and Shen [21], Mei and So [20] and recently in Lin and Mei [15]. For the local equation (1.7) with $b(v) = b_1(v)$ and $q = 1$, i.e. the Nicholson's blowflies equation, by using the weighted energy method, Mei, So, Li and Shen [21] proved that the wavefronts are stable for the wave speed $c > 2\sqrt{D_m(\varepsilon p - d_m)}$ with a sufficiently small initial perturbation, but the case when $c_* < c \leq 2\sqrt{D_m(\varepsilon p - d_m)}$ was left open, where $c_* > 0$ is the critical wave speed. Later by using the comparison principle together with the weighted energy method, Lin and Mei [15] further improved it to $c > c_*$ for the delay time $r \ll 1$, because $c_* \approx 2\sqrt{D_m(\varepsilon p - d_m)}$ as $r \ll 1$. The initial perturbation is also improved to be arbitrarily large in a weighted Sobolev space by the comparison principle and the squeeze technique. For the nonlocal equation (1.1) with $b(v) = b_1(v)$ and $q = 1$, the stability was obtained in [20] with a much strong restriction on the wave speed $c > 2\sqrt{D_m(3\varepsilon p - 2d_m)}$. The main goal in the present paper and our second paper [18] is to study the stability of all traveling wavefronts with $c > c_*$ and arbitrary delay time $r > 0$. In this paper, we focus on the local equation (1.7) and prove the stability of all wavefronts with arbitrary delay time r . Our approach is still the technical weighted energy method together with the comparison principle. But, inspired by Gourley [5] (see also an extension to the nonlinear stability by Li, Mei and Wong [11]) and based on the property of the critical wavefront speed, by selecting an ideal weight function and carefully taking the energy estimates, we obtain the stability for all waves including those slower waves. No restriction is needed for the delay time r and the speed c . For the nonlocal equation (1.1), the improved stability for all wavefronts with $c > c_*$ is the content of our second paper [18], where some new techniques are introduced to overcome the difficulty of the energy estimates caused by the nonlocal integral term.

For research related to other models of population dynamics, we refer to [1–3,5,6,11–13,16] and the references therein. See also the review articles [7,8] and the textbooks [32,35,36].

Notations. Throughout the paper, $C > 0$ denotes a generic constant, while $C_i > 0$ ($i = 0, 1, 2, \dots$) represents a specific constant. Let I be an interval, typically $I = \mathbb{R}$. $L^2(I)$ is the space of the square integrable functions defined on I , and $H^k(I)$ ($k \geq 0$) is the Sobolev space of the L^2 -functions $f(x)$

defined on the interval I whose derivatives $\frac{d^i}{dx^i} f$ ($i = 1, \dots, k$) also belong to $L^2(I)$. $L^2_w(I)$ denotes the weighted L^2 -space with a weight function $w(x) > 0$ and its norm is defined by

$$\|f\|_{L^2_w} = \left(\int_I w(x) |f(x)|^2 dx \right)^{1/2}.$$

$H^k_w(I)$ is the weighted Sobolev space with the norm given by

$$\|f\|_{H^k_w} = \left(\sum_{i=0}^k \int_I w(x) \left| \frac{d^i}{dx^i} f(x) \right|^2 dx \right)^{1/2}.$$

Let $T > 0$ be a number and \mathcal{B} be a Banach space. We denote by $C^0([0, T]; \mathcal{B})$ the space of the \mathcal{B} -valued continuous functions on $[0, T]$. $L^2([0, T]; \mathcal{B})$ as the space of the \mathcal{B} -valued L^2 -functions on $[0, T]$. The corresponding spaces of the \mathcal{B} -valued functions on $[0, \infty)$ are defined similarly.

The rest of the paper is organized as follows. In Section 2, we introduce the travelling wavefronts, their properties and state the nonlinear stability theorem. In Section 3, after establishing the comparison principle and some key energy estimates in the weighted Sobolev spaces, we prove the nonlinear stability result.

2. Traveling wavefronts and stability theorem

It is easily seen that Eq. (1.7) has two constant equilibria v_{\pm} , where

$$\text{for } b_1(v): \quad v_- = 0 \quad \text{and} \quad v_+ = \left(\frac{1}{a} \ln \frac{\varepsilon p}{d_m} \right)^{1/q}, \tag{2.1}$$

$$\text{for } b_2(v): \quad v_- = 0 \quad \text{and} \quad v_+ = \left(\frac{\varepsilon p - d_m}{ad_m} \right)^{1/q}. \tag{2.2}$$

A traveling wavefront of (1.7) connecting with v_{\pm} is a monotone solution $v(t, x) = \phi(x + ct)$ satisfying

$$\begin{cases} c\phi'(\xi) - D_m\phi''(\xi) + d_m\phi(\xi) = \varepsilon b(\phi(\xi - cr)), \\ \phi(\pm\infty) = v_{\pm}, \end{cases} \tag{2.3}$$

where $\xi = x + ct$ and $' = \frac{d}{d\xi}$.

By using the upper-lower solutions method as in [14,26,29], we can, similarly, prove the existence of the traveling wavefronts.

Proposition 2.1 (Existence of traveling wavefronts). Assume that

$$\text{for } b(v) = b_1(v): \quad 1 < \frac{\varepsilon p}{d_m} \leq e^{1/q}, \tag{2.4}$$

$$\begin{aligned} \text{for } b(v) = b_2(v): \quad &\text{either } 1 < \frac{\varepsilon p}{d_m} \leq \frac{q}{q-1} \quad \text{if } q > 1, \\ &\text{or } 1 < \frac{\varepsilon p}{d_m} < \infty \quad \text{if } 0 < q \leq 1. \end{aligned} \tag{2.5}$$




Fig. 2.1. The graphs of $F_c(\lambda)$ and $G_c(\lambda)$ for $c = c_*$ and $c > c_*$, respectively.

Then there exist a minimum speed $c_* = c_*(r, \varepsilon, D_m, d_m, p) > 0$ and a corresponding number $\lambda_* = \lambda(c_*) > 0$ satisfying

$$F_{c_*}(\lambda_*) = G_{c_*}(\lambda_*), \quad F'_{c_*}(\lambda_*) = G'_{c_*}(\lambda_*), \tag{2.6}$$

where

$$F_c(\lambda) = \varepsilon p e^{-\lambda c r}, \quad G_c(\lambda) = c\lambda - D_m \lambda^2 + d_m, \tag{2.7}$$

and (c_*, λ_*) is the tangent point of $F_c(\lambda)$ and $G_c(\lambda)$, such that for all $c > c_*$, the travelling wavefront $\phi(x + ct)$ of Eq. (1.7) connecting v_{\pm} exists uniquely (up to a shift).

The graphs of $F_c(\lambda)$ and $G_c(\lambda)$ are shown in Fig. 2.1 for $c = c_*$ and $c > c_*$, respectively.

From the graphs, we can easily see that $F_c(\lambda_*) = G_c(\lambda_*)$ for $c = c_*$ and $F_c(\lambda_*) < G_c(\lambda_*)$ for $c > c_*$, that is,

$$\varepsilon p e^{-\lambda_* c_* r} = c_* \lambda_* - D_m \lambda_*^2 + d_m, \quad \text{for } c = c_*, \tag{2.8}$$

$$\varepsilon p e^{-\lambda_* c r} < c \lambda_* - D_m \lambda_*^2 + d_m, \quad \text{for } c > c_*. \tag{2.9}$$

By a straightforward calculation, it can be verified that the wavefront $\phi(\xi)$ decays to the constant state v_- as follows

$$|\phi(\xi) - v_-| = O(1)e^{-\lambda_1 |\xi|}, \quad \text{as } \xi \rightarrow -\infty, \tag{2.10}$$

where $\lambda_1 = \lambda_1(c)$ is a c -depending function as showed in Fig. 2.1. It is also noted that

$$\varepsilon p e^{-\lambda c r} < c\lambda - D_m \lambda^2 + d_m, \quad \text{for } c > c_*, \lambda \in (\lambda_1, \lambda_*]. \tag{2.11}$$

Thus, we define a weight function as

$$w(x) = \begin{cases} e^{-2\lambda(x-x_0-cr)}, & \text{for } x \leq x_0 + cr, \\ 1, & \text{for } x > x_0 + cr, \end{cases} \tag{2.12}$$

where λ is a positive number between $\lambda_1 = \lambda_1(c)$ and $\lambda_* = \lambda_*(c_*)$, i.e., $\lambda_1 < \lambda \leq \lambda_*$ (cf. Fig. 2.1), and the number x_0 is chosen to be sufficiently large such that (3.31) holds, i.e.,

$$b'(\phi(x_0 - cr)) < \frac{\min\{\lambda c + d_m - D_m \lambda^2, d_m\}}{\varepsilon \cosh(\lambda cr)}.$$

We can now state our main theorem.

Theorem 2.2 (Nonlinear stability). *Let $\frac{\varepsilon p}{d_m}$ satisfy (2.4) for $b(v) = b_1(v)$, or (2.5) for $b(v) = b_2(v)$, and let $b'(v_+)$ be sufficiently small. For any given wavefront $\phi(x + ct)$ with the speed $c > c_*$, if the initial data satisfy*

$$v_- \leq v_0(s, x) \leq v_+, \quad \text{for } (s, x) \in [-r, 0] \times \mathbb{R}, \tag{2.13}$$

and the initial perturbation $v_0(s, x) - \phi(x + cs)$ is in $C([-r, 0]; H^1_w(\mathbb{R}))$, then the solution of (1.7) and (1.2) satisfies

$$v_- \leq v(t, x) \leq v_+, \quad \text{for } (t, x) \in \mathbb{R}_+ \times \mathbb{R}, \tag{2.14}$$

and

$$v(t, x) - \phi(x + ct) \in C([0, \infty); H^1_w(\mathbb{R})). \tag{2.15}$$

In particular, the solution $v(t, x)$ converges to the wavefront $\phi(x + ct)$ exponentially in time, that is,

$$\sup_{x \in \mathbb{R}} |v(t, x) - \phi(x + ct)| \leq C e^{-\mu t}, \quad t \geq 0, \tag{2.16}$$

for some positive number μ .

Remark 1.

- As showed in Theorem 2.2, the wavefront stability holds for all wavefronts $\phi(x + ct)$ with $c > c_*$, and for all delay time $r > 0$, which essentially improves the previous stability result for the wavefronts with speed $c > 2\sqrt{D_m - \varepsilon p}$ in [21]. In another word, when the initial perturbation around the wavefront at $x \rightarrow -\infty$ is sufficiently small, the solution of (1.7) and (1.2) can be sufficiently approximated by this specified traveling wavefront, which is very important and significant in the physical application, as well as the issue of numerical simulation.
- As the asymptotical profile, the wavefront $\phi(x + ct)$ satisfying $v_0(s, x) - \phi(x + cs) \in H^1_w(\mathbb{R})$ is uniquely selected. In fact, from the definition of the weight function (2.12) and the weighted Sobolev's space $H^1_w(\mathbb{R})$, we have

$$f \in H^1_w \Leftrightarrow \sqrt{w(x)}f \in H^1 \hookrightarrow C(\mathbb{R}),$$

see also a proof in [19]. Thus, the condition $v_0(s, x) - \phi(x + cs) \in H^1_w(\mathbb{R})$ implies that

$$|v_0(s, x) - \phi(x + cs)| = O(1)e^{-\lambda|x|}, \quad \text{as } x \rightarrow -\infty, \quad s \in [-r, 0], \tag{2.17}$$

for $\lambda \in (\lambda_1, \lambda_*]$, while, from (2.11), we know that, for any two wavefronts $\phi(x + ct)$ and $\phi(x + ct + x_1)$, where x_1 is a shift, the difference is

$$|\phi(x + cs) - \phi(x + cs + x_1)| = O(1)e^{-\lambda_1|x|}, \quad \text{as } x \rightarrow -\infty, \quad s \in [-r, 0]. \tag{2.18}$$

Since $\lambda > \lambda_1$, i.e., the decay of $|v_0(s, x) - \phi(x + cs)|$ is much faster than $|\phi(x + cs) - \phi(x + cs + x_1)|$, the condition (2.17) determines a unique wavefront, which is the asymptotical profile of the original solution $v(t, x)$. In another word, once the initial perturbation around the wavefront $\phi(x + cs)$ at $x \rightarrow -\infty$ decays faster than $O(e^{-\lambda_1|x|})$, then the original solution $v(t, x)$ to Eqs. (1.7) and (1.2) behaves exactly as this traveling wavefront $\phi(x + ct)$.

3. Regarding the condition $b'(v_+) \ll 1$, actually it is natural. In fact, we can easily check that, for $b(v) = b_1(v)$,

$$b'_1(v_+) = \frac{d_m}{\varepsilon} \left[1 - \ln \left(\frac{\varepsilon p}{d_m} \right)^q \right] \rightarrow 0, \quad \text{as } \frac{\varepsilon p}{d_m} \rightarrow e^{1/q}.$$

Thus,

$$b'_1(v_+) = 0, \quad \text{for } \frac{\varepsilon p}{d_m} = e^{1/q}.$$

In this case, the wavefront strength $|v_+ - v_-|$ is the largest, and the wavefront is called the strongest wave. So we prove the stability for the strong wavefronts.

Similarly, for $b(v) = b_2(v_+)$, we have

$$b'_2(v_+) = \frac{d_m^2}{\varepsilon^2 p} \left[1 - (q - 1) \frac{\varepsilon p - d_m}{d_m} \right] \rightarrow 0, \quad \text{as } \frac{\varepsilon p}{d_m} \rightarrow \frac{q}{q - 1}, \quad \text{for } q > 1,$$

and thus

$$b'_2(v_+) = 0, \quad \text{for } \frac{\varepsilon p}{d_m} = \frac{q}{q - 1}, \quad q > 1.$$

3. Proof of main theorem

First of all, we state the following two lemmas which are given in [15].

Lemma 3.1 (Boundedness). *Let*

$$v_- = 0 \leq v_0(s, x) \leq v_+, \quad \text{for } (s, x) \in [-r, 0] \times R. \tag{3.1}$$

Then the solution $v(t, x)$ of the Cauchy problem (1.7) and (1.2) satisfies

$$v_- \leq v(t, x) \leq v_+, \quad \text{for } (t, x) \in [0, \infty) \times R. \tag{3.2}$$

Lemma 3.2 (Comparison principle). *Let $\bar{v}(t, x)$ and $\underline{v}(t, x)$ be the solutions of (1.7) and (1.2) with the initial data $\bar{v}_0(s, x)$ and $\underline{v}_0(s, x)$, respectively. If*

$$v_- \leq \underline{v}_0(s, x) \leq \bar{v}_0(s, x) \leq v_+, \quad \text{for } (s, x) \in [-r, 0] \times R, \tag{3.3}$$

then

$$v_- \leq \underline{v}(t, x) \leq \bar{v}(t, x) \leq v_+, \quad \text{for } (t, x) \in R_+ \times R. \tag{3.4}$$

As in [11,15], we will use the comparison principle and the weighed energy method to prove Theorem 2.2. Let the initial data $v_0(s, x)$ be such that $v_- \leq v_0(s, x) \leq v_+$ for $(s, x) \in [-r, 0] \times R$, and define

$$\begin{cases} V_0^+(s, x) = \max\{v_0(s, x), \phi(x + cs)\}, \\ V_0^-(s, x) = \min\{v_0(s, x), \phi(x + cs)\}, \end{cases} \text{ for } (s, x) \in [-r, 0] \times R. \tag{3.5}$$

Obviously,

$$v_- \leq V_0^-(s, x) \leq v_0(s, x) \leq V_0^+(s, x) \leq v_+, \text{ for } (s, x) \in [-r, 0] \times R, \tag{3.6}$$

$$v_- \leq V_0^-(s, x) \leq \phi(x + cs) \leq V_0^+(s, x) \leq v_+, \text{ for } (s, x) \in [-r, 0] \times R. \tag{3.7}$$

Let $V^+(t, x)$ and $V^-(t, x)$ be the corresponding solutions of (1.7) and (1.2) with the initial data $V_0^+(s, x)$ and $V_0^-(s, x)$ respectively, i.e.,

$$\begin{cases} \frac{\partial V^\pm}{\partial t} - D_m \frac{\partial^2 V^\pm}{\partial x^2} + d_m V^\pm = \varepsilon b(V^\pm(t - r, x)), & (t, x) \in R_+ \times R, \\ V^\pm(s, x) = V_0^\pm(s, x), & x \in R, \quad s \in [-r, 0]. \end{cases} \tag{3.8}$$

By the comparison principle (Lemma 3.2), we have

$$v_- \leq V^-(t, x) \leq v(t, x) \leq V^+(t, x) \leq v_+, \text{ for } (t, x) \in R_+ \times R, \tag{3.9}$$

$$v_- \leq V^-(t, x) \leq \phi(x + ct) \leq V^+(t, x) \leq v_+, \text{ for } (t, x) \in R_+ \times R. \tag{3.10}$$

We will now prove the stability result (Theorem 2.2) in three steps.

Step 1. The convergence of $V^+(t, x)$ to $\phi(x + ct)$.

Let $\xi := x + ct$ and

$$u(t, \xi) := V^+(t, x) - \phi(x + ct), \quad u_0(s, \xi) := V_0^+(s, x) - \phi(x + cs). \tag{3.11}$$

Then by (3.10) and (3.7), we have

$$u(t, \xi) \geq 0, \quad u_0(s, \xi) \geq 0. \tag{3.12}$$

From Eq. (1.7), $u(t, \xi)$ satisfies

$$\begin{cases} \frac{\partial u}{\partial t} + c \frac{\partial u}{\partial \xi} - D_m \frac{\partial^2 u}{\partial \xi^2} + d_m u - \varepsilon b'(\phi(\xi - cr))u(t - r, \xi - cr) \\ = \varepsilon Q(t - r, \xi - cr), & (t, \xi) \in R_+ \times R, \\ u(s, \xi) = u_0(s, \xi), & (s, \xi) \in [-r, 0] \times R, \end{cases} \tag{3.13}$$

where

$$Q(t - r, \xi - cr) = b(\phi + u) - b(\phi) - b'(\phi)u \tag{3.14}$$

with $\phi = \phi(\xi - cr)$ and $u = u(t - r, \xi - cr)$.

Let $w(\xi) > 0$ be the weight function defined in (2.12). Multiplying (3.13) by $e^{2\mu t} w(\xi)u(t, \xi)$, where $\mu > 0$ will be specified later in Lemma 3.4, we obtain

$$\begin{aligned} & \left\{ \frac{1}{2} e^{2\mu t} w u^2 \right\}_t + \left\{ \frac{1}{2} e^{2\mu t} c w u^2 - D_m e^{2\mu t} w u u_\xi \right\}_\xi + D_m e^{2\mu t} w u_\xi^2 + D_m e^{2\mu t} w' u u_\xi \\ & + \left\{ -\frac{c}{2} \frac{w'}{w} + d_m - \mu \right\} e^{2\mu t} w u^2 - \varepsilon e^{2\mu t} w u b'(\phi(\xi - cr)) u(t - r, \xi - cr) \\ & = \varepsilon e^{2\mu t} w u Q(t - r, \xi - cr). \end{aligned} \tag{3.15}$$

By the Cauchy-Schwarz inequality $|xy| \leq \frac{\delta}{2} x^2 + \frac{1}{2\delta} y^2$ for any $\delta > 0$ and then taking $\delta = 2$, we have

$$|D_m e^{2\mu t} w' u u_\xi| = D_m e^{2\mu t} w \left| u_\xi \cdot \frac{w'}{w} v \right| \leq D_m e^{2\mu t} w u_\xi^2 + \frac{D_m}{4} e^{2\mu t} \left(\frac{w'}{w} \right)^2 w u^2.$$

Applying the above inequality to (3.15), we obtain

$$\begin{aligned} & \left\{ \frac{1}{2} e^{2\mu t} w u^2 \right\}_t + \left\{ \frac{1}{2} e^{2\mu t} c w u^2 - D_m e^{2\mu t} w u u_\xi \right\}_\xi + \left\{ -\frac{c}{2} \frac{w'}{w} + d_m - \mu - \frac{D_m}{4} \left(\frac{w'}{w} \right)^2 \right\} e^{2\mu t} w u^2 \\ & - \varepsilon e^{2\mu t} w u b'(\phi(\xi - cr)) u(t - r, \xi - cr) \\ & \leq \varepsilon e^{2\mu t} w u Q(t - r, \xi - cr). \end{aligned} \tag{3.16}$$

Integrating (3.16) over $R \times [0, t]$ with respect to ξ and t yields

$$\begin{aligned} & e^{2\mu t} \|u(t)\|_{L_w^2}^2 + \int_0^t \int_R e^{2\mu\tau} \left\{ -c \frac{w'(\xi)}{w(\xi)} + 2d_m - 2\mu - \frac{D_m}{2} \left(\frac{w'(\xi)}{w(\xi)} \right)^2 \right\} w(\xi) u^2(\tau, \xi) d\xi d\tau \\ & - 2\varepsilon \int_0^t \int_R e^{2\mu\tau} w(\xi) b'(\phi(\xi - cr)) u(\tau, \xi) u(\tau - r, \xi - cr) d\xi d\tau \\ & \leq \|u_0(0)\|_{L_w^2}^2 + 2\varepsilon \int_0^t \int_R e^{2\mu\tau} w(\xi) u(\tau, \xi) Q(\tau - r, \xi - cr) d\xi d\tau. \end{aligned} \tag{3.17}$$

Using (2.1) and (2.4) for $b(v) = b_1(v)$, or, (2.2) and (2.5) for $b(v) = b_2(v)$, a straightforward computation gives

$$b'_1(\phi) = p(1 - aq\phi^q) e^{-a\phi^q} \geq 0, \quad \text{or} \quad b'_2(\phi) = \frac{p[1 - a(q - 1)\phi^q]}{(1 + a\phi^q)^2} \geq 0$$

for $v_- \leq \phi \leq v_+$ with the corresponding constant equilibria v_\pm given in (2.1) or (2.2), which gives

$$0 \leq b'(\phi) \leq p, \quad \text{for } v_- \leq \phi \leq v_+. \tag{3.18}$$

Furthermore, using the Cauchy-Schwarz inequality again, we obtain

$$\begin{aligned} & |2\varepsilon e^{2\mu\tau} w(\xi) b'(\phi(\xi - cr)) u(\tau, \xi) u(\tau - r, \xi - cr)| \\ & \leq \varepsilon e^{2\mu\tau} w(\xi) b'(\phi(\xi - cr)) \left[\eta u^2(\tau, \xi) + \frac{1}{\eta} u^2(\tau - r, \xi - cr) \right] \end{aligned}$$

for any positive constant η , which will be specified later in Lemma 3.3. Thus, by the change of variables (see [15,21] for details), the third term on the left-hand side of (3.17) can be bounded as follows:

$$\begin{aligned}
 & \left| 2\varepsilon \int_0^t \int_R e^{2\mu\tau} w(\xi) b'(\phi(\xi - cr)) u(\tau, \xi) u(\tau - r, \xi - cr) d\xi d\tau \right| \\
 & \leq \varepsilon \eta \int_0^t \int_R e^{2\mu\tau} w(\xi) b'(\phi(\xi - cr)) u^2(\tau, \xi) d\xi d\tau \\
 & \quad + \frac{\varepsilon}{\eta} \int_0^t \int_R e^{2\mu\tau} w(\xi) b'(\phi(\xi - cr)) u^2(\tau - r, \xi - cr) d\xi d\tau \\
 & = \varepsilon \eta \int_0^t \int_R e^{2\mu\tau} w(\xi) b'(\phi(\xi - cr)) u^2(\tau, \xi) d\xi d\tau \\
 & \quad + \frac{\varepsilon}{\eta} e^{2\mu r} \int_{-r}^{t-r} \int_R e^{2\mu\tau} w(\xi + cr) b'(\phi(\xi)) u^2(\tau, \xi) d\xi d\tau \\
 & \leq \varepsilon \eta \int_0^t \int_R e^{2\mu\tau} w(\xi) b'(\phi(\xi - cr)) u^2(\tau, \xi) d\xi d\tau \\
 & \quad + \frac{\varepsilon}{\eta} e^{2\mu r} \int_0^t \int_R e^{2\mu\tau} w(\xi + cr) b'(\phi(\xi)) u^2(\tau, \xi) d\xi d\tau \\
 & \quad + \frac{\varepsilon}{\eta} e^{2\mu r} \int_{-r}^0 \int_R e^{2\mu\tau} w(\xi + cr) b'(\phi(\xi)) u_0^2(\tau, \xi) d\xi d\tau. \tag{3.19}
 \end{aligned}$$

Substituting (3.19) into (3.17), we then have

$$\begin{aligned}
 & e^{2\mu t} \|u(t)\|_{L_w^2}^2 + \int_0^t \int_R e^{2\mu\tau} B_{\eta, \mu, w}(\xi) w(\xi) u^2(\tau, \xi) d\xi d\tau \\
 & \leq \|u_0(0)\|_{L_w^2}^2 + 2\varepsilon \int_0^t \int_R e^{2\mu\tau} w(\xi) u(\tau, \xi) Q(\tau - r, \xi - cr) d\xi d\tau \\
 & \quad + \frac{\varepsilon}{\eta} e^{2\mu r} \int_{-r}^0 \int_R e^{2\mu\tau} w(\xi + cr) b'(\phi(\xi)) u_0^2(s, \xi) d\xi d\tau, \tag{3.20}
 \end{aligned}$$

where

$$B_{\eta, \mu, w}(\xi) := A_{\eta, w}(\xi) - 2\mu - \frac{\varepsilon}{\eta} (e^{2\mu r} - 1) b'(\phi(\xi)) \frac{w(\xi + cr)}{w(\xi)} \tag{3.21}$$

and

$$A_{\eta,w}(\xi) := -c \frac{w'(\xi)}{w(\xi)} + 2d_m - \frac{D_m}{2} \left(\frac{w'(\xi)}{w(\xi)} \right)^2 - \varepsilon \eta b'(\phi(\xi - cr)) - \frac{\varepsilon}{\eta} \frac{w(\xi + cr)}{w(\xi)} b'(\phi(\xi)). \tag{3.22}$$

Using (2.1) and (2.4) for $b(v) = b_1(v)$, or (2.2) and (2.5) for $b(v) = b_2(v)$, another straightforward computation shows

$$b_1''(\phi) = -paq\phi^{q-1}(2 - aq\phi^q)e^{-a\phi^q} \leq 0, \quad \text{or} \quad b_2''(\phi) = -\frac{paq\phi^{q-1}[1 + q - a(q - 1)\phi^q]}{(1 + a\phi^q)^3} \leq 0$$

for $v_- \leq \phi \leq v_+$ with the corresponding constant equilibria v_{\pm} given in (2.1) or (2.2). This implies

$$b''(\phi) \leq 0, \quad \text{for } v_- \leq \phi \leq v_+. \tag{3.23}$$

Thus, by (3.23) and the Taylor’s formula, we have

$$Q(t - r, \xi - cr) = b(\phi + u) - b(\phi) - b'(\phi)u = \frac{b''(\tilde{\phi})}{2!}u^2 \leq 0, \tag{3.24}$$

where $\tilde{\phi}$ is some function between ϕ and $\phi + u = V^+(t, x)$. With the help of (3.24) and the fact that $w > 0, u > 0$, we immediately obtain

$$2\varepsilon \int_0^t \int_R e^{2\mu\tau} w(\xi)u(\tau, \xi)Q(\tau - r, \xi - cr) d\xi d\tau \leq 0. \tag{3.25}$$

Finally, substituting (3.25) into (3.20), we obtain

$$\begin{aligned} & e^{2\mu t} \|u(t)\|_{L_w^2}^2 + \int_0^t \int_R e^{2\mu\tau} B_{\eta,\mu,w}(\xi)w(\xi)u^2(\tau, \xi) d\xi d\tau \\ & \leq \|u_0(0)\|_{L_w^2}^2 + \frac{\varepsilon}{\eta} e^{2\mu r} \int_{-r}^0 \int_R e^{2\mu\tau} w(\xi + cr)b'(\phi(\xi))u_0^2(s, \xi) d\xi d\tau \\ & \leq C_1 \left(\|u_0(0)\|_{L_w^2}^2 + \int_{-r}^0 \|u_0(\tau)\|_{L_w^2}^2 d\tau \right) \end{aligned} \tag{3.26}$$

for some positive constant C_1 .

The most important step now is to prove $B_{\eta,\mu,w}(\xi) \geq C > 0$ for some constant C . For this we need the following key lemma.

Lemma 3.3. *Let $\eta = e^{-\lambda cr}$. Then*

$$A_{\eta,w}(\xi) \geq C_2 > 0, \quad \xi \in R, \tag{3.27}$$

for some positive constant C_2 .

Proof. Case 1: $\xi \leq x_0$. From (2.12), we have $w(\xi) = e^{-2\lambda(\xi-x_0-cr)}$ and $w(\xi + cr) = e^{-2\lambda(\xi-x_0)}$. Notice that, $b'(\phi)$ is decreasing for $\phi \in (v_-, v_+)$ (since $b''(\phi) < 0$ as shown in (3.23)) and $\phi(\xi)$ is increasing for $\xi \in (-\infty, \infty)$. Thus, $b'(\phi(\xi))$ is decreasing in ξ . This implies

$$0 < b'(\phi(x_0)) < b'(\phi(\xi)) < b'(\phi(\xi - cr)) < b'(\phi(-\infty)) = b'(v_-) = p. \tag{3.28}$$

Notice also $\eta = e^{-\lambda cr}$. We then obtain

$$\begin{aligned} A_{\eta,w}(\xi) &= 2\lambda c + 2d_m - \frac{D_m}{2}(-2\lambda)^2 - \varepsilon\eta b'(\phi(\xi - cr)) - \frac{\varepsilon}{\eta}b'(\phi(\xi))e^{-2\lambda cr} \\ &\geq 2\lambda c + 2d_m - 2D_m\lambda^2 - \varepsilon\eta p - \frac{\varepsilon}{\eta}pe^{-2\lambda cr} \\ &= 2(c\lambda + d_m - D_m\lambda^2 - \varepsilon pe^{-\lambda cr}) \\ &=: C_3 > 0, \end{aligned} \tag{3.29}$$

where we used (2.11) for the last step.

Case 2: $x_0 < \xi \leq x_0 + cr$. In this case, we have $w(\xi) = e^{-2\lambda(\xi-x_0-cr)}$ and $w(\xi + cr) = 1$. Since $b'(\phi(\xi))$ is decreasing in ξ , i.e., $b'(\phi(x_0 + cr)) < b'(\phi(\xi)) \leq b'(\phi(x_0)) < b'(\phi(\xi - cr)) \leq b'(\phi(x_0 - cr))$, and using the fact $e^{2\lambda(\xi-x_0-cr)} \leq e^0 = 1$ due to $\xi - x_0 - cr \leq 0$, we then obtain

$$\begin{aligned} A_{\eta,w}(\xi) &= 2\lambda c + 2d_m - 2D_m\lambda^2 - \varepsilon\eta b'(\phi(\xi - cr)) - \frac{\varepsilon}{\eta}b'(\phi(\xi))e^{2\lambda(\xi-x_0-cr)} \\ &\geq 2\lambda c + 2d_m - 2D_m\lambda^2 - \varepsilon\eta b'(\phi(x_0 - cr)) - \frac{\varepsilon}{\eta}b'(\phi(x_0 - cr)) \\ &= 2\lambda c + 2d_m - 2D_m\lambda^2 - 2\varepsilon b'(\phi(x_0 - cr))\left(\frac{\eta}{2} + \frac{1}{2\eta}\right) \\ &= 2\lambda c + 2d_m - 2D_m\lambda^2 - 2\varepsilon b'(\phi(x_0 - cr)) \cosh(\lambda cr) \\ &=: C_4. \end{aligned} \tag{3.30}$$

Now we need to prove $C_4 > 0$. In fact, since $b'(v_+) \ll 1$ (we assume it as a sufficient condition in Theorem 2.2, see also Remark 1 for the explanation), we may take

$$b'(v_+) < \frac{\min\{\lambda c + d_m - D_m\lambda^2, d_m\}}{\varepsilon \cosh(\lambda cr)}.$$

Let x_0 be sufficiently large such that

$$|b'(v_+) - b'(\phi(x_0 - cr))| \ll 1.$$

In that case, we have

$$b'(\phi(x_0 - cr)) < \frac{\min\{\lambda c + d_m - D_m\lambda^2, d_m\}}{\varepsilon \cosh(\lambda cr)}. \tag{3.31}$$

With this help, we further obtain from (3.30) that

$$A_{\eta,w}(\xi) \geq 2[\lambda c + d_m - D_m\lambda^2 - \varepsilon b'(\phi(x_0 - cr)) \cosh(\lambda cr)] = C_4 > 0. \tag{3.32}$$

Case 3: $\xi > x_0 + cr$. In this case, $w(\xi) = w(\xi + cr) = 1$. Since $b'(\phi(\xi))$ is decreasing in ξ , we have

$$\begin{aligned} A_{\eta,w}(\xi) &= 2d_m - \varepsilon\eta b'(\phi(\xi - cr)) - \frac{\varepsilon}{\eta} b'(\phi(\xi)) \\ &\geq 2d_m - \varepsilon\eta b'(\phi(x_0 - cr)) - \frac{\varepsilon}{\eta} b'(\phi(x_0 - cr)) \\ &= 2d_m - 2\varepsilon b'(\phi(x_0 - cr)) \cosh(\lambda cr) \\ &=: C_5 > 0, \end{aligned} \tag{3.33}$$

because of (3.31).

Finally, let

$$C_2 := \min\{C_3, C_4, C_5\}. \tag{3.34}$$

Then (3.29), (3.32) and (3.33) imply (3.27). \square

Lemma 3.4. *It holds that*

$$B_{\mu,\eta,w}(\xi) > 0, \quad \xi \in (-\infty, \infty), \tag{3.35}$$

for $0 < \mu < \mu_1$, where $\mu_1 > 0$ is the unique root of the following equation

$$C_2 - 2\mu_1 - \frac{\varepsilon p}{\eta} (e^{2\mu_1 r} - 1) = 0. \tag{3.36}$$

Proof. As shown in Lemma 3.3, we have $0 < b'(\phi(\xi)) < p$ for $\xi \in (-\infty, \infty)$, and

$$\frac{w(\xi + cr)}{w(\xi)} = \begin{cases} e^{-2\lambda cr} < 1, & \text{for } \xi \leq x_0, \\ e^{2\lambda(\xi - x_0 - cr)} \leq e^0 = 1, & \text{for } x_0 < \xi \leq x_0 + cr, \\ 1, & \text{for } \xi > x_0 + cr. \end{cases}$$

It follows immediately that

$$\begin{aligned} B_{\mu,\eta,w}(\xi) &= A_{\eta,w}(\xi) - 2\mu - \frac{\varepsilon}{\eta} (e^{2\mu r} - 1) b'(\phi(\xi)) \frac{w(\xi + cr)}{w(\xi)} \\ &\geq C_2 - 2\mu - \frac{\varepsilon p}{\eta} (e^{2\mu r} - 1) > 0 \end{aligned} \tag{3.37}$$

for $0 < \mu < \mu_1$. \square

By dropping the positive term $\int_0^t \int_R e^{2\mu\tau} B_{\eta,\mu,w}(\xi) w(\xi) u^2(\tau, \xi) d\xi d\tau$ in (3.26), we obtain the first basic energy estimate.

Lemma 3.5. *It holds that*

$$e^{2\mu t} \|u(t)\|_{L^2_w}^2 \leq C_1 \left(\|u_0(0)\|_{L^2_w}^2 + \int_{-r}^0 \|u_0(\tau)\|_{L^2_w}^2 d\tau \right), \quad t \geq 0. \tag{3.38}$$

Next, differentiating (3.13) with respect to ξ , and multiplying it by $e^{2\mu t} w(\xi) u_\xi(t, \xi)$, then integrating the resultant equation with respect to (t, ξ) over $[0, t] \times R$, and using the basic energy estimates (3.38) in Lemma 3.5, we can similarly prove the second energy estimate.

Lemma 3.6. *It holds that*

$$e^{2\mu t} \|u_\xi(t)\|_{L^2_w}^2 \leq C_6 \left(\|u_0(0)\|_{H^1_w}^2 + \int_{-r}^0 \|u_0(\tau)\|_{H^1_w}^2 d\tau \right) \tag{3.39}$$

for some positive constant C_6 .

Combining Lemmas 3.5 and 3.6, and noting $w(\xi) \geq 1$ for all $\xi \in R$, from the definition of w , we obtain the following decay rate.

Lemma 3.7. *It holds that*

$$\|u(t)\|_{H^1} \leq \|u(t)\|_{H^1_w} \leq C_7 e^{-\mu t} \left(\|u_0(0)\|_{H^1_w}^2 + \int_{-r}^0 \|u_0(\tau)\|_{H^1_w}^2 d\tau \right)^{\frac{1}{2}}, \quad t \geq 0, \tag{3.40}$$

where $C_7 = \max\{\sqrt{C_1}, \sqrt{C_6}\}$.

Using Sobolev embedding theorem $H^1(R) \hookrightarrow C^0(R)$, we finally have the following stability result.

Lemma 3.8. *It holds that*

$$\sup_{x \in R} |V^+(t, x) - \phi(x + ct)| = \sup_{\xi \in R} |u(t, \xi)| \leq C_8 e^{-\mu t}, \quad t \geq 0, \tag{3.41}$$

for some positive constant C_8 .

Step 2. *The convergence of $V^-(t, x)$ to $\phi(x + ct)$.*

Let $\xi = x + ct$ and

$$u(t, \xi) = \phi(x + ct) - V^-(t, x), \quad u_0(s, \xi) = \phi(x + cs) - V_0^-(s, x). \tag{3.42}$$

As in Step 1, we can similarly prove that $V^-(t, x)$ converges to $\phi(x + ct)$.

Lemma 3.9. *It holds that*

$$\sup_{x \in R} |V^-(t, x) - \phi(x + ct)| = \sup_{\xi \in R} |u(t, \xi)| \leq C_9 e^{-\mu t}, \quad t \geq 0, \tag{3.43}$$

for some positive constant C_9 .

Step 3. *The convergence of $v(t, x)$ to $\phi(x + ct)$.*

Finally, we prove that $v(t, x)$ converges to $\phi(x + ct)$ as follows.

Lemma 3.10. *It holds that*

$$\sup_{x \in R} |v(t, x) - \phi(x + ct)| \leq C_{10} e^{-\mu t}, \quad t \geq 0, \quad (3.44)$$

for some positive constant C_{10} .

Proof. Since the initial data satisfy $V_0^-(x, s) \leq v_0(x, s) \leq V_0^+(x, s)$, from Lemma 3.2, it can be verified that the corresponding solutions of (1.7) and (1.2) satisfy

$$V^-(t, x) \leq v(t, x) \leq V^+(t, x), \quad (t, x) \in R_+ \times R.$$

Thanks to Lemmas 3.8 and 3.9, we have the following convergence results,

$$\sup_{x \in R} |V^-(t, x) - \phi(x + ct)| \leq C_9 e^{-\mu t}, \quad \sup_{x \in R} |V^+(t, x) - \phi(x + ct)| \leq C_8 e^{-\mu t}.$$

Combining these inequalities, the squeeze technique gives

$$\sup_{x \in R} |v(t, x) - \phi(x + ct)| \leq C_{10} e^{-\mu t}, \quad t > 0,$$

for some positive constant C_{10} . This completes the proof. \square

Acknowledgments

The authors would like to thank the anonymous referee for his helpful suggestions and comments, which lead us to have this improved stability result. The research of M.M. was supported in part by Natural Sciences and Engineering Research Council of Canada under the NSERC grant RGPIN 354724-08, the research of C.-K.L. was supported in part by National Science Council of Taiwan, ROC, under the grant 95-2115-M-009-019-MY3, and the research of C.-T.L. was supported in part by National Science Council of Taiwan, ROC, under the grants 96-2115-M-126-003 and 97-2115-M-126-002.

References

- [1] J. Al-Omari, S.A. Gourley, Monotone travelling fronts in an age-structured reaction–diffusion model of a single species, *J. Math. Biol.* 45 (2002) 294–312.
- [2] J. Al-Omari, S.A. Gourley, Monotone wave-fronts in a structured population model with distributed maturation delay, *IMA J. Appl. Math.* 70 (2005) 858–879.
- [3] J. Fang, J. Wei, X.-Q. Zhao, Spatial dynamics of a nonlocal and time-delayed reaction–diffusion system, *J. Differential Equations* 245 (2008) 2749–2770.
- [4] R.A. Fisher, The wave of advance of advantageous genes, *Ann. Eugen.* 7 (1937) 353–369.
- [5] S.A. Gourley, Linear stability of travelling fronts in an age-structured reaction–diffusion population model, *Quart. J. Mech. Appl. Math.* 58 (2005) 257–268.
- [6] S.A. Gourley, Y. Kuang, Wavefronts and global stability in a time-delayed population model with stage structure, *Proc. R. Soc. Lond. Ser. A* 459 (2003) 1563–1579.
- [7] S.A. Gourley, J.W.-H. So, J. Wu, Non-locality of reaction–diffusion equations induced by delay: Biological modeling and non-linear dynamics, in: D.V. Anosov, A. Skubachevskii (Eds.), *Contemporary Mathematics, Thematic Surveys*, Kluwer/Plenum, 2003, pp. 84–120 (in Russian); also appeared in: *J. Math. Sci.* 124 (2004) 5119–5153 (in English).
- [8] S.A. Gourley, J. Wu, Delayed nonlocal diffusive system in biological invasion and disease spread, in: *Fields Inst. Commun.*, vol. 48, 2006, pp. 137–200.
- [9] W.S.C. Gurney, S.P. Blythe, R.M. Nisbet, Nicholson's blowflies revisited, *Nature* 287 (1980) 17–21.
- [10] A. Kolmogorov, I. Petrowsky, N. Piskounov, Study of the diffusion equation with growth of the quantity of matter and its application to biological problems, *Bull. Univ. Etat Moscou* 6 (1937) 1–25.
- [11] G. Li, M. Mei, Y.S. Wong, Nonlinear stability of travelling wavefronts in an age-structured reaction–diffusion population model, *Math. Biosci. Eng.* 5 (2008) 85–100.
- [12] W.-T. Li, S. Ruan, Z.-C. Wang, On the diffusive Nicholson's blowflies equation with nonlocal delays, *J. Nonlinear Sci.* 17 (2007) 505–525.

- [13] W.-T. Li, Z.-C. Wang, J. Wu, Entire solutions in monostable reaction–diffusion equations with delayed nonlinearity, *J. Differential Equations* 245 (2008) 102–129.
- [14] D. Liang, J. Wu, Traveling waves and numerical approximations in a reaction–diffusion equation with nonlocal delayed effect, *J. Nonlinear Sci.* 13 (2003) 289–310.
- [15] C.-K. Lin, M. Mei, On travelling wavefronts of the Nicholson's blowflies equations with diffusion, submitted for publication.
- [16] S. Ma, X. Zou, Existence, uniqueness and stability of travelling waves in a discrete reaction–diffusion monostable equation with delay, *J. Differential Equations* 217 (2005) 54–87.
- [17] R.H. Martin, H.L. Smith, Abstract functional-differential equations and reaction–diffusion systems, *Trans. Amer. Math. Soc.* 321 (1990) 1–44.
- [18] M. Mei, C.-K. Lin, C.-T. Lin, J.W.-H. So, Traveling wavefronts for time-delayed reaction–diffusion equation: (II) Nonlocal nonlinearity, *J. Differential Equations* 247 (2) (2009) 511–529 (this issue).
- [19] M. Mei, K. Nishihara, Nonlinear stability of travelling waves for one-dimensional viscoelastic materials with non-convex nonlinearity, *Tokyo J. Math.* 20 (1997) 241–264.
- [20] M. Mei, J.W.-H. So, Stability of strong traveling waves for a nonlocal time-delayed reaction–diffusion equation, *Proc. Roy. Soc. Edinburgh Sect. A* 138 (2008) 551–568.
- [21] M. Mei, J.W.-H. So, M.Y. Li, S.S.P. Shen, Asymptotic stability of traveling waves for the Nicholson's blowflies equation with diffusion, *Proc. Roy. Soc. Edinburgh Sect. A* 134 (2004) 579–594.
- [22] J.A.J. Metz, O. Diekmann, *The Dynamics of Physiologically Structured Populations*, edited by J.A.J. Metz and O. Diekmann, Springer-Verlag, New York, 1986.
- [23] C. Ou, J. Wu, Persistence of wavefronts in delayed non-local reaction–diffusion equations, *J. Differential Equations* 235 (2007) 219–261.
- [24] K.W. Schaaf, Asymptotic behavior and traveling wave solutions for parabolic functional differential equations, *Trans. Amer. Math. Soc.* 302 (1987) 587–615.
- [25] H.L. Smith, X.-Q. Zhao, Global asymptotic stability of traveling waves in delayed reaction–diffusion equations, *SIAM J. Math. Anal.* 31 (2000) 514–534.
- [26] J.W.-H. So, J. Wu, X. Zou, A reaction–diffusion model for a single species with age structure: (I) Traveling wavefronts on unbounded domains, *Proc. R. Soc. Lond. Ser. A* 457 (2001) 1841–1853.
- [27] J.W.-H. So, J. Wu, Y. Yang, Numerical Hopf bifurcation analysis on the diffusive Nicholson's blowflies equation, *Appl. Math. Comput.* 111 (2000) 53–69.
- [28] J.W.-H. So, Y. Yang, Dirichlet problem for the diffusive Nicholson's blowflies equation, *J. Differential Equations* 150 (1998) 317–348.
- [29] J.W.-H. So, X. Zou, Traveling waves for the diffusive Nicholson's blowflies equation, *Appl. Math. Comput.* 22 (2001) 385–392.
- [30] H.R. Thieme, *Mathematics in Population Biology*, Princeton Univ. Press, Princeton, 2003.
- [31] H. Thieme, X.-Q. Zhao, Asymptotic speeds of spread and traveling waves for integral equation and delayed reaction–diffusion models, *J. Differential Equations* 195 (2003) 430–470.
- [32] A. Volpert, Vi. Volpert, Vi. Volpert, *Traveling Wave Solutions of Parabolic Systems*, *Transl. Math. Monogr.*, vol. 140, Amer. Math. Soc., Providence, RI, 1994.
- [33] D. Wei, J.-Y. Wu, M. Mei, Remark on critical speed of traveling wavefronts for Nicholson's blowflies equation with diffusion, *Acta Math. Sci.*, in press.
- [34] J.-Y. Wu, D. Wei, M. Mei, Analysis on critical speed of traveling waves, *Appl. Math. Lett.* 20 (2007) 712–718.
- [35] J.-H. Wu, *Theory and Applications of Partial Functional-Diffusion Equations*, *Appl. Math. Sci.*, vol. 119, Springer-Verlag, New York, 1996.
- [36] X.-Q. Zhao, *Dynamical Systems in Population Biology*, *CMS Books Math.*, vol. 16, Springer-Verlag, New York, 2003.
- [37] Y. Yang, J.W.-H. So, Dynamics for the diffusive Nicholson's blowflies equation, in: W. Chen, S. Hu (Eds.), *Dynamical Systems and Differential Equations*, vol. II, Southwest Missouri State University Press, Springfield, MO, 1998, pp. 333–352.