

Reference

Books and Thesis for futher reading

- [B1] Ralf Menzel, "Photonics: Linear and Nonlinear Interactions of Laser Light and Matter", ISBN: 3-540-67074-2 (Springer-Verlag, 2001).
- [B2] Walter Joechner, "Solid-State Laser Engineering", 4th ed., ISBN: 3-540-60237-2 (Springer-Verlag, 1996)
- [B3] <http://www.rp-photonics.com> (2006) (virtual library)
- [B4] Walter Joechner, Michiale Bass, "Solid-State Lasers: A Graduate Text", ISBN: 0-387-95590-9 (Springer-Verlag, 2003)
- [B5] Joseph T. Verdeyen, "Laser Electronics", 3rd ed., ISBN: 0-13-706666-X (Prentice Hall, 2000)
- [B6] V. G. Dmitriev, G. G. Gurzadyan, D. N. Nikogosyan, "Handbook of Nonlinear Optical Crystals", 3rd ed., ISBN: 3-540-65394-5 (Springer-Verlag, 1999)
- [T1] Hui Deng, "Dynamic Condensation of Semiconductor Microcavity Polaritons", Stanford University, Ph.D. thesis (2006)
- [T2] Riccardo Sapienza, "Photonic Nano Materials: Anisotropic Transport and Optical Bloch Oscillations", European Ph.D. school, XVII ciclo, Ph.D. thesis (2005)

Chapter 1 - Introduction

- [1.1] T. H. Maiman, "Stimulated optical radiation in ruby", Nature 187, 493 (1960)
- [1.2] 石中達, "全球雷射市場消長分析及 2000 年預測", PIDA (1998)
- [1.3] D. L. Spies, "Highly efficient neodymium:yttrium aluminum garnet laser end

- pumped by a semiconductor laser array", Appl. Phys. Lett. 47, 74-76 (1985)
- [1.4] D. C. Hanna and W. A. Clarkson, "A review of diode-pumped lasers", in "Advances in Lasers and Applications", D. M. Finlayson and B. Sinclair (editors), proceedings of the fifty-second Scottish Universities Summer School in Physics, St. Andrews, (September 1998) (eBook)
- ## Chapter 2 - QD SESAM
- [2.1] A. S. Grabtchikov, A. N. Kuzmin, V. A. Lisinetskii, V. A. Orlovich, A. A. Demidovich, K. V. Yumashev, N. V. Kuleshov, H. J. Eichler, and M. V. Danailov, "Passively Q -switched 1.35- μm diode-pumped Nd:KGW laser with V:YAG saturable absorbers", Opt. Mater. 16, 349-352 (2001)
- [2.2] K. V. Yumashev, I. A. Denisov, N. N. Posnov, P. V. Prokoshin, and V. P. Mikhailov, "Nonlinear absorption properties of $\text{Co}^{2+}:\text{MgAl}_2\text{O}_4$ crystal", Appl. Phys. B 70, 179-184 (2000)
- [2.3] V. G. Savitski, N. N. Posnov, P. V. Prokoshin, A. M. Malyarevich, K. V. Yumashev, M. I. Demchuk, and A. A. Lipovski, "PbS-doped phosphate glasses saturable absorbers for 1.3- μm neodymium lasers", Appl. Phys. B 75, 841-846 (2002)
- [2.4] R. Fluck, B. Braun, E. Gini, H. Melchior, and U. Keller, "Passively Q -switched 1.34- μm Nd:YVO₄ microchip laser with semiconductor saturable absorbers mirrors", Opt. Lett. 22, 991-993 (1997)
- [2.5] R. Fluck, G. Zhang, U. Keller, K. J. Weingarten, and M. Moser, "Diode-pumped passively mode-locked 1.3- μm Nd:YVO₄ and Nd:YLF lasers by use of semiconductor saturable absorbers," Opt. Lett. 21, 1378-1380 (1996)
- [2.6] M. Kondow, K. Uomi, A. Niwa, T. Kitatani, S. Watahiki, and Y. Yazawa, "GaInNAs: A novel material for long-wavelength-range laser diodes with excellent high-temperature performance," Japan. J. Appl. Phys. 35, 1273-1275 (1996)
- [2.7] D. L. Huffaker, G. Park, Z. Zou, O. B. Shchekin, and D. G. Deppe, "1.3 μm room-temperature GaAs-based quantum-dot laser," Appl. Phys. Lett. 73, 2564-2566 (1998)
- [2.8] H. D. Sun, G. J. Valentine, R. Macaluso, S. Calvez, D. Burns, M. D. Dawson, T. Jouhti, and M. Pessa, "Low-loss 1.3- μm GaInNAs saturable Bragg reflector for

- high-power picosecond neodymium lasers," Opt. Lett. **27**, 2124-2126 (2002).
- [2.9] V. Liverini, S. Schön, R. Grange, M. Haiml, S. C. Zeller, and U. Keller, "Low-loss GaInNAs saturable absorber mode locking a 1.3- μm solid-state laser," Appl. Phys. Lett. **84**, 4002-4004 (2004)
- [2.10] A. Markus, A. Fiore, J. D. Ganière, U. Oesterle, J. X. Chen, B. Deveaud, M. Illegems, H. Riechert, "Comparison of radiative properties of InAs quantum dots and GaInNAs quantum wells emitting around 1.3 μm ," Appl. Phys. Lett. **80**, 911-913 (2002)
- [2.11] A. Passaseo, V. Tasco, M. De Giorgi, M. T. Todaro, M. De Vittorio, and R. Cingolani, "Long wavelength emission in $\text{In}_x\text{Ga}_{1-x}\text{As}$ quantum dot structure grown in a GaAs barrier by metalorganic chemical vapor deposition," Appl. Phys. Lett. **84**, 1868-1870 (2004)
- [2.12] Y. V. Volk, I. A. Denisov, A. M. Malyarevich, K. V. Yumashev, O. S. Dymshits, A. V. Shashkin, A. A. Zhilin, U. Kang, and K. H. Lee, "Magnesium- and zinc-aluminosilicate cobalt-doped glass ceramics as saturable absorbers for diode-pumped 1.3 μm laser," Appl. Opt. **43**, 682-687 (2004)
- [2.13] O. Okhotnikov, A. Grudinin, M. Pessa, "Ultra-fast fibre laser systems based on SESAM technology: new horizons and applications", New J. Phys. **6**, 177 (2004)
- [2.14] U. Keller, D. A. B. Miller, G. D. Boyd, T. H. Chiu, J. F. Ferguson, and M. T. Asom, "Solid-state low-loss intracavity saturable absorber for Nd:YLF lasers: an A-FPSA," Opt. Lett. **17**, 505-508 (1992)
- [2.15] G. J. Spühler, T. Südmeyer, R. Paschotta, M. Moser, K. J. Weingarten, and U. Keller, "Passively mode-locked high-power Nd:YAG lasers with multiple laser heads," Appl. Phys. B **71**, 19-25 (2000)
- [2.16] J. L. He, Y. X. Fan, J. Du, Y. G. Wang, S. Liu, H. T. Wang, L. H. Zhang, and Y. Hang, "4-ps passively mode locked Nd:Gd_{0.5}Y_{0.5}VO₄ laser with a semiconductor saturable absorber mirror," Opt. Lett. **29**, 2803-2805 (2004)
- [2.17] A. A. Kaminskii, K. Ueda, H. J. Eichler, Y. Kuwano, H. Kouta, S. N. Bagaev, T. H. Chyba, J. C. Barnes, G. M. A. Gad, T. Murai, and J. Lu, "Tetragonal vanadates YVO₄ and GdVO₄ - new efficient $\chi^{(3)}$ -materials for Raman lasers," Opt. Commu. **194**, 201-206 (2001)

- [2.18] Y. F. Chen, "Compact efficient all-solid-state eye-safe laser with self-frequency Raman conversion in a Nd:YVO₄ crystal," Opt. Lett. 29, 2172-2174 (2004)
- [2.19] A. R. Kovsh, N. A. Maleev, A. E. Zhukov, S. S. Mikhrin, A. P. Vasil'ev, Yu. M. Shernyakov, M. V. Maximov, D. A. Livshits, V. M. Ustinov, Zh. I. Alferov, N. N. Ledentsov, and D. Bimberg, "InAs/InGaAs/GaAs quantum dot lasers of 1.3 μm range with high (88%) differential efficiency," Electron. Lett., 38, pp. 1104-1106 (2002)
- [2.20] E. U. Rafailov, S. J. White, A. A. Lagatsky, A. Miller, W. Sibbett, D. A. Livshits, A. E. Zhukov, and V. M. Ustinov, "Fast Quantum-dot saturable absorber for passive mode-locking of solid-state lasers," IEEE Photon. Technol. Lett. 16, pp. 2439-2441 (2004)
- [2.21] X. Liu, L. Qian, F. Wise, Z. Zhang, T. Itatani, T. Sugaya, T. Nakagawa, and W. Torizuka, "Femtosecond Cr:forsterite laser diode pumped by a double-clad fiber," Opt. Lett. 15, 129-131 (1998)
- [2.22] http://www.rp-photonics.com/semiconductor_saturable_absorber_mirrors.html
- [2.23] U. Keller et al., "Semiconductor saturable absorber mirrors (SESAMs) for femtosecond to nanosecond pulse generation in solid-state lasers," IEEE J. Sel. Top. Quantum Electron. 2, 435 (1996)
- [2.24] Y.F. Chen, Y.P. Lan, H.L. Chang, "Analytical model for design criteria of passively Q-switched lasers," IEEE J Quantum Electron. 37, 3, 462-468 (2001). Y.F. Chen, S.W. Tsai, "Simultaneous Q-switching and mode-locking in a diode-pumped Nd:YVO₄/Cr⁴⁺:YAG laser," IEEE J Quantum Electron. 37, 4, 580-586 (2001). Y.F. Chen, J.L. Lee, H.D. Hsieh, S.W. Tsai, "Analysis of passively Q-switched lasers with simultaneous mode-locking," IEEE J Quantum Electron. 38, 3, 312-317 (2002)
- [2.25] Ling-Yi Tsai, Ch. 3 in "Passively Q-switched intracavity optical parametric oscillator," National Chiao Tung University, M.S. thesis (2005)
- [2.26] M. Sheik, A.A. Said, and E.W. Van Stryland, "High-sensitivity, single-beam n₂ measurements," Opt. Lett. 14, 955 (1989). M. Sheik, A.A. Said, T.H. Wei, D.J. Hagan, and E.W. Van Stryland, "Sensitive measurement of optical nonlinearities using a single beam," IEEE J. Quantum Electron. QE-26, 760 (1990). J. Wang, M. Sheik-Bahae, A.A. Said, D.J. Hagan, E.W. Van Stryland, "Time-resolved Z-scan measurements of optical nonlinearities," J. Opt. Soc. Am. B 11, 6, 1009

(1994)

- [2.27] W. Zhao and P. Palffy-Muhoray, "Z-scan technique using top-hat beams," *Appl. Phys. Lett.* 63, 12, 1613-1615 (1993)
- [2.28] <http://britneyspears.ac/physics/dos/dos.htm> (Britney's Guide to Semiconductor Physics)
- [2.29] A. McWilliam, A. A. Lagatsky, C. T. A. Brown, W. Sibbett, A. E. Zhukov, V. M. Ustinov, A. P. Vasil'ev, and E. U. Rafailov, "Quantum-dot-based saturable absorber for femtosecond mode-locked operation of a solid-state laser," *Opt. Lett.* 31, 1444-1446 (2006)

Chapter 3 - QW SESAM

- [3.1] A. M. Malyarevich, I. A. Denisov, K. V. Yumashev, V. P. Mikhailov, R. S. Conroy, and B. D. Sinclair, *Appl. Phys. B* 67, 555-558 (1998)
- [3.2] A. S. Grabtchikov, A. N. Kuzmin, V. A. Lisinetskii, V. A. Orlovich, A. A. Demidovich, K. V. Yumashev, N. V. Kuleshov, H. J. Eichler, and M. V. Danailov, *Opt. Mater.* 16, 349-352 (2001)
- [3.3] A. Agnesi, A. Guandalini, G. Reali, J. K. Jabczynski, K. Kopczynski, and Z. Mierczyk, *Opt. Commun.* 194, 429-433 (2001)
- [3.4] K. V. Yumashev, I. A. Denisov, N. N. Posnov, P. V. Prokoshin, and V. P. Mikhailov, *Appl. Phys. B* 70, 179-184 (2000)
- [3.5] Y. V. Volk, I. A. Denisov, A. M. Malyarevich, K. V. Yumashev, O. S. Dymshits, A. V. Shashkin, A. A. Zhilin, U. Kang, and K. H. Lee, *Appl. Opt.* 43, 682-687 (2004)
- [3.6] V. G. Savitski, N. N. Posnov, P. V. Prokoshin, A. M. Malyarevich, K. V. Yumashev, M. I. Demchuk, and A. A. Lipovski, *Appl. Phys. B* 75, 841-846 (2002)
- [3.7] R. Fluck, G. Zhang, U. Keller, K. J. Weingarten, and M. Moser, "Diode-pumped passively mode-locked 1.3- μm Nd:YVO₄ and Nd:YLF lasers by use of semiconductor saturable absorbers," *Opt. Lett.* 21, 1378-1380 (1996)
- [3.8] H. D. Sun, G. J. Valentine, R. Macaluso, S. Calvez, D. Burns, M. D. Dawson, T. Jouhti, and M. Pessa, *Opt. Lett.* 27, 2124-2126 (2002).

- [3.9] V. Liverini, S. Schön, R. Grange, M. Haiml, S. C. Zeller, and U. Keller, *Appl. Phys. Lett.* 84, 4002-4004 (2004)
- [3.10] H. C. Lai, A. Li, K. W. Su, M. L. Ku, Y. F. Chen, and K. F. Huang, *Opt. Lett.* 30, 480-483 (2005).
- [3.11] R. Fluck, B. Braun, E. Gini, H. Melchior, and U. Keller, *Opt. Lett.* 22, 991-993 (1997)
- [3.12] O. Ostinelli, W. Bächtold, M. Haiml, R. Grange, U. Keller, M. Ebnöther, E. Gini, G. Almuneau, *J. Cryst. Growth* 286, 247 (2006)
- [3.13] Y. Tsou, E. Garmire, W. Chen, M. Birnbaum, and R. Asthana, *Opt. Lett.* 18, 1514-1516 (1993)
- [3.14] W. Koechner, *Solid-State Laser Engineering*, vol. 1, Optical Sciences, 5th ed., (Springer, Berlin, 1999)
- [3.15] Y. Kaneda, M. Oka, H. Masuda, and S. Kubota, “7.6 W of continuous-wave radiation in a TEM₀₀ mode from a laser-diode end-pumped Nd:YAG laser,” *Opt. Lett.* 17, 1003-1005 (1992)
- [3.16] T. Kellner, F. Heine, and G. Huber, “Efficient laser performance of Nd:YAG at 946 nm and intracavity frequency doubling with LiJO₃, β -BaB₂O₄, and LiB₃O₅,” *Appl. Phys. B* 65, 789–902 (1997)
- [3.17] N. Moore, W. A. Clarkson, D.C. Hanna, S. Lehmann, and J. Bösenberg, “Efficient operation of a diode-bar-pumped Nd:YAG laser on the low-gain 1123-nm line,” *Appl. Opt.* 38, 5761–5764 (1999)
- [3.18] X. Guo, M. Chen, G. Li, B. Zhang, J. Yang, Z. Zhang, and Y. Wang, “Diode-pumped 1123-nm Nd:YAG laser,” *Chin. Opt. Lett.* 2, 402–404 (2004)
- [3.19] Z. Cai, M. Chen, Z. Zhang, R. Zhou, W. Wen, X. Ding, and J. Yao, “Diode end-pumped 1123-nm Nd:YAG laser with 2.6-W output power,” *Chin. Opt. Lett.* 3, 281–282 (2005)
- [3.20] Y.F. Chen and Y.P. Lan, “Diode-pumped passively Q-switched Nd:YAG laser at 1123 nm,” *Appl. Phys. B* 79, 29–31 (2004)
- [3.21] Y. F. Chen, Y. P. Lan, and S. W. Tsai, “High-power diodepumped actively Q-switched Nd:YAG laser at 1123 nm,” *Opt. Commun.* 234, 309–313 (2004)
- [3.22] R. Paschotta, N. Moore, W.A. Clarkson, A.C. Tropper, D.C. Hannnd, and G. Mazé,

- "230 mW of blue light from a thulium doped upconversion fiber," IEEE J. Sel. Top. Quantum Electron. 3, 1100–1102 (1997)
- [3.23] G.J. Spühler, R. Paschotta, R. Fluck, B. Braun, M. Moser, G. Zhang, E. Gini, and U. Keller, "Experimentally confirmed design guidelines for passively Q-switched microchip lasers using semiconductor saturable absorbers," J. Opt. Soc. Am. B 16, 376–388 (1999)
- [3.24] R. Fluck, G. Zhang, U. Keller, K. J. Weingarten, and M. Moser, "Diode-pumped passively mode-locked 1.3- μm Nd:YVO₄ and Nd:YLF lasers," Opt. Lett. 21, 1378–1380 (1996)
- [3.25] F. Bugge, G. Erbert, J. Fricke, S. Gramlich, R. Staske, H. Wenzel, U. Zeimer, and M. Weyers, "12 W continuous-wave diode lasers at 1120 nm with InGaAs quantum wells," Appl. Phys. Lett. 79, 1965–1967 (2001)
- [3.26] S. Mogg, N. Chitica, R. Schatz, and M. Hammar, "Properties of highly strained InGaAs GaAs quantum wells for 1.2- μm laser diodes," Appl. Phys. Lett. 81, 2334–2336 (2002)
- [3.27] T. Kondo, D. Schlenker, T. Miyamoto, Z. Chen, M. Kawaguchi, E. Gouardes, F. Koyama, and K. Iga, "Lasing characteristics of 1.2 μm highly strained GaInAs GaAs quantum well lasers," Jpn. J. Appl. Phys. 40, 467–471 (2001)

Chapter 4 - Optical-pumped VECSEL

- [4.1] A. S. Grabtchikov, A. N. Kuzmin, V. A. Lisinetskii, V. A. Orlovich, A. A. Demidovich, K. V. Yumashev, N. V. Kuleshov, H. J. Eichler, and M. V. Danailov, "Passively Q-switched 1.35- μm diode-pumped Nd:KGW laser with V:YAG saturable absorbers", Opt. Mater. 16, 349-352 (2001)
- [4.2] R. Fluck, R. Häring, R. Paschotta, R. Gini, H. Melchior, and U. Keller, "Eyesafe pulsed microchip laser using semiconductor saturable absorber mirrors", Appl. Phys. Lett. 72, 3273-3275 (1998)
- [4.3] R. D. Stultz, V. Leyva, and K. Spariosu, "Short pulse, high-repetition rate, passively Q-switched Er:yttrium-aluminum-garnet laser at 1.6 microns," Appl. Phys. Lett. 87, 241118 (2005)
- [4.4] V. G. Savitski, N. N. Posnov, P. V. Prokoshin, A. M. Malyarevich, K. V. Yumashev, M. I. Demchuk, and A. A. Lipovski, "PbS-doped phosphate glasses saturable absorbers for 1.3- μm neodymium lasers", Appl. Phys. B 75, 841-846

- (2002)
- [4.5] D. L. Spies, "Highly efficient neodymium:yttrium aluminum garnet laser end-pumped by a semiconductor laser array," *Appl. Phys. Lett.* **47**, 74-76 (1985)
- [4.6] M. Kuznetsov, F. Hakimi, R. Sprague, and A. Mooradian, "Design and characteristics of high-power (>0.5-W CW) diode-pumped vertical-external-cavity surface-emitting semiconductor lasers with circular TEM₀₀ beams," *IEEE J. Sel. Topics Quantum Electron.* **5**, 561-573 (1999)
- [4.7] A. C. Tropper, H. D. Foreman, A. Garnache, K. G. Wilcox, and S. H. Hoogland, "Vertical-external-cavity semiconductor lasers," *J. Phys. D: Appl. Phys.* **39**, R74-R85 (2004)
- [4.8] M. Kondow, K. Uomi, A. Niwa, T. Kitatani, S. Watahiki, and Y. Yazawa, "GaInNAs: A novel material for long-wavelength-range laser diodes with excellent high-temperature performance," *Japan. J. Appl. Phys.* **35**, 1273-1275 (1996)
- [4.9] J. M. Hopkins, S. A. Smith, C. W. Jeon, H. D. Sun, D. Burns, S. Calvez, M. D. Dawson, T. Jouhti, and M. Pessa, "0.6 W CW GaInAs vertical external-cavity surface-emitting laser operating at 1.32 μm," *Electron. Lett.* **4**, 30-31 (2004).
- [4.10] C. E. Zah, R. Bhat, B. N. Pathak, F. Favire, W. Lin, M. C. Wang, N. C. Andreadakis, D. M. Hwang, M. A. Koza, T. P. Lee, Z. Wang, D. Darby, D. Flanders, and J. J. Hsieh, "High-performance uncooled 1.3-μm Al_xGa_yIn_{1-x-y}As/InP strained-layer quantum-well lasers for subscriber loop applications," *IEEE J. Quantum Electron.*, vol. QE-30, 511–521 (1994).
- [4.11] J. Minch, S. H. Park, T. Keating, and S. L. Chuang, "Theory and experiment of In_{1-x-y}Ga_xAs_yP_{1-y} and In_{1-x-y}Ga_xAl_yAs long-wavelength strained quantum-well lasers," *IEEE J. Quantum Electron.*, vol. 35, 771–782 (1999).
- [4.12] S. R. Selmic, T. M. Chou, J. P. Sih, J. B. Kirk, A. Mantie, J. K. Butler, D. Bour, and G. A. Evans, "Design and characterization of 1.3-μm AlGaInAs-InP multiple-quantum-well lasers," *IEEE J. Sel. Topics Quantum Electron.*, vol. 7, 340–349 (2001).
- [4.13] N. Nishiyama, C. Caneau, B. Hall, G. Guryanov, M. H. Hu, X. S. Liu, M.-J. Li, R. Bhat, and C. E. Zah, "Long-wavelength vertical-cavity surface-emitting lasers on InP with lattice matched AlGaInAs-InP DBR grown by MOCVD," *IEEE J. Sel. Topics Quantum Electron.*, vol. 11, 990–998 (2005).

- [4.14] O. Hanaizumi, K. T. Jeong, S. Y. Kashiwada, I. Syuaib, K. Kawase, and S. Kawakami, "Observation of gain in an optically pumped surface-normal multiple-quantum-well optical amplifier," Opt. Lett. 21, 269-271 (1996).
- [4.15] F. Sanchez, M. Brunel, and K. Aït-Ameur, "Pump-saturation effects in end-pumped solid-state lasers," J. Opt. Soc. Am. B 15, 2390-2394 (1998).
- [4.16] K. S. Kim et al., "1060-nm vertical-external-cavity surface-emitting lasers with an optical-to-optical efficiency of 44% at room temperature," Appl. Phys. Lett. 88, 091107 (2006)
- [4.17] J.E. Hastie, C.W. Jeon, D. Burns, J.M. Hopkins, S. Calvez, R. Abram, and M.D. Dawson, "Thermal management of VECSELs with the use of intra-cavity crystalline heatspreaders," University of Strathclyde
- [4.18] Y. Tzuk, A. Tal, S. Goldring, Y. Glick, E. Lebiush, G. Kaufman, and R. Lavi, "Diamond Cooling of High-Power Diode-Pumped Solid-State Lasers," IEEE J. Quantum Electron. 40, 3, 262–269 (2004)

Chapter 5 - Second-order Effects

- [5.1] E. Abrahams, P. W. Anderson, D. C. Licciardello, and T. V. Ramakrishnan, Phys. Rev. Lett. 42, 673 (1979); S. John, Phys. Rev. Lett. 53, 2169 (1984); P. W. Anderson, Philos. Mag. B 52, 505 (1985); F. Scheffold and G. Maret, Phys. Rev. Lett. 81, 5800 (1998).
- [5.2] P. Sheng, *Introduction to Wave Scattering, Localization, and Mesoscopic Phenomena* (Academic Press, New York, 1995).
- [5.3] V. L. Kuzmin and V. P. Romanov, Phys. Usp. 39, 231 (1996).
- [5.4] M. C. W. van Rossum and Th. M. Nieuwenhuizen, Rev. Mod. Phys. 71, 313 (1999).
- [5.5] B. A. van Tiggelen, Phys. Rev. Lett. 75, 422 (1995); B. A. van Tiggelen and R. Maynard, in *Wave Propagation in Complex Media*, edited by G. Papanicolaou. IMA volumes in Mathematics and its Applications vol. 96 (Springer, New York, 1998).
- [5.6] M. P. van Albada and A. Lagendijk, Phys. Rev. Lett. 55, 2692 (1985); P. E. Wolf and G. Maret, Phys. Rev. Lett. 55, 2696 (1985).
- [5.7] D. S. Wiersma, M. P. van Albada, B. A. Van Tiggelen, and A. Lagendijk, Phys.

- Rev. Lett. 74, 4193 (1995).
- [5.8] G. Labeyrie, F. De Tomasi, J. C. Bernard, C. A. Müller, C. Miniatura, and R. Kaiser, Phys. Rev. Lett. 83, 5266 (1999); Y. Bidel, B. Klappauf, J. C. Bernard, D. Delande, G. Labeyrie, C. Miniatura, D. Wilkowski, and R. Kaiser, Phys. Rev. Lett. 88, 203902 (2002).
- [5.9] I. Freund, M. Rosenbluh, R. Berkovitsm and M. Kaveh, Phys. Rev. Lett. 61, 1214 (1988).
- [5.10] D. S. Wiersma, M. P. van Albada, and A. Lagendijk, Phys. Rev. Lett. 75, 1739 (1995).
- [5.11] R. Sapienza, S. Mujumdar, C. Cheung, A. G. Yodh, and D. Wiersma, Phys. Rev. Lett. 92, 33903 (2004); L. V. Kuzmin, V. P. Romanov, and L. A. Zubkov, Phys. Rev. E 54, 6798 (1996).
- [5.12] M. Gurioli, F. Bogani, L. Cavigli, H. Gibbs, G. Khitrova, and D. S. Wiersma, Phys. Rev. Lett. 94, 183901 (2005).
- [5.13] P. Pradhan and S. Sridhar, Phys. Rev. Lett. 85, 2360 (2000); A. Kudrolli, V. Kidambi, and S. Sridhar, Phys. Rev. Lett. 75, 822 (1995); M. Barth, U. Kuhl, and H. J. Stöckmann, Phys. Rev. Lett. 82, 2026 (1999).
- [5.14] A. R. Tunyagi, M. Ulex, and K. Betzler, Phys. Rev. Lett. 90, 243901 (2003); P. Xu, S. H. Ji, S. N. Zhu, X. Q. Yu, J. Sun, H. T. Wang, J. L. He, Y. Y. Zhu, and N. B. Ming, Phys. Rev. Lett. 93, 133904 (2004)
- [5.15] Y. V. Fyodorov and A. D. Mirlin, Phys. Rev. B 51, 13403 (1995); A. D. Mirlin, Phys. Rep. 326, 259 (2000); V. Prigodin and B. Altshuler, Phys. Rev. Lett. 80, 1944 (1998); K. B. Efetov, *Supersymmetry in Disorder and Chaos* (Cambridge University press, 1997).
- [5.16] M. V. Pack, D. J. Armstrong, A. V. Smith, G. Aka, B. Ferrand, and D. Pelenc, J. Opt. Soc. Am. B 22, 417 (2005).
- [5.17] M. Lefeld-Sosnowska, E. Olszyńska, A. Pajaczkowska, and A. Kłos, J. Cryst. Growth 262, 388 (2004).
- [5.18] P. O'Connor, J. Gehlen, and E. J. Heller, Phys. Rev. Lett. 58, 1296 (1987).
- [5.19] S. Feng, C. Kane, P. A. Lee, and A. D. Stone, Phy. Rev. Lett. 61, 834 (1988).
- [5.20] P. Sebbah, B. Hu, A. Z. Genack, R. Pnini, and B. Shapiro, Phys. Rev. Lett. 88,

- 123901 (2002).
- [5.21] S. H. Chang, A. Taflove, A. Yamilov, A. Burin, and H. Cao, Opt. Lett. 29, 917 (2004).
- [5.22] K. Staliunas, and V. J. Sanchez-Morcillo, “Transverse Patterns in Nonlinear Optical Resonators”, Springer Tracts in Modern Physics, 183, ISBN: 3-540-00434-3 (2005)
- [5.23] E. Gregor, D. E. Nieuwsma, and R. D. Stultz, “20 Hz eyesafe laser rangefinder for air defense,” Proc. SPIE, 1207, 124-134 (1990).
- [5.24] J. E. Nettleton, B. W. Schilling, D. N. Barr, and J. S. Lei, “Monoblock laser for a low-cost, eyesafe, microlaser range finder,” Appl. Opt. 39, 2428-2432 (2000).
- [5.25] S. Kück, K. Petermann, U. Pohlmann, U. Schönhoff, and G. Huber, “Tunable room-temperature laser action of Cr⁴⁺-doped Y₃Sc_xAl_{5-x}O₁₂,” Appl. Phys. B 58, 153-156 (1994).
- [5.26] N. V. Kuleshov, A. A. Lagatsky, A. V. Podlipensky, V. P. Mikhailov, A. A. Kornienko, E. B. Dunina, S. Hartung, and G. Huber, “Fluorescence dynamics, excited-state absorption and stimulated emission of Er³⁺ in KY(WO₄)₂,” J. Opt. Soc. Am. B 15, 1205-1212 (1998).
- [5.27] I. Sokólska, E. Heumann S. Kück, and T. Łukasiewicz, “Laser oscillation of Er³⁺:YVO₄ and Er³⁺, Yb³⁺:YVO₄ crystals in the spectral range around 1.6 μm,” Appl. Phys. B 71, 893-896 (2000).
- [5.28] A. Sennaroglu, “Broadly tunable Cr⁴⁺-doped solid-state lasers in the near infrared and visible,” Prog. in Quantum Electron. 26, 287–352 (2002).
- [5.29] P. Černý, H. Jelínkova, P. Zverev, and T. T. Basiev, “Solid state lasers with Raman frequency conversion,” Prog. in Quantum Electron. 28, 113-143 (2004).
- [5.30] Y. F. Chen, “Compact efficient all-solid-state eye-safe laser with self-frequency Raman conversion in a Nd:YVO₄ crystal,” Opt. Lett. 29, 2172-2174 (2004).
- [5.31] Y. F. Chen, “Efficient 1521-nm Nd :GdVO₄ Raman laser,” Opt. Lett. 29, 2632-2634 (2004).
- [5.32] J. T. Murray, R. C. Powell, D. Smith, W. Austin, and R. A. Stolzenberger, “Generation of 1.5 m radiation through intracavity solid-state Raman shifting in Ba(NO₃)₂ nonlinear crystals,” Opt. Lett. 20, 1017-1019 (1995).

- [5.33] Y. Yashkir and H. M. van Driel, "Passively Q-switched 1.57- μm intracavity optical parametric oscillator," *Appl. Opt.* 38, 2554-2559 (1999).
- [5.34] A. Agnesi, S. Dell'Acqua, and G. Reali, "Diode-pumped quasi-cw intracavity optical parametric oscillator at 1.57 μm with efficient pulse shortening," *Appl. Phys. B* 70, 751-753 (2000).
- [5.35] R. Dabu, C. Fenic, and A. Stratan, "Intracavity pumped nanosecond optical parametric oscillator emitting in the eye-safe range," *Appl. Opt.* 40, 4334-4340 (2001).
- [5.36] W. Zendzian, J. K. Jabczyński, and J. Kwiatkowski, "Intracavity optical parametric oscillator at 1572-nm wavelength pumped by passively Q-switched diode-pumped Nd:YAG laser," *Appl. Phys. B* 76, 355-358 (2003).
- [5.37] Y. F. Chen, S. W. Chen, Y. C. Chen, Y. P. Lan, and S. W. Tsai, "Compact efficient intracavity optical parametric oscillator with a passively Q-switched Nd:YVO₄/Cr⁴⁺:YAG laser in a hemispherical cavity," *Appl. Phys. B* 77, 493-495 (2003).
- [5.38] B. W. Schilling, S. R. Chinn, A. D. Hays, L. Goldberg, and C. W. Trussell, "End-pumped 1.5 μm monoblock laser for broad temperature operation," *Appl. Opt.* 45, 6607-6615 (2006).
- [5.39] Y. F. Chen, S. W. Chen, L. Y. Tsai, Y. C. Chen, C. H. Chien, "Efficient sub-nanosecond intracavity optical parametric oscillator pumped with a passively Q-switched Nd:GdVO₄ laser," *Appl. Phys. B* 79, 823-825 (2004).
- [5.40] Y. F. Chen and L. Y. Tsai, "Comparison between shared and coupled resonators for passively Q-switched Nd:GdVO₄ intracavity optical parametric oscillators," *Appl. Phys. B* 82, 403-406 (2006).
- [5.41] R. J. Beach, "Theory and optimization of lens ducts," *Appl. Opt.* 35, 2005-2015 (1996).
- [5.42] R. Fu, G. Wang, Z. Wang, E. Ba, G. Mu, and X. Hu, "Design of efficient lens ducts," *Appl. Opt.* 37, 4000-4003 (1998).
- [5.43] Yi-Shin Chen, Ch. 3 in "Intracavity Frequency Conversion in Diode-pumped Actively Q-switched Nd:YVO₄ Laser: From NIR to Visible and Violet," National Chiao Tung University, M.S. thesis (2004)
- [5.44] K. Staliunas and V.J. Sanchez-Morcillo, Ch. 1 in "Transverse Patterns in

- Nonlinear Optical Resonators,” ISBN: 3-540-00434-3 (Springer-Verlag, 2003)
- [5.45] K. Pennicott, “Random laser feels the heat (D Wiersma and S Cavalieri 2001 Nature 414 708),” <http://physicsweb.org/articles/news/5/12/5>
- [5.46] I.S. Aranson, L. Kramer, “The world of the complex Ginzburg-Landau equation,” Rev. of Modern Phys., 74, 99-143 (2002)
- [5.47] G. J. Zhang, S. Horinouchi, T. Kinoshita, and K. Sasaki, “Theoretical analysis of the spatial phase-matching loci for second-harmonic generation and multiwave-mixing interactions,” Appl. Opt. 34, 24, 5301-5311 (1995)
- [5.48] Ling-Yi Tsai, “Passively Q-switched intracavity optical parametric oscillator,” National Chiao Tung University, M.S. thesis (2005)
- [5.49] J. Nettleton, D. Barr, B. Schilling, J. Lei, and S.M. Goldwasser, “Micro-Laser Range Finder Development: Using the Monolithic Approach,” <http://www.repairfaq.org/sam/lr/> (1999)
- [5.50] D.L. Kaufman, I. Kosztin, and K. Schulten, “Expansion method for stationary states of quantum billiards,” Am. J. Phys. 67, 133 (1999)
- [5.51] K.F. Huang, Y.F. Chen, H.C. Lai, and Y.P. Lan, “Observation of the Wave Function of a Quantum Billiard from the Transverse Patterns of Vertical Cavity Surface Emitting Lasers,” Phys. Rev. Lett. 89, 224102 (2002)
- [5.52] C.C. Chen, C.C. Liu, K.W. Su, T.H. Lu, Y.F. Chen, and K.F. Huang, “Statistical properties of experimental coherent waves in microcavity lasers: Analogous study of quantum billiard wave functions,” Phys. Rev. E 76, 1 (2007)

Chapter 6 - Third-order Effects

- [6.1] T.T. Basiev, V.V. Osiko, A.M. Prokhorov and E.M. Dianov, “Crystalline and Fiber Raman Lasers,” Topics Appl. Phys. 89, 351 (2003)
- [6.2] T.T. Basiev, R.C. Powell, “Solid state Raman lasers” in *Handbook of Laser Technology and Applications*, B1.7, 469 (2003)
- [6.3] H.M. Pask, “The design and operation of solid-state Raman lasers”, Prog. Quantum Electron. 27, 3 (2003)
- [6.4] R.G. Zevrev, T.T. Basiev, and A.M. Prokhorov, “Stimulated Raman scattering of laser radiation in Raman crystals”, Opt. Mater. 11, 335 (1999)

- [6.5] T.T. Basiev, A.A. Sobol, P.G. Zverev, L.I. Ivelleva, V.V. Osiko, R.C. Powell, “Raman spectroscopy of crystals for stimulated Raman scattering,” Opt. Mater. 11, 4, 307-314 (1999)
- [6.6] P. Cerný, P.G. Zverev, H. Jelímková, T.T. Basiev, “Efficient Raman shifting of picosecond pulses using BaWO₄ crystal,” Opt. Commun. 177, 397-404 (2000)
- [6.7] J.T. Murray, W.L. Austin, and R.C. Powell, in Advanced Solid-State Lasers, W. R. Bosenberg and M. J. Fejer, eds., Vol. 19 of OSA Trends in Optics and Photonics Series (Optical Society of America, Washington, D.C., 1998), pp. 129–135.
- [6.8] J.T. Murray, W.L. Austin, and R.C. Powell, in Advanced Solid-State Lasers, M. J. Fejer, H. Injeyan, and U. Keller, eds., Vol. 26 of OSA Trends in Optics and Photonics Series (Optical Society of America, Washington, D.C., 1999), pp. 575–578.
- [6.9] C. He and T.H. Chyba, “Solid-state barium nitrate Raman laser in the visible region”, Opt. Commun. 135, 273 (1997).
- [6.10] J.T. Murray, W.L. Austin, and R.C. Powell, “Intracavity Raman conversion and Raman beam cleanup”, Opt. Mater. 11, 353 (1999).
- [6.11] T.T. Basiev, S.V. Vassiliev, M.E. Doroshenko, and V.V. Osiko, V.M. Puzikov and M.B. Kosmyna, “Laser and self-Raman-laser oscillations of PbMoO₄:Nd³⁺ crystal under laser diode pumping,” Opt. Lett. 31, 1, 65-67 (2006); T.T. Basiev, S.V. Vassiliev, M.E. Doroshenko, and V.V. Osiko, “Laser oscillation and self-Raman frequency conversion in Nd:PbMoO₄ and Nd:SrMoO₄ crystals under laser diode pumping,” Technical Digest, Advanced Solid State Photonics, Jan. 29, 2006, Lake Tahoe, OSA, TuB21 (2006)
- [6.12] G. Eckhardt, D.P. Bortfeld, and M. Geller, “Stimulated emission of Stokes and anti-Stokes Raman lines from diamond, calcite, and α -sulfur single crystals”, Appl. Phys. Lett. 3, 137 (1963).
- [6.13] A.S. Eremenko, S.N. Karpukhin, and A.I. Stepanov, “Stimulated Raman scattering of the second harmonic of a neodymium laser in nitrate crystals”, Sov. J. Quantum Electron. 10, 113 (1980).
- [6.14] E.O. Ammann and C.D. Decker, “0.9-W Raman oscillator”, J. Appl. Phys. 48, 1973 (1977).

- [6.15] A.S. Grabtachikov, A.N. Kuzmin, V.A. Lisinetskii, V.A. Orlovich, G.I. Ryabtsev, and A. A. Demidovich, "All solid-state diode-pumped Raman laser with self-frequency conversion", *Appl. Phys. Lett.* 75, 3742 (1999).
- [6.16] J. Findeisen, H.J. Eichler, and A.A. Kaminskii, "Efficient picosecond PbWO₄ and two-wavelength KGd(WO₄)₂ Raman lasers in the IR and visible", *IEEE J. Quantum Electron.* 35, 173 (1999).
- [6.17] T.T. Basiev, A.A. Sobol, P.G. Zverev, L.I. Ivleva, V.V. Osiko, R.C. Powell, "Laser material for stimulated Raman scattering," Patent of Russian Federation, 2178938 C1 1, pending 25.04.2000, Jb2000110182/28 bul. 3, 27.01.2002
- [6.18] T.T. Basiev, P.G. Zverev, A.A. Sobol, V.V. Skornyakov, L.I. Ivleva, N.M. Polozkov, V.V. Osiko, "Stimulated Raman scattering in crystals of alkali-halide tungstates," *Sov. J. Quantum Electron.* 30, Jb1, 55-59 (2000)
- [6.19] P.G. Zverev, T.T. Basiev, M.E. Doroshenko, V.V. Osiko, "Barium tungstate Raman laser – a new coherent source for sodium star experiments," *Trends in Optics and Photonics* 34, Advanced Solid State Lasers, H. Injeyan, U. Keller, and C. Marshall, eds.(OSA Washington D.C.) 348-354 (2000)
- [6.20] P.G. Zverev, T.T. Basiev, L.I. Ivleva, V.V. Osiko, N.M. Polozkov, I.S. Voronina, "Raman laser on strontium tungstate crystal," *Trends in Optics and Photonics* 68, (OSA Washington D.C.) 70-73 (2002)
- [6.21] P. Cerný, H. Jelínková, T.T. Basiev, and P.G. Zverev, "Highly efficient picosecond Raman generators based on the BaWO₄ crystal in the near infrared, visible, and ultraviolet," *IEEE J. Quantum Electron.* 38, 11, 1471-1478 (2002)
- [6.22] A.A. Lagatsky, A. Abdolvand, and N.V. Kuleshov, "Passive Q switching and self-frequency Raman conversion in a diode-pumped Yb:KGd(WO₄)₂ laser", *Opt. Lett.* 25, 616 (2000)
- [6.23] K. Andryunas, Y. Vishakas, V. Kabelka, I.V. Mochalov, A.A. Pavlyuk, G.T. Petrovskii, V. Syrus, *JETP Lett.* 42, 333 (1985); K. Andryunas, Y. Vishakas, V. Kabelka, I.V. Mochalov, A.A. Pavlyuk, N.V. Ionina, V. Syrus, Patent of USSR #1227074
- [6.24] A.M. Ivanyuk, M.A. Ter-Pogosyan, P.A. Shakhverdov, V.D. Belyaev, V.L. Ermolaev, H.P. Tikhonova, "Picosecond radition pulses at intracavity stimulated Raman scattering inside the active element of neodymium laser," *Opt. and Spectroscopy* 59, 950-952 (1985)

- [6.25] Y.K. Vishakas, I.V. Mochalov, A.V. Mikhailov, R.F. Klevtsova, A.V. Lyubimov, "Crystal structure and Raman scattering in KGd(WO₄)₂ crystals," Lietuvos fizikos rinkinys 28, 2, 224-234 (1988)
- [6.26] I.V. Mochalov, "Laser and nonlinear properties of the potassium gadolinium tungstate laser crystal KGd(WO₄)₂:Nd³⁺ (KGW:Nd)," Opt. Eng. 36, 6, 1660-1669 (1997)
- [6.27] A.M. Ivanyuk, V.A. Sandulenko, and M.A. Ter-Pogosyan, P.A. Shakhverdov, V.G. Chervinskii, A.V. Lukin, V.L. Ermolaev, "Intracavuty stimulated Raman scattering in a nanosecond neodymium laser based on potassium-gadolinium tungstate," Opt. and Spectroscopy 62, 5, 961-962 (1987)
- [6.28] V.A. Berenberg, S.N. Karpukhin, I.V. Mochalov, "SRS of nanosecond pulses in KGd(WO₄)₂ crystals," Sov. J. Quantum Electron. 14, 9, 1849-1850 (1987)
- [6.29] J. Findeisen, H.J. Eichler, and P. Peuser, "Self-stimulating, transversally diode pumped Nd³⁺: KGd(WO₄)(2) Raman laser", Opt. Commun. 181, 129 (2000).
- [6.30] W. Chen, Y. Inagawa, T. Omatsu, M. Tateda, N. Takeuchi, and Y. Usuki, "Diode-pumped, self-stimulating, passively Q-switched Nd³⁺: PbWO₄ Raman laser", Opt. Commun. 194, 401 (2001).
- [6.31] A. A. Kaminskii, K. Ueda, H.J. Eichler, Y. Kuwano, H. Kouta, S.N. Bagaev, T.H. Chyba, J.C. Barnes, G.M. A. Gad, T. Murai, and J. Lu, "Tetragonal vanadates YVO₄ and GdVO₄ – new efficient $\chi^{(3)}$ -materials for Raman lasers", Opt. Commun. 194, 201 (2001).
- [6.32] Y.F. Chen, M.L. Ku, L.Y. Tsai, and Y.C. Chen, "Diode-pumped passively Q-switched picosecond Nd:GD_xY_{1-x}VO₄ self-stimulated Raman laser", Opt. Lett. 29, 19, 2279-2281 (2004)
- [6.33] Y.F. Chen, "Efficient subnanosecond diode-pumped passively Q-switched Nd : YVO₄ self-stimulated Raman laser", Opt. Lett. 29, 11, 1251 (2004).
- [6.34] Y.F. Chen, "High-power diode-pumped actively Q-switched Nd : YVO₄ self-Raman laser: influence of dopant concentration", Opt. Lett. 29, 16, 1915 (2004)
- [6.35] T.T. Basiev, S.V. Vassiliev, V.A. Konjushkin, V.V. Osiko, A.I. Zagumennyi, Y.D. Zavartsev, S.A. Kutovoi, I.A. Shecherbakov, "Diode pumped 500-picosecond Nd:GdVO₄ Raman laser," Laser Phys. Lett. 1, 5, 237-240 (2004)

- [6.36] Y.F. Chen, “Compact efficient self-frequency Raman conversion in diode-pumped passively Q-switched Nd : GdVO₄ laser”, *Appl. Phys. B* 78, 6, 685 (2004).
- [6.37] Y.F. Chen, “Efficient 1521-nm Nd : GdVO₄ Raman laser”, *Opt. Lett.* 29, 22, 2632 (2004).
- [6.38] Y.P. Lan, Y.F. Chen, and S.C. Wang, “Repetition-rate dependence of thermal loading in diode-end-pumped Q-switched lasers: influence of energy-transfer upconversion”, *Appl. Phys. B* 71, 27 (2000).
- [6.39] P.G. Zverev, A.Y. Karasik, A.A. Sobol, D.S. Chunaev, T.T. Basiev, A.I. Zagumennyi, Y.D. Zavartsev, S.A. Kutowoi, V.V. Osiko, I.A. Shcherbakov, “Stimulated Raman scattering of picosecond pulses in GdVO₄ and YVO₄ crystals,” Advanced Solid State Photonics Conference, Technical digest, Feb. 1-4, 2004, Santa Fe, USA, WB9. OSA TOPS ASSL, TuB10, 25-27
- [6.40] S. N. Karpukhin and A. I. Stepanov, “SRS oscillation in the cavity in Ba(NO₃)₂, NaNO₃ and CaCO₃ crystals,” *Sov. J. Quantum Electron.* 16, 1027- 1032 (1986)
- [6.41] H. M. Pask and J. A. Piper, “ Diode-pumped LiIO₃ intracavity Raman lasers,” *IEEE J. Quantum Electron.*, 36, 949 (2000)
- [6.42] T. T. Basiev, A. A. Sobol, P. G. Zverev, V. V. Osiko, and R. C. Powell, “Comparative spontaneous Raman spectroscopy of crystals for Raman lasers,” *Appl. Opt.* 38, 594-598 (1999)
- [6.43] A. V. Gulin, G. I. Narkhova, N. S. Ustimenko, “Multiwave generation of the Stokes components in lasers with an SRS self-converter based on an Nd³⁺:KGd(WO₄)₂ crystal,” *Quantum Electron.* 28, 804-805 (1998)
- [6.44] K. Andryunas, Yu. Vishchakas, V. Kabelka, I. V. Mochalov, A. A. Pavlyuk, G. T. Petrovskii, and V. Syrus, “SRS-self-conversion of Nd³⁺ laser emission in tungstate crystals,” *JETP Lett.* 42, 410-412 (1985)
- [6.45] J. Findeisen, H. J. Eichler, and A. A. Kaminskii, “Efficient picosecond PbWO₄ and two-wavelength KGd(WO₄)₂ Raman lasers in the IR and visible,” *IEEE J. Quantum Electron.* 35, 173-178 (1999)
- [6.46] A. Kaminskii, H. Eichler, K. Ueda, N. Klassen, B. Redkin, L. Li, J. Findeisen, D. Jaque, J. Garcia-Sole, J. Fernandez, and R. Balda, “Nd³⁺-doped and undoped tetragonal PbWO₄, NaY(WO₄)₂, CaWO₄, and undoped monoclinic ZnWO₄

- and CdWO₄ as laser- and SRS-active crystals: low-threshold pulsed and CW near-IR stimulated emission, multiple Stokes and anti-Stokes generation, and efficient picosecond Raman laser," Appl. Opt., vol. 38, 4533-4547 (1999)
- [6.47] P. Černý, H. Jelínková, P. G. Zverev, and T. T. Basiev, "Solid state lasers with Raman frequency conversion," Prog. Quantum Electron., 28 113-143 (2004)
- [6.48] P. Černý, W. Žendzian, J. Jabczyński, H. Jelínková, J. Šulc, and K. Kopczyński, "Efficient diode-pumped passively Q-switched Raman laser on barium tungstate crystal," Opt. Commu. 209, 403-409 (2002)
- [6.49] Y.F. Chen, "Efficient subnanosecond diode-pumped passively Q-switched Nd : YVO₄ self-stimulated Raman laser", Opt. Lett. 29, 11, 1251 (2004).
- ## Chapter 7 - Summary
- [7.1] L.V. Butov, C.W. Lai, A.L. Ivanov, A. C. Gossard and D.S. Chemla, "Towards Bose-Einstein condensation of excitons in potential traps," Nature 417, 47-52 (2002)
- [7.2] L.V. Butov, A.C. Gossard and D.S. Chemla, "Macroscopically ordered state in an exciton system," Nature 418, 751-754 (2002)
- [7.3] D. Snoke, S. Denev, Y. Liu, L. Pfeiffer, and K. West, "Long-range transport in excitonic dark states in coupled quantum wells," Nature 418, 754-757 (2002)
- [7.4] D. Snoke, S. Denev, Y. Liu, S. Simon, R. Rapaport, L. Pfeiffer, and K. West, "Moving beyond a simple model of luminescence rings in quantum well structures," J. Phys.: Condens Matter 16, S3621-S3627 (2004)
- [7.5] A.T. Hammack, M.Griswold, L.V. Butov, L.E. Smallwood, A.L. Ivanov, A. C. Gossard, "Trapping of Cold Excitons in Quantum Well Structures with Laser Light," Phys. Rev. Lett. 96, 227402 (2006)
- [7.6] Z. Vörös, D.W. Snoke, L. Pfeiffer, and K. West, "Trapping Excitons in a Two-Dimensional In-Plane Harmonic Potential: Experimental Evidence for Equilibration of Indirect Excitons," Phys. Rev. Lett. 97, 016803 (2006)
- [7.7] H. Deng, G. Weihs, C. Santori, J. Bloch, Y. Yamamoto, "Condensation of Semiconductor Microcavity Exciton Polaritons," Science 298, 5591, 199 – 202 (2002)
- [7.8] H. Deng, G. Weihs, D. Snoke, J. Bloch, and Y. Yamamoto, "Polariton lasing vs.

- photon lasing in a semiconductor microcavity," PNAS 100, 15318-15323 (2003)
- [7.9] P. Pradhan and S. Sridhar, "Correlations due to Localization in Quantum Eigenfunctions of Disordered Microwave Cavities" Phys. Rev. Lett. 85, 2360 (2000)
- [7.10] L. Zenteno, "High-power double-clad fiber lasers," J. Lightwave Technol. 11, 1435-1446 (1993)
- [7.11] M. J. F. Digonnet, *Rare-Earth-Doped Fiber Lasers and Amplifiers*, 2nd ed. (Marcel Dekker, 2001)

Publication List

- [J1] H.C. Lai, A. Li, **K.W. Su**, M.L. Ku, Y.F. Chen, K.F. Huang, "InAs/GaAs quantum-dot saturable absorbers for diode-pumped passively Q-switched Nd-doped 1.3- μ m lasers," Opt. Lett. **30**, 5, 480-482 (2005)
- [J2] **K.W. Su**, H.C. Lai, A. Li, Y.F. Chen, K.F. Huang, "InAs/GaAs quantum-dot saturable absorbers for diode-pumped self-starting continuous mode-locked Nd-doped 1.3- μ m laser," Opt. Lett. **30**, 12, 1482-1484 (2005)
- [J3] Y.F. Chen, M.L. Ku, **K.W. Su**, "High-power efficient tunable Nd : GdVO₄ laser at 1083 nm," Opt. Lett. **30**, 16, 2107-2109 (2005)
- [J4] Y.F. Chen, Y.S. Chen, T.H. Ou, **K.W. Su**, "Compact efficient diode-pumped Nd : YVO₄ Q-switched blue laser with intracavity frequency tripling," Appl. Phys. B **81**, 4, 517-520 (2005)
- [J5] Y.F. Chen, T.H. Lu, **K.W. Su**, K.F. Huang, "Quantum signatures of nonlinear resonances in mesoscopic systems Efficient extension of localized wave functions," Phys. Rev. E **72**, 056210 (2005)
- [J6] Y.F. Chen, **K.W. Su**, H.J. Zhang, J.Y. Wang, M.H. Jiang, "Efficient diode-pumped actively Q-switched Nd:YAG/BaWO₄ intracavity Raman laser," Opt. Lett. **30**, 24, 3335-3337 (2005)
- [J7] Y.F. Chen, **K.W. Su**, T.H. Lu, K.F. Huang, "Manifestation of weak localization and long-range correlation in disordered wave functions from conical second harmonic generation," Phys. Rev. Lett. **96**, 3, 033905 (2006)
- [J8] Y.F. Chen, T.H. Lu, **K.W. Su**, K.F. Huang, "Devil's Staircase in Three-Dimensional Coherent Waves Localized on Lissajous Parametric

- Surfaces," Phys. Rev. Lett. **96**, 21, 213902 (2006)
- [J9] **K.W. Su**, S.C. Huang, A. Li, S.C. Liu, Y.F. Chen, and K.F. Huang, "High-peak-power AlGaInAs quantum wells 1.3- μ m laser pumped by a diode-pumped actively Q-switched solid-state laser," Opt. Lett. **31**, 13, 2009-2011 (2006)
- [J10] A. Li, S.C. Liu, **K.W. Su**, Y.L. Liao, S.C. Huang, Y.F. Chen, K.F. Huang, "InGaAsP quantum-wells saturable absorber for diode-pumped passively Q-switched 1.3- μ m lasers," Appl. Phys. B **84**, 3, 429-431 (2006)
- [J11] J.Y. Huang, H.C. Liang, **K.W. Su**, H.C. Lai, Y.F. Chen and K.F. Huang, "InGaAs quantum-well saturable absorbers for a diode-pumped passively Q-switched Nd:YAG laser at 1123 nm," Appl. Opt. **46**, 2, 239-242 (2007)
- [J12] J.Y. Huang, H.C. Liang, **K.W. Su**, and Y.F. Chen, "High power passively Q-switched ytterbium fiber laser with Cr⁴⁺:YAG as a saturable absorber," Opt. Express **15**, 473-479 (2007)
- [J13] H.C. Liang, J.Y. Huang, **K.W. Su**, H.C. Lai, Y.F. Chen, K.F. Huang, H.J. Zhang, J.Y. Wang, and M.H. Jiang, "Passively Q-switched Yb³⁺:YCa₄O(BO₃)₃ laser with InGaAs quantum wells as a saturable absorbers," Appl. Opt. **46**, (2007)
- [J14] Y.F. Chen, **K.W. Su**, Y.T. Chang, and W.C. Yen, "Compact efficient eye-safe intracavity optical parametric oscillator with a shared cavity configuration," Appl. Opt. **46**, (2007)
- [C1] 蘇冠暉, 張雅婷, 陳永富, 閻偉中, “利用完全重疊共振腔架構製造精巧與高效率的人眼安全腔內光學參數震盪器,” 95 年度國防科技學術合作計畫成果發表會 B-27, 論文集(上) pp. B178-B183 (2006)
- [C2] 黃哲彥, 梁興弛, 蘇冠暉, 陳永富, 林志平, “光纖雷射研製,” 95 年度國防科技學術合作計畫成果發表會 B-28, 論文集(上) pp. B184-B187 (2006)