

行政院國家科學委員會專題研究計畫 成果報告

網格動態系統的型及混沌現象(3/3)

計畫類別：整合型計畫

計畫編號：NSC91-2115-M-009-004-

執行期間：91年08月01日至93年07月31日

執行單位：國立交通大學應用數學系

計畫主持人：林松山

報告類型：完整報告

報告附件：國外研究心得報告

出席國際會議研究心得報告及發表論文

處理方式：本計畫可公開查詢

中 華 民 國 93 年 10 月 7 日

THE DEVIL'S STAIRCASE DIMENSIONS AND MEASURE-THEORETICAL ENTROPY OF MAPS WITH HORIZONTAL GAP

JUNG-CHAO BAN AND SONG-SUN LIN

Dedicated to Professor. S-N Chow on the occasion of his 60th birthday

ABSTRACT. This work elucidates the measure-theoretical entropy and dimensions of a unimodal map with a horizontal gap.

The measure-theoretical entropy and dimensions of the F_t (which is defined later) are shown to form a devil's staircase structure with respect to the gap size t . Pesin's formula for gap maps is also considered .

1. INTRODUCTION

This study addresses a special family of maps (gap maps), which arise in the communication with chaos in cellular neural networks. (Readers should refer to [BHL]). Such maps are constructed by cutting a gap into a given unimodal map f .

The most interesting invariants to be considered in determining whether a map is chaotic are topological entropy, measure-theoretical entropy, dimensions and Lyapunov exponents. In [ZB], the authors examined such maps and observed the devil's staircase structure of the entropy

Key words and phrases. Upper(lower) box Dimension, Hausdoff dimension , Devil's Staircase, Pesin's formula.

This research was supported in part by National Center for Theoretical Sciences.

This research was supported in part by the National Science Council of ROC on Taiwan, project number NSC 92-2115-M-009-018 and the Lee and MTI Center for Networking Research.

function with respect to the gap size. Then in [BHL], the authors applied kneading theory to prove rigorously this presence of such structure for a map f with strong transitivity. In [M1], Misiurewicz proved that this structure can be applied to more general maps by considering a dynamic mean rather than kneading theory.

Nature questions arise regarding what happens to the measure-theoretical entropy for certain measure for these families of maps, and what about their dimensions (Hausdorff, upper box and lower box dimension are considered).

This investigation addresses whether for some function f there exist some measure μ (defined later), where the measure-theoretical entropy $h_\mu(F_t)$ (F_t is a family of gap maps induced by f) with respect to this measure μ the devil's staircase structure remains; that is, $h_\mu(t) \equiv h_\mu(F_t)$ is a monotonic and continuous function. Furthermore, the union of the constant part of $h_\mu(t)$ is open and dense in parameter space, i.e., $\{t \in [0, 1] \mid h_\mu(t) \text{ is constant}\}$ is open and dense in $[0, 1]$.

Moreover, for a given f , the Hausdorff, lower box and upper box dimension can be determined. For these three dimensions, the devil's staircase structure also obtains. Additionally, these three dimensions are equal for some function f .

Similar results of the well-known Pesin's dimension theorem for these maps were obtained.

This paper is organized as follows. Section 2 provides some well-known definitions, results and the main theorem. Section 3 presents a geometric structure developed by Moran, Pesin and Weiss. Section 4 includes relevant proof of the main theorems.

2. PRELIMINARIES

This section presents some definitions and well-known theorems without proof. f is assumed to be a continuous unimodal map.

Definition 1. *A continuous map $f : [0, 1] \rightarrow [0, 1]$ is unimodal if there exist $c \in [0, 1]$ such that*

$$(i). \quad f(0) = f(1) = 0$$

(ii). $f(x)$ is monotonically increasing on $[0, c]$ and decreasing on $[c, 1]$.

c is called turning point herein.

Definition 2. *Tent map $f : [0, 1] \rightarrow [0, 1]$ with slope $\lambda > 0$ is defined*

$$T_\lambda(x) = \begin{cases} \lambda x & \text{if } 0 \leq x \leq \frac{1}{2} \\ \lambda(1-x) & \text{if } \frac{1}{2} \leq x \leq 1 \end{cases}.$$

Definition 3. *Given a continuous map f , for $t \in [0, 1]$, we define $U_t = \{x \mid f(x) > t\}$ and the gap map $F_t(x)$ with respect to $f(x)$ is*

$$F_t(x) = \begin{cases} t & \text{if } x \in U_t \\ f(x) & \text{if } x \in [0, 1] \setminus U_t \end{cases}.$$

Furthermore,

$$C(t) = C(F_t) \equiv \{x \in [0, 1] \mid f^n(x) \notin U_t, \text{ for all } n \geq 1\},$$

$$C_m(t) = C_m(F_t) \equiv \{x \in [0, 1] \mid f^k(x) \notin U_t, \text{ for } 1 \leq k \leq m\},$$

$$D(t) = D(F_t) \equiv \{x \in [0, 1] \mid f^k(x) \in U_t, \text{ for some } k \geq 1\}.$$

The box dimensions are defined as follows.

Definition 4. *For a compact set $A \subset \mathbb{R}^m$, the lower box dimension of A , $\underline{\dim}_B(A)$, is defined by*

$$\underline{\dim}_B(A) = \liminf_{\varepsilon \rightarrow 0} \frac{\log N_A(\varepsilon)}{\log \varepsilon^{-1}}$$

where $N_A(\varepsilon)$ is the least number of balls of radius ε needed to cover A . The upper box dimension of A is similarly defined, by

$$\overline{\dim}_B(A) = \limsup_{\varepsilon \rightarrow 0} \frac{\log N_A(\varepsilon)}{\log \varepsilon^{-1}}.$$

In particular, the lower and upper box dimension of $C(t)$ for $t \in (0, 1)$ are defined by

$$\underline{\dim}_B(t) \equiv \underline{\dim}_B(C(F_t)).$$

and

$$\overline{\dim}_B(t) \equiv \overline{\dim}_B(C(F_t)),$$

respectively.

Definition 5. Let X be a metric space with metric d . For $x \in X$ and $\rho > 0$, let

$$B_\rho(x) = \{y : d(x, y) \leq \rho\}.$$

The diameter of a cover κ of X is $\sup\{\text{diam}A : A \in \kappa\}$. Then the Hausdorff dimension of X is defined by

$$\dim_H(X) = \inf\{\alpha : \liminf_{\varepsilon \rightarrow 0} \sum_{\substack{A \in \kappa \\ \kappa: \text{cover of } X \\ \text{with } \text{diam}\kappa \leq \varepsilon}} (\text{diam}A)^\alpha = 0\}$$

In particular, $\dim_H(t) \equiv \dim_H(C(F_t))$.

The definition of topological entropy is defined as follows.

Definition 6. Let $f : X \rightarrow X$ be uniformly continuous on the metric space X . For $E, F \subset X$ we say that E (n, δ) -spans F (with respect to f), if for each $y \in F$ there is an $x \in E$ so that $d(f^k(x), f^k(y)) \leq \delta$ for all $0 \leq k \leq n$. We let $r_n(F, \delta) = r_n(F, \delta, f)$ denote the minimum

cardinality of a set which (n, δ) -spans F . If K is compact, then the continuity of f guarantees $r_n(F, \delta) < \infty$. For compact K we define

$$\bar{r}_f(K, \delta) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log r_n(K, \delta)$$

and

$$h(f, K) = \lim_{\delta \rightarrow 0} \bar{r}_f(K, \delta).$$

Finally let $h_{\text{top}}(f) = \sup_K h_{\text{top}}(f, K)$ where K varies over all compact subsets of X .

Here, some well-known theorems concerning topological entropy are presented without proof.

Theorem 1. *If f is a continuous map, then*

$$h_{\text{top}}(f) = \lim_{n \rightarrow \infty} \frac{\log \text{card}(P_n(f))}{n} = \lim_{n \rightarrow \infty} \frac{\log \text{lap}(f^n)}{n}$$

where $P_n(f)$ is the set of periodic points of period $\leq n$, $\text{card}(A)$ presents cardinality of A , and $\text{lap}(f^n)$ is the number laps of f^n , that is, the minimal number of intervals on which f^n is monotonic.

Proof. Ref[ALM]. □

The measure-theoretical entropy is defined as follows.

Definition 7. *Suppose $f : X \rightarrow X$ is a map on a metric space (X, d) , where d is a metric. Let μ be a Borel measure on X with $\mu(X) = 1$, and μ is f -invariant (i.e. $\mu(f^{-1}(A)) = \mu(A)$ for every Borel set A). One can then define a measure theoretic entropy $h_\mu(f)$ as follows: Call $\alpha = \{A_1, \dots, A_r\}$ a (finite) measurable partition of X if the A_i are disjoint measurable subsets of X covering X . Now set*

$$H_m(\alpha) = \sum_{1 \leq i_0, \dots, i_{m-1} \leq r} -\mu\left(\bigcap_{k=0}^{m-1} f^{-k} A_{i_k}\right) \log \mu\left(\bigcap_{k=0}^{m-1} f^{-k} A_{i_k}\right).$$

Then the limit $h_\mu(f, \alpha) = \lim_{m \rightarrow \infty} (1/m)H_m(\alpha)$ exists and one defines

$$h_\mu(f) = \sup\{h_\mu(f, \alpha) : \alpha \text{ is a finite measurable partition of } X\}.$$

The well-known theorem that relates the topological and measure-theoretical entropy is stated as follows.

Theorem 2. (Goodwyn). *Let (X, d) be a compact metric space, let $f : X \rightarrow X$ be continuous map and μ be a f -invariant Borel measure on X with $\mu(X) = 1$. Then $h_\mu(f) \leq h_{top}(f)$.*

The main results in the work of [BHL] and [M1] are included.

Theorem 3. (BHL & Misiurewicz) *If f is a unimodal map with $h_{top}(f) > 0$, the topological entropy function $h_{top}(t) = h_{top}(F_t)$ of the gap map F_t forms a devil's staircase function with respect to t ; that is, $h_{top}(t)$ is a monotonic, continuous function, and the constant parts of $h_{top}(t)$ is open and dense in parameter space, i.e., $\{t \in [0, 1] \mid h_{top}(t) \text{ is constant}\}$ is open and dense in $[0, 1]$.*

Then, the main theorem of this work is as follows.

Theorem A. *If f is a unimodal map with $h_{top}(f) > 0$ and let F_t be the gap map induced by f , then*

(i) $\underline{\dim}_B(t)$, $\overline{\dim}_B(t)$ and $\dim_H(t)$ are monotonic increasing functions.

(ii) *The union of the constant parts of each of the three dimensions are open and dense in parameter space; that is, $\{t \in [0, 1] \mid \underline{\dim}_B(t)$ ($\dim_H(t)$ and $\overline{\dim}_B(t)$, respectively) is constant} is open and dense in $[0, 1]$.*

Furthermore, *If f is a tent map with slope $\lambda > 1$, then*

(iii) $\underline{\dim}_B(t) = \overline{\dim}_B(t) = \dim_H(t)$ and is devil's staircase function with respect to t ,

(iv) there exist a measure μ (which is defined in Theorem 4) such that $h_\mu(t)$ of the gap map F_t forms a devil's staircase function with respect to t .

Theorem B. *Under the same assumption as in Theorem A. Let V be some maximal interval of the constant part of $h_{top}(t)$; V is also a the maximal constant part in $\underline{\dim}_B$, $\overline{\dim}_B$ and \dim_H .*

Furthermore, there exist $\bar{t} \in \partial V$ such that for all $t \in V$,

$$(2.1) \quad \underline{\dim}_B(t) \leq \frac{h_{top}(\bar{t})}{\log \lambda} \overline{\lim}_{n \rightarrow \infty} \left(\frac{\log \bar{\varepsilon}_n^{-1}}{\log \hat{\varepsilon}_n^{-1}} \right),$$

where $\lambda = \lim_{n \rightarrow \infty} |Df^n(\bar{t})|^{-\frac{1}{n}}$ the Lyapunov exponent, $\bar{\varepsilon}_n(t)$ is the minimum length of the elements in $C_n(t)$, and $\hat{\varepsilon}_n(t)$ is the maximum length of the elements in $C_n(t)$.

In addition, if f is a tent map with slope $\lambda > 1$, the inequality is actually equality; that is,

$$\underline{\dim}_B(t) = \frac{h_{top}(\bar{t})}{\log \lambda}.$$

3. MORAN-LIKE GEOMETRIC CONSTRUCTIONS

In this section, we recall the Moran-like geometric construction given by Y.Pesin and H.Weiss [P], [PW], and some theorems are stated without proof.

Let \sum_p^+ be the set of all one-sided infinite sequences (i_0, i_1, \dots) of p symbols, and σ be the shift map, i.e., $\sigma(w)_i = w_{i+1}$ if $w = (i_0, i_1, \dots)$. The basic geometric construction is considered first. Starting from p arbitrary closed subsets $\Delta_1, \dots, \Delta_p$ of R^m , a Cantor-like set is defined

by

$$(3.1) \quad F = \bigcap_{n=0}^{\infty} \bigcup_{\substack{(i_0, \dots, i_n) \\ Q\text{-admissible}}} \Delta_{i_0, \dots, i_n},$$

where the basic sets on the n -th step of the geometric construction, Δ_{i_0, \dots, i_n} , $i_k = 1, \dots, p$ ($n \geq 0$) are closed, and $Q \subset \sum_p^+$ is compact set invariant under the shift map σ .

In 1996 [PW], Pesin and Weiss established the following geometric construction whose basic sets satisfy the following conditions:

- (i) $\Delta_{i_0, \dots, i_n j} \subset \Delta_{i_0, \dots, i_n}$ for $j = 1, \dots, p$;
- (ii) $\underline{B}_{i_0, \dots, i_n} \subset \Delta_{i_0, \dots, i_n} \subset \bar{B}_{i_0, \dots, i_n}$, where $\underline{B}_{i_0, \dots, i_n}$ and $\bar{B}_{i_0, \dots, i_n}$ are closed balls of radii $\underline{r}_{i_0, \dots, i_n}$ and $\bar{r}_{i_0, \dots, i_n}$;
- (iii) $\underline{r}_{i_0, \dots, i_n} = K_1 \prod_{j=0}^n \lambda_{i_j}$, $\bar{r}_{i_0, \dots, i_n} = K_2 \prod_{j=0}^n \lambda_{i_j}$, where $\lambda_{i_j} \in \{\lambda_1, \dots, \lambda_k\}$ with $0 < \lambda_k < 1$ for $k = 1, \dots, p$, $K_1 > 0$, and $K_2 > 0$ are constants;
- (iv) $\text{int } B_{i_0, \dots, i_n} \cap \text{int } B_{j_0, \dots, j_m} = \emptyset$ for any $(i_0, \dots, i_n) \neq (j_0, \dots, j_m)$ and $m \geq n$.

Let (Q, σ) be the dynamical systems generated by the above construction, where $Q \subset \sum_p^+$ is a compact shift invariant set and σ is shift map. Pesin and Weiss were able to prove following results for F and (Q, σ) , [PW] and [P].

Theorem 4. *Let F be the limit set for a geometric construction ((i)-(iv)) modeled by a symbolic dynamical system (Q, σ) . Then*

- (1) $\underline{\dim}_B F = \overline{\dim}_B F = \dim_H F = s_\lambda$ where s_λ is such that

$$P_Q(s_\lambda \log \lambda_{i_0}) = 0,$$

P_Q denote the topological pressure on Q with respect to the shift map σ

(2) $s_\lambda = -\frac{h_{\mu_\lambda}(\sigma|_Q)}{\int_Q \log \lambda_{i_0} d\mu_\lambda}$ where μ_λ be an equilibrium measure for the function $(i_0, \dots, i_n) \mapsto s_\lambda \log \lambda_{i_0}$ on Q (Ref Theorem 13.1 of [P], p125).

Theorem 5. *Let F be the limit set for a geometric construction ((i)-(iv)) modeled by a symbolic dynamical system (Q, σ) . Then*

(1) $s_\lambda \leq \frac{h_Q(\sigma)}{-\log \lambda_{\max}}$, where $\lambda_{\max} = \max\{\lambda_k : 1 \leq k \leq p\}$ and $h_Q(\sigma)$ is the topological entropy of σ on Q ; equality occurs if $\lambda_i = \lambda$ for $i = 1, \dots, p$; in particular, if $h_Q(\sigma) = 0$, then

$$\underline{\dim}_B F = \overline{\dim}_B F = \dim_H F = 0;$$

(2) if $\lambda_i = \lambda$ for $i = 1, \dots, p$, then

$$\underline{\dim}_B F = \overline{\dim}_B F = \dim_H F = s_\lambda = \frac{h_Q(\sigma)}{-\log \lambda}.$$

(Ref Theorem 13.2 of [P], p129)

4. PROOF OF THE MAIN THEOREM

This section presents detail proofs of the Theorem A and Theorem B . f is assumed throughout to be a unimodal map with positive topological entropy.

We first prove (iv) of Theorem A .

Lemma 6. *If f is a tent map with slope $\lambda > 1$, let F_t be the gap maps induced by f , then the measure-theoretical entropy $h_{\mu_\lambda}(t)$ with respect to the equilibrium measure defined in Theorem 4 is monotonic, continuous and the constant part of $h_{\mu_\lambda}(t)$ is open and dense in parameter space; that is, $\{s \mid h_{\mu_\lambda}(t) = \text{constant}\}$ is open and dense in $[0, 1]$.*

Proof. The geometric construction of Pesin and Weiss can be easily demonstrated as follows.

(1) Take $p = 2$, and let $\Delta_1 = C_{1,1} = C_{1,1}(t)$ be the left part and $\Delta_2 = C_{1,2} = C_{1,2}(t)$ be the right part of $[0, 1] \setminus U_t$, where $C_{m,i}$ are the elements of C_m . Now $\Delta_{i_0 \dots i_{n-1}j} \in C_{n+1}$ is such that $f(\Delta_{i_0 \dots i_{n-1}j}) = \Delta_{i_1 \dots i_{n-1}j}$, where $\Delta_{i_1 \dots i_{n-1}j}$ is some element of C_n ,

(2) Conditions (ii) and (iv) clearly hold for C_m and C_{m+1} ,

(3) Condition (iii) holds by showing $K_1 = \text{diam}C_{1,1} = \text{diam}\Delta_1$ or $K_2 = \text{diam}C_{1,2} = \text{diam}\Delta_2$ and $\lambda_k = \lambda$, for $k = 1, 2$.

Thus for such a measure μ_λ defined in Theorem 4, and by applying Theorem 5 we have

$$s_\lambda = -\frac{h_{\mu_\lambda}(\sigma | Q)}{\int_Q \log \lambda_{i_0} d\mu_\lambda} = -\frac{h_{\mu_\lambda}(\sigma | Q)}{\log \lambda}.$$

Since $\lambda_k = \lambda$, for $k = 1, 2$,

$$\int_Q \log \lambda_{i_0} d\mu_\lambda = \log \lambda.$$

Therefore we obtain

$$h_{\mu_\lambda}(t) = h_{\mu_\lambda}(\sigma | Q) = h_Q(\sigma) = h_{\text{top}}(t).$$

The result follow by applying Theorem 3 to $h_{\text{top}}(t)$. The proof is complete. \square

Next, (iii) of Theorem A is proven.

Lemma 7. *If f is a tent map with slope $\lambda > 1$, then the lower and upper box dimension and Hausdorff dimension of the gap map F_t form devil's staircase functions with respect to t . Furthermore, these three dimensions are equal.*

Proof. By Theorem 5 and the proof of Lemma 6, we have

$$\underline{\dim}_B C(t) = \overline{\dim}_B C(t) = \dim_H C(t) = \frac{h_Q(\sigma)}{-\log \lambda} = \frac{h_{\text{top}}(F_t)}{-\log \lambda},$$

and from the definition of F_t , λ is independent of t . The results follow. \square

Next, we prove (i) of Theorem A.

Lemma 8. (*Monotonicity*) *If $s < t$, $\underline{\dim}_B(s) \leq \underline{\dim}_B(t)$, $\overline{\dim}_B(s) \leq \overline{\dim}_B(t)$ and $\dim_H(s) \leq \dim_H(t)$.*

Proof. We only prove the lemma for $\underline{\dim}_B(t)$, $\overline{\dim}_B(t)$ and $\dim_H(t)$ can be treated analogously. If $s < t$, $C(s) \subseteq C(t)$ is easily checked. If κ is a cover of $C(t)$ with $\text{diam}\kappa = \varepsilon$, then κ is also a cover of $C(s)$, and a subcover $\hat{\kappa}$ of κ can be chosen that, covers $C(s)$; with $\text{card}(\hat{\kappa}) \leq \text{card}(\kappa)$. Hence $\underline{\dim}_B C(s) \leq \underline{\dim}_B C(t)$, and the proof is completed. \square

To prove (ii) of Theorem A, we need to introduce the kneading sequence $K(t)$ of $t \in [0, 1]$ with respect to F_t .

Definition 8. *Let f be a unimodal map with turning point c .*

(i) *The itinerary of x with respect to f , defined by $I(x)$ is the sequence $i = (i_0(x), i_1(x), \dots, i_n(x), \dots)$ where*

$$i_j(x) = \begin{cases} 0, & \text{if } f^j(x) < c, \\ 1, & \text{if } f^j(x) > c, \\ c, & \text{if } f^j(x) = c. \end{cases} ,$$

(ii) *A signed lexicographic ordering \prec on $\Sigma = \{0, c, 1\}^N$ is defined as follows. Let $s_i = t_i$ for $i = 1$ to $n - 1$, then $s \prec t$ if either*

(a) $\tau_{n-1}(s)$ is even and $s_n < t_n$

or

(b) $\tau_{n-1}(s)$ is odd and $s_n > t_n$,

where $\tau_k(s) = \sum_{i=0}^k s_i$. We also write $s \preceq t$ if $s \prec t$ or $s = t$.

(iii) The kneading sequence of f , denoted by Kf , is defined as follows,

(a) If c is aperiodic point,

$$Kf = I(f(c)),$$

(b) If c is periodic point of f with period n ,

$$Kf = \lim_{x \rightarrow c^+} I(f(x)),$$

Moreover, $K(t) = KF_t \equiv I(f(t))$, and $K(t)$ is called even(odd) if $\tau_{n-1}(K(t))$ is even(odd).

We need some lemma from [BHL].

Lemma 9. *If f is a continuous unimodal map, and F_t is induced by f and $t \in [0, 1]$ is a periodic point of f with period n ; that is, $f^n(t) = t$, then there exists a neighborhood V_t of t in $[0, 1]$ such that if $K(t)$ is even(odd), then t is the left (right) end point of V_t such that for all $s \in V_t$, $C(s) = C(t)$. Furthermore,*

$$h_{top}(s) = h_{top}(t), \forall s \in V_t.$$

Proof. Ref [BHL]. □

Lemma 10. *If $h(f) > 0$, then the union of the constant part of $\underline{\dim}_B(t)$ ($\overline{\dim}_B(t)$ and $\dim_H(t)$) of the gap maps is dense in parameter space.*

Proof. The lemma is proven by contradiction. Suppose there exist open interval K of $[0, 1]$ such that for all s, t in K , and $s < t$, $\underline{\dim}_B(s) < \underline{\dim}_B(t)$.

First, we may assume $0 < h_{top}(s) \leq h_{top}(t)$ for some s, t in K , and $s < t$, thus there exist a periodic orbit $p \in (s, t)$ with period n . We assume $K(p)$ is even(odd), then there exists a neighborhood $V_p \subset K$ of p from Lemma 9 with p is the left (right) end point of V_p such that for

all $q \in V_p$, $C(q) = C(p)$, which implies $\underline{\dim}_B(q) = \underline{\dim}_B(p)$, this leads a contradiction. $\overline{\dim}_B(t)$ and $\dim_H(t)$ can be treated analogously. \square

Combined with results of Lemmas 6, 7, 8, 9 and 10, the proof Theorem A is complete.

We first prove the first part of Theorem B.

Lemma 11. *If V is a maximal interval of the constant part of $h_{top}(t)$, then V is a maximal constant part in $\underline{\dim}_B(s)$, $\overline{\dim}_B(t)$ and $\dim_H(t)$.*

Proof. Let $t \in [0, 1]$, and V_t be such that $h_{top}(t)$ is constant in V_t . V_t is assumed to be maximal; that is $\exists \hat{t} \in [0, 1]$ such that \hat{t} is a periodic point of period n , and if $K(\hat{t})$ is even(odd), then \hat{t} is the left (right) end point of V_t by Lemma 9, thus we have

$$C(s) = C(\hat{t}), \forall s \in V_t.$$

It implies $\underline{\dim}_B(s) = \underline{\dim}_B(\hat{t}), \forall s \in V_t$. Similar arguments can apply to $\overline{\dim}_B(t)$ and $\dim_H(t)$, the proof is complete. \square

To prove the second part of Theorem B, we need following lemmas.

Lemma 12. *If f is a unimodal map, and F_t are the family of gap maps induced by f , we have*

$$\lim_{n \rightarrow \infty} \frac{\log N_n(t)}{n} = h_{top}(t)$$

where $N_n(t)$ is the cardinality of the maximal interval of $C_n(t)$.

Proof. Ref [M1] \square

Lemma 13. *If f is a unimodal map, and F_t are the family of gap maps induced by f , we have*

$$(4.1) \quad \lim_{n \rightarrow \infty} \frac{\log \bar{\varepsilon}_n^{-1}}{n} \geq \lim_{n \rightarrow \infty} \log |DF^n(t)|^{\frac{1}{n}}$$

for all $t \in C$.

Furthermore, if f is a tent map with constant slope $\lambda > 1$, then we have

$$(4.2) \quad \lim_{n \rightarrow \infty} \frac{\log \bar{\varepsilon}_n^{-1}}{n} = \log \lambda$$

Proof. By the definition of $C_n(t)$, $F_t|_{C_n(t)}$ is a homeomorphism; Then, by the mean value theorem, for each $A \in C_n(t)$, there exists $\bar{t} \in A$ such that

$$\mu(F_t^n(A)) = |DF_t^n(\bar{t})| \mu(A),$$

where μ is Lebesgue measure of $[0, 1]$.

Let $k_1 = \min\{\mu(C_{1,1}), \mu(C_{1,2})\}$ and $k_2 = \max\{\mu(C_{1,1}), \mu(C_{1,2})\}$, thus we have

$$k_1 \leq \mu(F_t^n(A)) \leq k_2.$$

By the definition, $0 < \bar{\varepsilon}_n \leq \mu(A)$ for all $A \in C_n$, this implies

$$\frac{\log \bar{\varepsilon}_n^{-1}}{n} \geq \frac{\log |DF_s^n(\bar{t})| - \log k_1}{n}.$$

Since k_1 is constant independent of n and $\bigcap_{n \geq 1} C_n = C$. Taking $n \rightarrow \infty$, then (4.1) holds.

If f is tent map with slope $\lambda > 1$, then $\mu(A) = k_1 \lambda^{n-1}$ or $k_2 \lambda^{n-1}$ for all $A \subset C_n(t)$. Hence, for all $A, B \in C_n(t)$,

$$\frac{\mu(A)}{\mu(B)} \leq \frac{k_2}{k_1} = k_3$$

where k_3 is a constant independent the choice of A, B and n . Hence,

$$k_3 \mu(A) \geq \bar{\varepsilon}_n \geq \mu(A).$$

This implies

$$\frac{\log k_3 + \log |DF_s^n(\bar{t})| - \log k_2}{n} \geq \frac{\log \bar{\varepsilon}_n^{-1}}{n} \geq \frac{\log |DF_s^n(\bar{t})| - \log k_1}{n}.$$

Since k_1, k_2 and k_3 are constants independent of the subcover of C_n , and $\frac{\log|DF_s^n(\bar{t})|}{n} = \lambda$. Letting $n \rightarrow \infty$, then (4.2) follows. The proof is complete. \square

Lemma 14. *If f is a unimodal map, and F_t are the family of gap maps induced by f , then*

$$\liminf_{n \rightarrow \infty} \frac{\log N_n(t)}{\log \hat{\varepsilon}_n^{-1}(t)} \geq \underline{\dim}_B(t) (\overline{\dim}_B(t), \dim_H(t)).$$

Furthermore, if f is a tent map with constant slope $\lambda > 1$, then we have

$$\liminf_{n \rightarrow \infty} \frac{\log N_n(t)}{\log \hat{\varepsilon}_n^{-1}(t)} = \underline{\dim}_B(t) = \overline{\dim}_B(t) = \dim_H(t).$$

Proof. Since C_n is a cover of C with $\text{diam}C_n = \hat{\varepsilon}_n$, thus we have $N_n \geq N_C(\hat{\varepsilon}_n)$, which implies

$$\begin{aligned} \underline{\dim}_B(t) &= \liminf_{\varepsilon \rightarrow 0} \frac{\log N_C(\varepsilon)}{\log \varepsilon^{-1}} \\ &= \liminf_{n \rightarrow \infty} \frac{\log N_C(\hat{\varepsilon}_n^{-1})}{\log \hat{\varepsilon}_n^{-1}} \\ &\leq \liminf_{n \rightarrow \infty} \frac{\log N_n(t)}{\log \hat{\varepsilon}_n^{-1}}. \end{aligned}$$

If f is tent map with slope $\lambda > 1$, as in Lemma 13, we have

$$\frac{\hat{\varepsilon}_n}{\bar{\varepsilon}_n} \leq k_3.$$

Thus we have

$$N_C(k_3 \bar{\varepsilon}_n) \geq N_n \geq N_C(\hat{\varepsilon}_n),$$

which implies

$$\frac{\log N_C(k_3 \bar{\varepsilon}_n)}{\log(\hat{\varepsilon}_n)^{-1}} \geq \frac{\log N_n}{\log(\hat{\varepsilon}_n)^{-1}} \geq \frac{\log N_C(\hat{\varepsilon}_n)}{\log(\hat{\varepsilon}_n)^{-1}}.$$

Since k_3 is independent of n , letting $n \rightarrow \infty$, implies

$$\underline{\dim}_B(t) \geq \liminf_{n \rightarrow \infty} \frac{\log N_n}{\log(\hat{\varepsilon}_n)^{-1}} \geq \underline{\dim}_B(t),$$

the result follows. The proofs of Hausdoff and upper box dimensions are similar. \square

To complete the proof of Theorem B. Given $\varepsilon_1 > 0$, choose $n_1 \in Z^+$ such that

$$\frac{\log N_n(t)}{\log \hat{\varepsilon}_n^{-1}(t)} \geq \underline{\dim}_B(t) + \varepsilon_1$$

whenever $n \geq n_1$. Given $\varepsilon_2 > 0$, choose $n_2 \in Z^+$ such that

$$\frac{\log N_n}{n} \leq h_{top}(F_s) - \varepsilon_2$$

whenever $n \geq n_2$. Given $\varepsilon_3 > 0$, choose $n_3 \in Z^+$ such that

$$\frac{\log \bar{\varepsilon}_n^{-1}}{n} \geq \log |DF^n(t)|^{\frac{1}{n}} + \varepsilon_3$$

whenever $n \geq n_3$. Let $N = \max\{n_1, n_2, n_3\}$, as $n \geq N$,

$$\begin{aligned} \underline{\dim}_B(t) + \varepsilon_1 &\leq \frac{\log N_n(t)}{\log \hat{\varepsilon}_n^{-1}(t)} \\ &= \left(\frac{\log N_n(t)}{n}\right) \left(\frac{n}{\log \bar{\varepsilon}_n^{-1}}\right) \left(\frac{\log \bar{\varepsilon}_n^{-1}}{\log \hat{\varepsilon}_n^{-1}}\right) \\ &\leq (h_{top}(F_t) - \varepsilon_2) (\log |DF^n(t)|^{\frac{1}{n}} + \varepsilon_3)^{-1} \left(\frac{\log \bar{\varepsilon}_n^{-1}}{\log \hat{\varepsilon}_n^{-1}}\right). \end{aligned}$$

Since $\varepsilon_1, \varepsilon_2$ and ε_3 are arbitrary, now letting $n \rightarrow \infty$ yields

$$\lim_{n \rightarrow \infty} \left(\frac{\log \bar{\varepsilon}_n^{-1}}{\log \hat{\varepsilon}_n^{-1}}\right) \left(\frac{h_{top}(F_t)}{\lim_{n \rightarrow \infty} \log |DF^n(t)|^{\frac{1}{n}}}\right) \geq \underline{\dim}_B(t).$$

If f is tent map with slope $\lambda > 1$, since

$$\bar{\varepsilon}_n \leq \hat{\varepsilon}_n \leq k_3 \bar{\varepsilon}_n$$

where k_3 is defined in the proof of Lemma 13, then equality in Lemma 14 can be replaced by

$$\liminf_{n \rightarrow \infty} \frac{\log N_n(t)}{\log \bar{\varepsilon}_n^{-1}(t)} = \underline{\dim}_B(t).$$

From Lemmas 12, 13 and 14, we have

$$\underline{\dim}_B(t) = \frac{h_{top}(F_t)}{\log \lambda}.$$

Then the proof of Theorem B is complete.

Remark 1. *Some relationship exists between $h_{top}(f)$, Hausdoff dimension, and $lip(f)$, where $lip(f)$ is the lipschitz constant of f , says*

$$\underline{\dim}_H(f) \leq \frac{h_{top}(f)}{\log lip(f)}$$

, see [M2].

REFERENCES

- [BHL] J.-C Ban, C.-H. Hsu and S.-S Lin, Devil's Staircase of Gap Maps, *Internat. J. Bifur. Chaos. Appl. Sci. Engrg.* 13 (2003), 115-122.
- [M1] M.-Misiurewicz, Entropy of Maps with Horizontal Gaps, *preprint*.
- [M2] M.-Misiurewicz, On Bowen's Definition of Topological Entropy, *preprint*.
- [ALM] L. Alsedà, J. Llibre and M. Misiurewicz, Combinatorial Dynamics and Entropy in Dimension One, *Advanced Series in Nonlinear Dynamics, vol. 5, World Scientific, Singapore* (2000), second Edition.
- [D] R.L. Devaney, An Introduction to Chaotic Dynamical Systems, *Addison Wesley, Menlo Park* (1986).
- [ZB] K. Zyczkowski and E.M. Bollt, On the Entropy Devil's Staircase in a Family of Gap-Tent Maps, *Physica D* 132 (1999), 392-410.
- [P] Y.-Pesin, Dimension Theory in Dynamical Systems: contemporary views and applications, *the university of chicago press* (1997).
- [PW] Ya.Pesin, H. Weiss, On the Dimension of Deterministic and Random Cantor-like Sets, Symbolic Dynamics and the Eckmann-Ruelle Conjecture. *Comm. Math. Phys.* 182 vol1 (1996), 105-153.

- [Mo] P.-Moran, Additive Functions of Intervals and Hausdoff Dimension, *Math. Proc. Comb. Phil. Soc.*42 (1946),15-23.

National Center for Theoretical Sciences, National Tsing Hua University, Hsinchu, Taiwan 30043

Department of Applied Mathematics, National Chiao-Tung University, Hsinchu, Taiwan

E-mail address: jcban@math.cts.nthu.edu.tw, sslin@math.nctu.edu.tw