

成果報告 (執行時段： 89 年 8 月 1 日至 90 年 7 月 31 日)

白色念珠菌之致病基因的鑑定

Identification of Virulence Genes of *Candida albicans*

計畫編號：NSC 89-2320-B-009-002

執行期限： 89 年 8 月 1 日至 90 年 7 月 31 日

主持人：楊昶良 國立交通大學生物科技學系

計畫參與人員：

協同主持人 羅秀容 國家衛生研究院臨床研究組

學生 邱美惠 國立交通大學生物科技學系博士班

徐貝帆 國立交通大學生物科技學系碩士班(89年8月-90年1月)

石欣怡 國立交通大學生物科技學系碩士班

林啟陽 國立交通大學生物科技學系碩士班

羅瀚倫 國立交通大學生物科技學系碩士班(90年2月-90年7月)

## 一、中文摘要

黴菌感染在近十年已為院內感染的主要病原菌之一。黴菌不但造成局部性的表皮感染，還會引起散播性的全身感染。甚至還會造成生命的威脅。目前所用的抗黴菌藥物都有副作用、不能有效控制一些黴菌及引起抗藥性的致病原的問題。

白色念珠菌(*Candida albicans*)是最常引起感染的黴菌之一。它可由酵母菌型細胞轉換為菌絲型細胞，這種形態的交替和其致病力有關。不會形成菌絲型細胞的 *cph1/cph1 efg1/efg1* 雙重突變體(double mutant)也沒有致病力。因此，由 Cph1p 及 Efg1p 所調控的基因或其產物和白色念珠菌的致病力是息息相關的。

這個研究就是為尋找白色念珠菌的致病基因(virulence genes)所設計的。前題是：野生種的白色念珠菌(wild-type)有致病力，而 *cph1/cph1 efg1/efg1* 雙重突變體沒有。推論是：若基因只在野生種表現而不在 *cph1/cph1 efg1/efg1* 突變體表現，這些基因可能和白色念珠菌的致病有關；相反地，若基因只在突變體表現，這些基因或其產物可能有阻止白色念珠菌感染的功能。

利用相減雜交法(subtractive hybridization)的技術，相互比對野生種及雙重突變體的 total RNA，我們成功的分離了 991 個只在野生種表現或在野生種表現比在雙重突變體多(Wt - Mutant)之 cDNA clones 及 340 個只在雙重突變體表現或在雙重突變體表現比在野生種多 (Mutant-Wt)之 cDNA clones。從只在野生種表現或在

野生種表現比在雙重突變體多之 cDNA clones 中，我們任意挑選了 11 個 cDNA clones 做一次序列檢定(Sequencing) 反應。其中有 5 個是 *EFG1* 的片段(表一)。因為在 *cph1/cph1 efg1/efg1* 雙重突變體，*EFG1* 已經被突變。因此，在此實驗中能分離出 *EFG1* 的片段，證明此技術之可行性。接著利用 restriction fragment mapping, 將 Wt-Mutant 組分成 381 類，Mutant-Wt 組 266 類。將所得分類依其所含 cDNA clone 之數目排序，排序由多至少。再依序每類取 2 個 cDNA clones 做 DNA 定序。Wt-Mutant 中已知有 14 類是不同片段之 *EFG1* cDNA，其他 16 類則為不同之基因。

為了更清楚地瞭解這些基因的作用，我們會將和白色念珠菌致病可能相關的基因突變，再觀察這些突變體的特性。

這個研究所得之基因及其特性不但可以幫助我們對白色念珠菌的致病機制有進一步的了解，而且對抗黴菌新藥物的發展亦有貢獻。

**關鍵詞：致病基因，念珠菌，抗黴菌新藥物**

English Abstract:

In the past decade, yeast infections have played an increasingly important role in nosocomial infections in the hospital setting. *Candida albicans* is the most frequently isolated fungal pathogen in humans [4].

*Candida albicans* can switch from a unicellular yeast form into a multicellular filamentous form. It has been reported that the *cph1/cph1 efg1/efg1* double mutant in *C. albicans* is defective in filamentous growth and is also avirulent in a mouse model [5]. Thus, identifying the genes regulated by the Cph1p and Efg1p in *C. albicans* may reveal important therapeutic targets in addition to the mechanism of pathogenicity.

This study is designed to identify the virulence genes in *C. albicans*. We would like to compare the patterns of gene expression between the wild-type strain and the

*cph1/cph1 efg1/efg1* double mutant in *C. albicans* by the method of subtractive hybridization [3]. The genes expressed in the wild-type strain but less or not in the *cph1/cph1 efg1/efg1* double mutant (Wt-Mutant) may be virulence genes which are potential candidates for targets of antifungal agents. In contrast, the genes expressed in the *cph1/cph1 efg1/efg1* double mutant but less or not in the wild-type strain (Mutant-Wt) may be negative regulators for pathogenicity of *C. albicans*. In the experiment from which the levels of gene expression of the double mutant was subtracted from that of the wild-type strain, we have obtained 991 clones, while only 340 clones from the reversed subtraction (Mutant-Wt). Eleven clones were picked randomly for sequencing. Five of them are *EFG1* gene of *C. albicans* (Table 1). Since *EFG1* has been mutated in the *cph1/cph1 efg1/efg1* double mutant, identification of the *EFG1* gene in this experiment is an excellent indication and positive control that the techniques used in this research have worked as intended. The rest of the clones were then analyzed by restriction fragment mapping. Wt-Mutant has 381 different groups of restriction fragment pattern, while Mutant-Wt has 266 groups. Two from each group were sent for sequencing analysis. In the Wt-Mutant groups that have been sequenced, 14 different portion of the EFG-1 cDNA fragment and 11 other gene fragments have been identified. The particular functions of those genes expressing specifically in the virulent strain or the avirulent mutant will be determined by gene disruption approaches.

The knowledge gained from this study will help us to understand more about the pathogenicity of *C. albicans* and also to design better antifungal drugs that block the virulence of *C. albicans* and, potentially, other fungal pathogens.

Keywords : Virulence gene, Candida, antifungal agent

## 二、緣由與目的

In the past decade, nosocomial infections caused by yeast pathogens have increased dramatically. For instance, the prevalence of nosocomial candidemia increased 27-fold from 1981 through 1993 at National Taiwan University Hospital [2]. In the United States yeast infections rank as the fourth most common cause of nosocomial bloodstream infection [6]. The rise in the prevalence of fungal infections has exacerbated the need for the next generation anti-fungal agents. Many current available anti-fungal agents have several problems, for example, having side effects, being ineffective against new or reemerging fungi, and leading to the rapid development of resistance. To design and develop a new effective anti-fungal agent requires the understanding of the mechanism of pathogenicity.

To establish an infection, fungal pathogens have to survive and divide in the host environment and also invade new tissue and phagocytic cells. The molecule dissection of virulence genes of fungal pathogens has hence focused on adhesion, proteinase secretion, and hyphal formation. Since *C. albicans* is one of the major fungal pathogens in humans and has been established for the molecular study, this study is designed to identify the virulence genes of *C. albicans*. A strain of *C. albicans* with *cph1/cph1 efg1/efg1* double mutant has been shown to lose its virulence in a mouse model [5]. This strain was used for our study to isolate genes regulated by the Cph1p and Efg1p, which control the virulence in *C. albicans*.

## 三、結果與討論

In order to identify virulence genes of *C. albicans*, we determined the differential gene expression, under the condition that induces wild-type cells into hyphal form, of the virulent wild-type strain and the avirulent *cph1/cph1 efg1/efg1* double mutant by means of the subtractive hybridization [3]. The genes expressed in the wild-type

strain but less or not in the *cph1/cph1 efg1/efg1* double mutant may be virulence genes of *C. albicans*. In contrast, the genes expressed in the *cph1/cph1 efg1/efg1* double mutant but less or not in the wild-type strain may be negative regulators for virulence of *C. albicans*.

The RNA isolations are preceded according to the procedure of Carlson and Botstein [1]. The total RNA from the fungal cells was used for the synthesis of cDNA for the subtractive hybridization described by Diatchenko et.al. [3]. The products from the subtractive hybridization were then subcloned onto plasmid vectors.

In the experiment from which the gene expression pattern of the double mutant was subtracted from that of the wild-type strain, we have obtained 991 clones, while only 340 clones from the reversed subtraction. Eleven clones were picked randomly for sequencing. The raw sequence results were used for sequence alignment by the blast program provided by NCBI. Five of them are *EFG1* gene of *C. albicans* (Fig. 1). Identification of the *EFG1* gene is an excellent indication and positive control that the techniques used in this research have worked as intended since *EFG1* has been deleted in the *cph1/cph1 efg1/efg1* double mutant. Another indication of success is the cloning of the homolog of yeast *STE20* (Table 1). *STE20* is known to be involved in hyphal formation. In the Mutant-Wt groups, identification of Tup1 is the indicator that our experimental has worked accordingly, since Tup1 is known to express higher in the double mutants. The categorization of the cDNA clones was performed by comparing the patterns of restriction enzyme digestion of each clone. The Wt-Mutant has 381 groups, and the Mutant-Wt has 266 groups. Groups with more clones in the groups have been sequenced first. Only two clones from each group were subjected to the sequence process.

So far, in the Wt-Mutant groups, 14 groups are different cDNA fragments of *EFG1*. Besides, 11 different gene fragments have been identified. To understand their

function in *Candida* virulence, those genes will be subjected to genetic and functional studies after assessing by northern analysis.

#### 四、計畫結果自評

The identification of the *EFGI* sequence from the clones obtaining from the experiment in which the gene expression pattern of the *cph1/cph1 efg1/efg1* double mutant was subtracted from that of the wild-type strain provides a solid evidence that the subtractive hybridization has functioned as it was designed to. Another indication of success is the cloning of the homolog of yeast *STE20* (Table 1). *STE20* is known to be involved in hyphal formation. At the same time, five out of eleven clones chosen for sequencing are *EFGI* indicating that this method is also very sensitive to the level of gene expression. It is likely that the level of expression of *EFGI* is higher in the hyphal form than that in the yeast form. However, it is necessary to confirm the expression (both quality and quantity) of interested genes in wild-type and double mutant cells by the Northern blots analysis.

More sequences will become available and thus other genes that involves in the virulence will be identified. However, it is worth while to emphasize that in our experimental design, the hyphal formation is used only as a marker for virulence, genes regulates these two pathways do not have to be complete identical. In fact, it is likely that they only share some genes, for instance, *EFGI* and *CPHI*. It will be necessary to use mutagenesis approach to determine the functions of interested genes.

#### 五、參考文獻

1. Carlson M Botstein D. Two differentially regulated mRNAs with different 5' ends encode secreted with intracellular forms of yeast invertase. Cell 1982; 28:145-154.

2. Chen YC, Chang SC, Sun CC, Yang LS, Hsieh WC, Luh KT. Secular trends in the epidemiology of nosocomial fungal infections at a teaching hospital in Taiwan, 1981 to 1993. *Infect.Control Hosp.Epidemiol.* 1997; 18:369-375.
3. Diatchenko L, Lukyanov S, Lau YF, Siebert PD. Suppression subtractive hybridization: a versatile method for identifying differentially expressed genes. *Methods Enzymol.* 1999; 303:349-80:349-380.
4. Edwards, J. E., Jr. *Candida species.* In *Principles and Practice of Infectious Diseases.* Mandell, G. L., Douglas, R. G., and Bennett, J. E. 1943-1958. 1990. New York: Churchill Livingstone. 淘淘  
Ref Type: Serial (Book,Monograph)
5. Lo HJ, Kohler JR, DiDomenico B, Loebenberg D, Cacciapuoti A, Fink GR. Nonfilamentous *C. albicans* mutants are avirulent. *Cell* 1997; 90:939-949.
6. Pfaller MA, Jones RN, Messer SA, Edmond MB, Wenzel RP. National surveillance of nosocomial blood stream infection due to species of *Candida* other than *Candida albicans*: frequency of occurrence and antifungal susceptibility in the SCOPE Program. SCOPE Participant Group. *Surveillance and Control of Pathogens of Epidemiologic.* *Diagn.Microbiol.Infect.Dis.* 1998; 30:121-129.